

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
29 July 2004 (29.07.2004)

PCT

(10) International Publication Number
WO 2004/063351 A2

(51) International Patent Classification⁷: C12N

(21) International Application Number:
PCT/US2004/000643

(22) International Filing Date: 9 January 2004 (09.01.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/439,498 9 January 2003 (09.01.2003) US
60/456,041 19 March 2003 (19.03.2003) US
60/514,549 23 October 2003 (23.10.2003) US

(71) Applicant (for all designated States except US): MACRO-
GENICS, INC. [US/US]; 1500 East Gude Drive,
Rockville, MD 20850-5307 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): STAVENHAGEN,

Jeffrey [US/US]; 19417 Treadway Road, Brookville, MD
20833 (US). VIJH, Sujata [US/US]; 912-100 Orchard
Ridge Drive, Gaithersburg, MD 20878 (US).

(74) Agents: INSOGNA, Anthony, M. et al.; Jones Day, 222
East 41st Street, New York, NY 10017-6702 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,

[Continued on next page]

(54) Title: IDENTIFICATION AND ENGINEERING OF ANTIBODIES WITH VARIANT Fc REGIONS AND METHODS OF
USING SAME

(57) Abstract: The present invention relates to molecules, particularly polypeptides, more particularly immunoglobulins (e.g., antibodies), comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, which variant Fc region binds FcγRIIIA and/or FcγRIIA with a greater affinity, relative to a comparable molecule comprising the wild-type Fc region. The molecules of the invention are particularly useful in preventing, treating, or ameliorating one or more symptoms associated with a disease, disorder, or infection. The molecules of the invention are particularly useful for the treatment or prevention of a disease or disorder where an enhanced efficacy of effector cell function (e.g., ADCC) mediated by FcγR is desired, e.g., cancer, infectious disease, and in enhancing the therapeutic efficacy of therapeutic antibodies the effect of which is mediated by ADCC.

BEST AVAILABLE COPY

WO 2004/063351 A2



GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**IDENTIFICATION AND ENGINEERING OF ANTIBODIES WITH
VARIANT Fc REGIONS AND METHODS OF USING SAME**

[0001] This application claims priority to U.S. Provisional Application Nos. 60/439,498; 60/456,041; and 60/514,549 filed on January 9, 2003; March 19, 2003, and October 23, 2003 respectively; all of which are incorporated herein by reference in their entireties.

5 **1. FIELD OF THE INVENTION**

[0002] The present invention relates to molecules, particularly polypeptides, more particularly immunoglobulins (*e.g.*, antibodies), comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, which variant Fc region binds Fc γ RIIIA and/or Fc γ RIIA with a greater affinity, relative to a comparable molecule comprising the wild-type Fc region. The molecules of the invention are particularly useful in preventing, treating, or ameliorating one or more symptoms associated with a disease, disorder, or infection. The molecules of the invention are particularly useful for the treatment or prevention of a disease or disorder where an enhanced efficacy of effector cell function (*e.g.*, ADCC) mediated by Fc γ R is desired, *e.g.*, cancer, infectious disease, and in enhancing the therapeutic efficacy of therapeutic antibodies the effect of which is mediated by ADCC.

2. **BACKGROUND OF THE INVENTION**

2.1 **Fc RECEPTORS AND THEIR ROLES IN THE IMMUNE SYSTEM**

[0003] The interaction of antibody-antigen complexes with cells of the immune system results in a wide array of responses, ranging from effector functions such as antibody-dependent cytotoxicity, mast cell degranulation, and phagocytosis to immunomodulatory signals such as regulating lymphocyte proliferation and antibody secretion. All these interactions are initiated through the binding of the Fc domain of antibodies or immune complexes to specialized cell surface receptors on hematopoietic cells. The diversity of cellular responses triggered by antibodies and immune complexes results from the structural heterogeneity of Fc receptors. Fc receptors share structurally related ligand binding domains which presumably mediate intracellular signaling.

[0004] The Fc receptors, members of the immunoglobulin gene superfamily of proteins, are surface glycoproteins that can bind the Fc portion of immunoglobulin molecules. Each member of the family recognizes immunoglobulins of one or more isotypes through a recognition domain on the α chain of the Fc receptor. Fc receptors are defined by their specificity for immunoglobulin subtypes. Fc receptors for IgG are referred

to as Fc γ R, for IgE as Fc ϵ R, and for IgA as Fc α R. Different accessory cells bear Fc receptors for antibodies of different isotype, and the isotype of the antibody determines which accessory cells will be engaged in a given response (reviewed by Ravetch J.V. *et al.* 1991, *Annu. Rev. Immunol.* 9: 457-92; Gerber J.S. *et al.* 2001 *Microbes and Infection*, 3: 131-139; Billadeau D.D. *et al.* 2002, *The Journal of Clinical Investigation*, 2(109): 161-168; Ravetch J.V. *et al.* 2000, *Science*, 290: 84-89; Ravetch J.V. *et al.*, 2001 *Annu. Rev. Immunol.* 19:275-90; Ravetch J.V. 1994, *Cell*, 78(4): 553-60). The different Fc receptors, the cells that express them, and their isotype specificity is summarized in Table 1 (adapted from Immunobiology: The Immune System in Health and Disease, 4th ed. 1999, Elsevier Science Ltd/Garland Publishing, New York).

Fc γ Receptors

[0005] Each member of this family is an integral membrane glycoprotein, possessing extracellular domains related to a C2-set of immunoglobulin-related domains, a single membrane spanning domain and an intracytoplasmic domain of variable length.

15 There are three known Fc γ Rs, designated Fc γ RI(CD64), Fc γ RII(CD32), and Fc γ RIII(CD16). The three receptors are encoded by distinct genes; however, the extensive homology between the three family members suggest they arose from a common progenitor perhaps by gene duplication.

Fc γ RII(CD32)

20 [0006] Fc γ RII proteins are 40KDa integral membrane glycoproteins which bind only the complexed IgG due to a low affinity for monomeric Ig (10^6 M⁻¹). This receptor is the most widely expressed Fc γ R, present on all hematopoietic cells, including monocytes, macrophages, B cells, NK cells, neutrophils, mast cells, and platelets. Fc γ RII has only two immunoglobulin-like regions in its immunoglobulin binding chain and hence a much lower

25 affinity for IgG than Fc γ RI. There are three human Fc γ RII genes (Fc γ RII-A, Fc γ RII-B, Fc γ RII-C), all of which bind IgG in aggregates or immune complexes.

[0007] Distinct differences within the cytoplasmic domains of Fc γ RII-A and Fc γ RII-B create two functionally heterogenous responses to receptor ligation. The fundamental difference is that the A isoform initiates intracellular signaling leading to cell

30 activation such as phagocytosis and respiratory burst, whereas the B isoform initiates inhibitory signals, *e.g.*, inhibiting B-cell activation.

Signaling through FcγRs

[0008] Both activating and inhibitory signals are transduced through the FcγRs following ligation. These diametrically opposing functions result from structural differences among the different receptor isoforms. Two distinct domains within the cytoplasmic signaling domains of the receptor called immunoreceptor tyrosine based activation motifs (ITAMs) or immunoreceptor tyrosine based inhibitory motifs (ITIMs) account for the different responses. The recruitment of different cytoplasmic enzymes to these structures dictates the outcome of the FcγR-mediated cellular responses. ITAM-containing FcγR complexes include FcγRI, FcγRIIA, FcγRIIA, whereas ITIM-containing complexes only include FcγRIIB.

[0009] Human neutrophils express the FcγRIIA gene. FcγRIIA clustering via immune complexes or specific antibody cross-linking serves to aggregate ITAMs along with receptor-associated kinases which facilitate ITAM phosphorylation. ITAM phosphorylation serves as a docking site for Syk kinase, activation of which results in activation of downstream substrates (*e.g.*, PI₃K). Cellular activation leads to release of proinflammatory mediators.

[0010] The FcγRIIB gene is expressed on B lymphocytes; its extracellular domain is 96% identical to FcγRIIA and binds IgG complexes in an indistinguishable manner. The presence of an ITIM in the cytoplasmic domain of FcγRIIB defines this inhibitory subclass of FcγR. Recently the molecular basis of this inhibition was established. When colligated along with an activating FcγR, the ITIM in FcγRIIB becomes phosphorylated and attracts the SH2 domain of the inositol polyphosphate 5'-phosphatase (SHIP), which hydrolyzes phosphoinositol messengers released as a consequence of ITAM-containing FcγR-mediated tyrosine kinase activation; consequently preventing the influx of intracellular Ca⁺⁺. Thus crosslinking of FcγRIIB dampens the activating response to FcγR ligation and inhibits cellular responsiveness. B cell activation, B cell proliferation and antibody secretion is thus aborted.

TABLE 1. Receptors for the Fc Regions of Immunoglobulin Isotypes

Receptor	Fc γ RI (CD64)	Fc γ RII-A (CD32)	Fc γ RII-B2 (CD32)	Fc γ RII-B1 (CD32)	Fc γ RIII (CD16)	Fc ϵ RI	Fc α RI (CD89)
Binding	IgG1 10^8 M^{-1}	IgG1 $2 \times 10^6 \text{ M}^{-1}$	IgG1 $2 \times 10^6 \text{ M}^{-1}$	IgG1 $2 \times 10^6 \text{ M}^{-1}$	IgG1 $5 \times 10^5 \text{ M}^{-1}$	IgE 10^{10} M^{-1}	IgA1, IgA2 10^7 M^{-1}
Cell Type	Macrophages Neutrophils Eosinophils Dendritic cells	Macrophages Neutrophils Eosinophils Dendritic cells Platelets Langerhan cells	Macrophages Neutrophils Eosinophils	B cells Mast cells	NK cells Eosinophil Macrophages Neutrophils Mast Cells	Mast cells Eosinophil Basophils	Macrophages Neutrophils Eosinophils
Effect of Ligation	Uptake Stimulation Activation of respiratory burst Induction of killing	Uptake Granule release	Uptake Inhibition of Stimulation	No uptake Inhibition of Stimulation	Induction of Killing	Secretion of granules	Uptake Induction of killing

2.2 DISEASES OF RELEVANCE

2.2.1 CANCER

[0011] A neoplasm, or tumor, is a neoplastic mass resulting from abnormal uncontrolled cell growth which can be benign or malignant. Benign tumors generally remain localized. Malignant tumors are collectively termed cancers. The term "malignant" generally means that the tumor can invade and destroy neighboring body structures and spread to distant sites to cause death (for review, *see* Robbins and Angell, 1976, *Basic Pathology*, 2d Ed., W.B. Saunders Co., Philadelphia, pp. 68-122). Cancer can arise in many sites of the body and behave differently depending upon its origin. Cancerous cells destroy the part of the body in which they originate and then spread to other part(s) of the body where they start new growth and cause more destruction.

[0012] More than 1.2 million Americans develop cancer each year. Cancer is the second leading cause of death in the United States and if current trends continue, cancer is expected to be the leading cause of the death by the year 2010. Lung and prostate cancer are the top cancer killers for men in the United States. Lung and breast cancer are the top cancer killers for women in the United States. One in two men in the United States will be diagnosed with cancer at some time during his lifetime. One in three women in the United States will be diagnosed with cancer at some time during her lifetime.

[0013] A cure for cancer has yet to be found. Current treatment options, such as surgery, chemotherapy and radiation treatment, are oftentimes either ineffective or present serious side effects.

Cancer Therapy

[0014] Currently, cancer therapy may involve surgery, chemotherapy, hormonal therapy and/or radiation treatment to eradicate neoplastic cells in a patient (*See*, for example, Stockdale, 1998, "Principles of Cancer Patient Management", in Scientific American: Medicine, vol. 3, Rubenstein and Federman, eds., Chapter 12, Section IV). Recently, cancer therapy could also involve biological therapy or immunotherapy. All of these approaches pose significant drawbacks for the patient. Surgery, for example, may be contraindicated due to the health of the patient or may be unacceptable to the patient. Additionally, surgery may not completely remove the neoplastic tissue. Radiation therapy is only effective when the neoplastic tissue exhibits a higher sensitivity to radiation than normal tissue, and radiation therapy can also often elicit serious side effects. Hormonal therapy is rarely given as a single agent and although can be effective, is often used to

prevent or delay recurrence of cancer after other treatments have removed the majority of the cancer cells. Biological therapies/immunotherapies are limited in number and may produce side effects such as rashes or swellings, flu-like symptoms, including fever, chills and fatigue, digestive tract problems or allergic reactions.

5 [0015] With respect to chemotherapy, there are a variety of chemotherapeutic agents available for treatment of cancer. A significant majority of cancer chemotherapeutics act by inhibiting DNA synthesis, either directly, or indirectly by inhibiting the biosynthesis of the deoxyribonucleotide triphosphate precursors, to prevent DNA replication and concomitant cell division (*See, for example, Gilman et al., Goodman and Gilman's: The*
10 *Pharmacological Basis of Therapeutics*, Eighth Ed. (Pergamom Press, New York, 1990)). These agents, which include alkylating agents, such as nitrosourea, anti-metabolites, such as methotrexate and hydroxyurea, and other agents, such as etoposides, campathecins, bleomycin, doxorubicin, daunorubicin, *etc.*, although not necessarily cell cycle specific, kill
15 cells during S phase because of their effect on DNA replication. Other agents, specifically colchicine and the vinca alkaloids, such as vinblastine and vincristine, interfere with microtubule assembly resulting in mitotic arrest. Chemotherapy protocols generally involve administration of a combination of chemotherapeutic agents to increase the efficacy of treatment.

[0016] Despite the availability of a variety of chemotherapeutic agents,
20 chemotherapy has many drawbacks (*See, for example, Stockdale, 1998, "Principles Of Cancer Patient Management" in Scientific American Medicine*, vol. 3, Rubenstein and Federman, eds., ch. 12, sect. 10). Almost all chemotherapeutic agents are toxic, and chemotherapy causes significant, and often dangerous, side effects, including severe nausea, bone marrow depression, immunosuppression, *etc.* Additionally, even with administration
25 of combinations of chemotherapeutic agents, many tumor cells are resistant or develop resistance to the chemotherapeutic agents. In fact, those cells resistant to the particular chemotherapeutic agents used in the treatment protocol often prove to be resistant to other drugs, even those agents that act by mechanisms different from the mechanisms of action of the drugs used in the specific treatment; this phenomenon is termed pleiotropic drug or
30 multidrug resistance. Thus, because of drug resistance, many cancers prove refractory to standard chemotherapeutic treatment protocols.

[0017] There is a significant need for alternative cancer treatments, particularly for treatment of cancer that has proved refractory to standard cancer treatments, such as surgery, radiation therapy, chemotherapy, and hormonal therapy. A promising alternative is
35 immunotherapy, in which cancer cells are specifically targeted by cancer antigen-specific

antibodies. Major efforts have been directed at harnessing the specificity of the immune response, for example, hybridoma technology has enabled the development of tumor selective monoclonal antibodies (*See Green M.C. et al., 2000 Cancer Treat Rev., 26: 269-286; Weiner LM, 1999 Semin Oncol. 26(suppl. 14):43-51*), and in the past few years, the Food and Drug Administration has approved the first MABs for cancer therapy: Rituxin (anti-CD20) for non-Hodgkin's Lymphoma and Herceptin [anti-(c-erb-2/HER-2)] for metastatic breast cancer (Suzanne A. Eccles, 2001, *Breast Cancer Res.*, 3: 86-90). However, the potency of antibody effector function, *e.g.*, to mediate antibody dependent cellular cytotoxicity ("ADCC") is an obstacle to such treatment. Methods to improve the efficacy of such immunotherapy are thus needed.

2.2.2 INFLAMMATORY DISEASES AND AUTOIMMUNE DISEASES

[0018] Inflammation is a process by which the body's white blood cells and chemicals protect our bodies from infection by foreign substances, such as bacteria and viruses. It is usually characterized by pain, swelling, warmth and redness of the affected area. Chemicals known as cytokines and prostaglandins control this process, and are released in an ordered and self-limiting cascade into the blood or affected tissues. This release of chemicals increases the blood flow to the area of injury or infection, and may result in the redness and warmth. Some of the chemicals cause a leak of fluid into the tissues, resulting in swelling. This protective process may stimulate nerves and cause pain. These changes, when occurring for a limited period in the relevant area, work to the benefit of the body.

[0019] In autoimmune and/or inflammatory disorders, the immune system triggers an inflammatory response when there are no foreign substances to fight and the body's normally protective immune system causes damage to its own tissues by mistakenly attacking self. There are many different autoimmune disorders which affect the body in different ways. For example, the brain is affected in individuals with multiple sclerosis, the gut is affected in individuals with Crohn's disease, and the synovium, bone and cartilage of various joints are affected in individuals with rheumatoid arthritis. As autoimmune disorders progress destruction of one or more types of body tissues, abnormal growth of an organ, or changes in organ function may result. The autoimmune disorder may affect only one organ or tissue type or may affect multiple organs and tissues. Organs and tissues commonly affected by autoimmune disorders include red blood cells, blood vessels, connective tissues, endocrine glands (*e.g.*, the thyroid or pancreas), muscles, joints, and

skin. Examples of autoimmune disorders include, but are not limited to, Hashimoto's thyroiditis, pernicious anemia, Addison's disease, type 1 diabetes, rheumatoid arthritis, systemic lupus erythematosus, dermatomyositis, Sjogren's syndrome, dermatomyositis, lupus erythematosus, multiple sclerosis, autoimmune inner ear disease myasthenia gravis, Reiter's syndrome, Graves disease, autoimmune hepatitis, familial adenomatous polyposis and ulcerative colitis.

[0020] Rheumatoid arthritis (RA) and juvenile rheumatoid arthritis are types of inflammatory arthritis. Arthritis is a general term that describes inflammation in joints. Some, but not all, types of arthritis are the result of misdirected inflammation. Besides rheumatoid arthritis, other types of arthritis associated with inflammation include the following: psoriatic arthritis, Reiter's syndrome, ankylosing spondylitis arthritis, and gouty arthritis. Rheumatoid arthritis is a type of chronic arthritis that occurs in joints on both sides of the body (such as both hands, wrists or knees). This symmetry helps distinguish rheumatoid arthritis from other types of arthritis. In addition to affecting the joints, rheumatoid arthritis may occasionally affect the skin, eyes, lungs, heart, blood or nerves.

[0021] Rheumatoid arthritis affects about 1% of the world's population and is potentially disabling. There are approximately 2.9 million incidences of rheumatoid arthritis in the United States. Two to three times more women are affected than men. The typical age that rheumatoid arthritis occurs is between 25 and 50. Juvenile rheumatoid arthritis affects 71,000 young Americans (aged eighteen and under), affecting six times as many girls as boys.

[0022] Rheumatoid arthritis is an autoimmune disorder where the body's immune system improperly identifies the synovial membranes that secrete the lubricating fluid in the joints as foreign. Inflammation results, and the cartilage and tissues in and around the joints are damaged or destroyed. In severe cases, this inflammation extends to other joint tissues and surrounding cartilage, where it may erode or destroy bone and cartilage and lead to joint deformities. The body replaces damaged tissue with scar tissue, causing the normal spaces within the joints to become narrow and the bones to fuse together. Rheumatoid arthritis creates stiffness, swelling, fatigue, anemia, weight loss, fever, and often, crippling pain. Some common symptoms of rheumatoid arthritis include joint stiffness upon awakening that lasts an hour or longer; swelling in a specific finger or wrist joints; swelling in the soft tissue around the joints; and swelling on both sides of the joint. Swelling can occur with or without pain, and can worsen progressively or remain the same for years before progressing.

[0023] The diagnosis of rheumatoid arthritis is based on a combination of factors, including: the specific location and symmetry of painful joints, the presence of joint

stiffness in the morning, the presence of bumps and nodules under the skin (rheumatoid nodules), results of X-ray tests that suggest rheumatoid arthritis, and/or positive results of a blood test called the rheumatoid factor. Many, but not all, people with rheumatoid arthritis have the rheumatoid-factor antibody in their blood. The rheumatoid factor may be present
5 in people who do not have rheumatoid arthritis. Other diseases can also cause the rheumatoid factor to be produced in the blood. That is why the diagnosis of rheumatoid arthritis is based on a combination of several factors and not just the presence of the rheumatoid factor in the blood.

[0024] The typical course of the disease is one of persistent but fluctuating joint
10 symptoms, and after about 10 years, 90% of sufferers will show structural damage to bone and cartilage. A small percentage will have a short illness that clears up completely, and another small percentage will have very severe disease with many joint deformities, and occasionally other manifestations of the disease. The inflammatory process causes erosion or destruction of bone and cartilage in the joints. In rheumatoid arthritis, there is an
15 autoimmune cycle of persistent antigen presentation, T-cell stimulation, cytokine secretion, synovial cell activation, and joint destruction. The disease has a major impact on both the individual and society, causing significant pain, impaired function and disability, as well as costing millions of dollars in healthcare expenses and lost wages. (See, for example, the NIH website and the NIAID website).

20 [0025] Currently available therapy for arthritis focuses on reducing inflammation of the joints with anti-inflammatory or immunosuppressive medications. The first line of treatment of any arthritis is usually anti-inflammatories, such as aspirin, ibuprofen and Cox-2 inhibitors such as celecoxib and rofecoxib. "Second line drugs" include gold, methotrexate and steroids. Although these are well-established treatments for arthritis, very
25 few patients remit on these lines of treatment alone. Recent advances in the understanding of the pathogenesis of rheumatoid arthritis have led to the use of methotrexate in combination with antibodies to cytokines or recombinant soluble receptors. For example, recombinant soluble receptors for tumor necrosis factor (TNF)- α have been used in combination with methotrexate in the treatment of arthritis. However, only about 50% of
30 the patients treated with a combination of methotrexate and anti-TNF- α agents such as recombinant soluble receptors for TNF- α show clinically significant improvement. Many patients remain refractory despite treatment. Difficult treatment issues still remain for patients with rheumatoid arthritis. Many current treatments have a high incidence of side effects or cannot completely prevent disease progression. So far, no treatment is ideal, and

there is no cure. Novel therapeutics are needed that more effectively treat rheumatoid arthritis and other autoimmune disorders.

2.2.3 INFECTIOUS DISEASES

[0026] Infectious agents that cause disease fall into five groups: viruses, bacteria, fungi, protozoa, and helminths (worms). The remarkable variety of these pathogens has caused the natural selection of two crucial features of adaptive immunity. First, the advantage of being able to recognize a wide range of different pathogens has driven the development of receptors on B and T cells of equal or greater diversity. Second, the distinct habitats and life cycles of pathogens have to be countered by a range of distinct effector mechanisms. The characteristic features of each pathogen are its mode of transmission, its mechanism of replication, its pathogenesis or the means by which it causes disease, and the response it elicits.

[0027] The record of human suffering and death caused by smallpox, cholera, typhus, dysentery, malaria, etc. establishes the eminence of the infectious diseases. Despite the outstanding successes in control afforded by improved sanitation, immunization, and antimicrobial therapy, the infectious diseases continue to be a common and significant problem of modern medicine. The most common disease of mankind, the common cold, is an infectious disease, as is the feared modern disease AIDS. Some chronic neurological diseases that were thought formerly to be degenerative diseases have proven to be infectious. There is little doubt that the future will continue to reveal the infectious diseases as major medical problems.

[0028] An enormous number of human and animal diseases result from virulent and opportunistic infections from any of the above mentioned infectious agents (*see* Belshe (Ed.) 1984 Textbook of Human Virology, PSG Publishing, Littleton, MA).

25 [0029] One category of infectious diseases are viral infections for example. Viral diseases of a wide array of tissues, including the respiratory tract, CNS, skin, genitourinary tract, eyes, ears, immune system, gastrointestinal tract, and musculoskeletal system, affect a vast number of humans of all ages (*see* Table 328-2 In: Wyngaarden and Smith, 1988, Cecil Textbook of Medicine, 18th Ed., W.B. Saunders Co., Philadelphia, pp.1750-1753).

30 Although considerable effort has been invested in the design of effective anti-viral therapies, viral infections continue to threaten the lives of millions of people worldwide. In general, attempts to develop anti-viral drugs have focused on several stages of viral life cycle (*See e.g.*, Mitsuya *et al.*, 1991, *FASEB J.* 5:2369-2381, discussing HIV). However, a

common drawback associated with using of many current anti-viral drugs is their deleterious side effects, such as toxicity to the host or resistance by certain viral strains.

3. SUMMARY OF THE INVENTION

[0030] The present invention is based, in part, on the identification of mutant human
5 IgG1 heavy chain Fc regions, with altered affinities for Fc γ R receptors (*e.g.*, activating Fc γ Rs, inhibitory Fc γ Rs), using a yeast display system. Accordingly, the invention relates to molecules, preferably polypeptides, and more preferably immunoglobulins (*e.g.*, antibodies), comprising a variant Fc region, having one or more amino acid modifications (*e.g.*, substitutions, but also including insertions or deletions) in one or more regions, which
10 modifications alter, *e.g.*, increase or decrease, the affinity of the variant Fc region for an Fc γ R. Preferably, said one or more amino acid modification increases the affinity of the variant Fc region for Fc γ RIIA and/or Fc γ RIIA. In a preferred embodiment, the molecules of the invention further specifically bind Fc γ RIIB (via the Fc region) with a lower affinity than a comparable molecule (*i.e.*, having the same amino acid sequence as the molecule of
15 the invention except for the one or more amino acid modifications in the Fc region) comprising the wild-type Fc region binds Fc γ RIIB. In some embodiments, the invention encompasses molecules with variant Fc regions, having one or more amino acid modifications, which modifications increase the affinity of the variant Fc region for Fc γ RIIA and/or Fc γ RIIA and enhance the affinity of the variant Fc region for Fc γ RIIB
20 relative to a comparable molecule with a wild type Fc region. In other embodiments, the invention encompasses molecules with variant Fc regions, having one or more amino acid modifications, which modifications increase the affinity of the variant Fc region for Fc γ RIIA and/or Fc γ RIIA but do not alter the affinity of the variant Fc regions for Fc γ RIIB relative to a comparable molecule with a wild type Fc region.

25 [0031] The invention encompasses molecules that are homodimers or heterodimers of Fc regions. Heterodimers comprising Fc regions refer to molecules where the two Fc chains have the same or different sequences. In some embodiments, in the heterodimeric molecules comprising variant Fc regions, each chain has one or more different modifications from the other chain. In other embodiments, in the heterodimeric molecules
30 comprising variant Fc regions, one chain contains the wild-type Fc region and the other chains comprises one or or more modifications. Methods of engineering heterodimeric Fc containing molecules are known in the art and encompassed within the invention.

[0032] In some embodiments, the invention encompasses molecules comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid

modification relative to a wild type Fc region, which variant Fc region does not bind any Fc γ R or binds with a reduced affinity, relative to a comparable molecule comprising the wild-type Fc region, as determined by standard assays (*e.g.*, *in vitro* assays) known to one skilled in the art. In a specific embodiment, the invention encompasses molecules

5 comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild type Fc region, which variant Fc region only binds one Fc γ R, wherein said Fc γ R is Fc γ IIIA. In another specific embodiment, the invention encompasses molecules comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild type Fc region, which

10 variant Fc region only binds one Fc γ R, wherein said Fc γ R is Fc γ RIIA. In yet another embodiment, the invention encompasses molecules comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild type Fc region, which variant Fc region only binds one Fc γ R, wherein said Fc γ R is Fc γ RIIB.

[0033] The affinities and binding properties of the molecules of the invention for an

15 Fc γ R are initially determined using *in vitro* assays (biochemical or immunological based assays) known in the art for determining Fc-Fc γ R interactions, *i.e.*, specific binding of an Fc region to an Fc γ R including but not limited to ELISA assay, surface plasmon resonance assay, immunoprecipitation assays (*See* Section 5.2.1). Preferably, the binding properties of the molecules of the invention are also characterized by *in vitro* functional assays for

20 determining one or more Fc γ R mediator effector cell functions (*See* Section 5.2.6). In most preferred embodiments, the molecules of the invention have similar binding properties in *in vivo* models (such as those described and disclosed herein) as those in *in vitro* based assays. However, the present invention does not exclude molecules of the invention that do not exhibit the desired phenotype in *in vitro* based assays but do exhibit the desired phenotype

25 *in vivo*.

[0034] In a specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIA with a greater affinity than a comparable molecule comprising the wild-type Fc

30 region binds Fc γ RIIA, provided that said variant Fc region does not solely have a substitution at any one of positions 329, 331, or 332, and do not include or are not solely substitution with any one of: alanine at any of positions 256, 290, 298, 312, 333, 334, 359, 360, 326, or 430; a lysine at position 330; a threonine at position 339; a methionine at position 320; a serine at position 326; an asparagine at position 326; an aspartic acid at

35 position 326; a glutamic acid at position 326; a glutamine at position 334; a glutamic acid at

position 334; a methionine at position 334; a histidine at position 334; a valine at position 334; or a leucine at position 334; a lysine at position 335.

[0035] In another specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds FcγRIIA with a greater affinity than a comparable molecule comprising the wild-type Fc region binds FcγRIIA, provided that the one or more amino acid modifications do not include or are not solely substitution with an alanine at any of positions 256, 290, 326, 255, 258, 267, 272, 276, 280, 283, 285, 286, 331, 337, 268, 272, or 430; an asparagine at position 268; a glutamine at position 272; a glutamine, serine, or aspartic acid at position 286; a serine at position 290; a methionine, glutamine, glutamic acid, or arginine at position 320; a glutamic acid at position 322; a serine, glutamic acid, or aspartic acid at position 326; a lysine at position 330; a glutamine at position 335; or a methionine at position 301.

[0036] In a preferred specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said molecule has an altered affinity for an FcγR, provided that said variant Fc region does not have a substitution at positions that make a direct contact with FcγR based on crystallographic and structural analysis of Fc-FcγR interactions such as those disclosed by Sondermann *et al.*, (2000 *Nature*, 406: 267-273, which is incorporated herein by reference in its entirety). Examples of positions within the Fc region that make a direct contact with FcγR are amino acids 234-239 (hinge region), amino acids 265-269 (B/C loop), amino acids 297-299 (C'/E loop), and amino acids 327-332 (F/G) loop. In some embodiments, the molecules of the invention comprising variant Fc regions comprise modification of at least one residue that does not make a direct contact with an FcγR based on structural and crystallographic analysis, *e.g.*, is not within the Fc-FcγR binding site.

[0037] In another preferred embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said molecule binds an FcγR with an altered affinity relative to a molecule comprising a wild-type Fc region, provided that said at least one amino acid modification do not include or are not solely a substitution at any of positions 255, 256, 258, 267, 268, 269, 270, 272, 276, 278, 280, 283, 285, 286, 289, 290, 292, 293, 294, 295, 296, 298, 300, 301, 303, 305, 307, 309, 312, 320, 322, 326, 329, 330, 332, 331, 333, 334, 335, 337, 338, 339, 340, 359, 360, 373, 376, 416, 419, 430,

434, 435, 437, 438, 439. In a specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said molecule binds an FcγR with an altered affinity relative to a molecule comprising a wild-type Fc region, provided
5 that said variant Fc region does not include or are not solely a substitution at any of positions 255, 258, 267, 269, 270, 276, 278, 280, 283, 285, 289, 292, 293, 294, 295, 296, 300, 303, 305, 307, 309, 322, 329, 332, 331, 337, 338, 340, 373, 376, 416, 419, 434, 435, 437, 438, 439 and does not have an alanine at any of positions 256, 290, 298, 312, 333, 334, 359, 360, 326, or 430; a lysine at position 330; a threonine at position 339; a methionine at
10 position 320; a serine at position 326; an asparagine at position 326; an aspartic acid at position 326; a glutamic acid at position 326; a glutamine at position 334; a glutamic acid at position 334; a methionine at position 334; a histidine at position 334; a valine at position 334; or a leucine at position 334; a lysine at position 335 an asparagine at position 268; a glutamine at position 272; a glutamine, serine, or aspartic acid at position 286; a serine at
15 position 290; a methionine, glutamine, glutamic acid, or arginine at position 320; a glutamic acid at position 322; a serine, glutamic acid, or aspartic acid at position 326; a lysine at position 330; a glutamine at position 335; or a methionine at position 301.

[0038] In a specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region does not include or are not solely a
20 substitution at any of positions 268, 269, 270, 272, 276, 278, 283, 285, 286, 289, 292, 293, 301, 303, 305, 307, 309, 331, 333, 334, 335, 337, 338, 340, 360, 373, 376, 416, 419, 430, 434, 435, 437, 438 or 439 and does not have a histidine, glutamine, or tyrosine at position 280; a serine, glycine, threonine or tyrosine at position 290, a leucine or isoleucine at position 300; an asparagine at position 294, a proline at position 296; a proline, asparagine,
25 aspartic acid, or valine at position 298; a lysine at position 295. In yet another preferred embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said molecule binds an FcγR with a reduced affinity relative to a molecule comprising a wild-type Fc region provided that said variant Fc region does not
30 have or are not solely have a substitution at any of positions 252, 254, 265, 268, 269, 270, 278, 289, 292, 293, 294, 295, 296, 298, 300, 301, 303, 322, 324, 327, 329, 333, 335, 338, 340, 373, 376, 382, 388, 389, 414, 416, 419, 434, 435, 437, 438, or 439. In yet another preferred embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification
35 relative to a wild-type Fc region, such that said molecule binds an FcγR with an enhanced

affinity relative to a molecule comprising a wild-type Fc region provided that said variant Fc region does not have or are not solely a substitution at any of positions 280, 283, 285, 286, 290, 294, 295, 298, 300, 301, 305, 307, 309, 312, 315, 331, 333, 334, 337, 340, 360, 378, 398, or 430.

- 5 [0039] In a specific embodiment, molecules of the invention comprise a variant Fc region having one or more amino acid modifications (*e.g.*, substitutions), which modifications increase the affinity of the variant Fc region for Fc γ RIIA and/or Fc γ RIIA by at least 2-fold, relative to a comparable molecule comprising a wild-type Fc region. In certain embodiments, molecules of the invention comprise a variant Fc region having one or
10 more amino acid modifications (*e.g.*, substitutions), which modifications increase the affinity of the variant Fc region for Fc γ RIIA and/or Fc γ RIIA by greater than 2-fold, at least 4-fold, at least 5-fold, at least 6-fold, at least 8-fold, or at least 10-fold relative to a comparable molecule comprising a wild-type Fc region. In other embodiments of the invention, molecules of the invention comprising a variant Fc region specifically bind
15 Fc γ RIIA and/or Fc γ RIIA with at least 65%, at least 75%, at least 85%, at least 95%, at least 100%, at least 150%, at least 200% greater affinity relative to a molecule comprising a wild-type Fc region. Such measurements are preferably *in vitro* assays.

- [0040] The invention encompasses molecules with altered affinities for the activating and/or inhibitory Fc γ receptors. In particular, the invention contemplates
20 molecules with variant Fc regions, having one or more amino acid modifications, which modifications increase the affinity of the variant Fc region for Fc γ RIIB but decrease the affinity of the variant Fc region for Fc γ RIIA and/or Fc γ RIIA, relative to a comparable molecule with a wild-type Fc region. In other embodiments, the invention encompasses molecules with variant Fc regions, having one or more amino acid modifications, which
25 modifications decrease the affinity of the variant Fc region for Fc γ RIIB and also decrease the affinity of the variant Fc regions for Fc γ RIIA and/or Fc γ RIIA relative to a comparable molecule with a wild-type Fc region. In yet other embodiments, the invention encompasses molecules with variant Fc regions, which modifications decrease the affinity of the variant Fc region for Fc γ RIIA and/or Fc γ RIIA but do not alter the affinity of the variant Fc region
30 for Fc γ RIIB relative to a comparable molecule with a wild-type Fc region.

[0041] In a specific embodiment, the molecules of the invention comprise a variant Fc region, having one or more amino acid modifications (*e.g.*, substitutions), which one or more modifications increase the affinity of the variant Fc region for Fc γ RIIA and decrease the affinity of the variant Fc region for Fc γ RIIB, relative to a comparable molecule

comprising a wild-type Fc region which binds Fc γ RIIA and Fc γ RIIB with wild-type affinity. In a certain embodiment, the one or more amino acid modifications are not a substitution with alanine at any of positions 256, 298, 333, or 334.

[0042] In another specific embodiment, the molecules of the invention comprise a variant Fc region, having one or more amino acid modifications (e.g., substitutions), which one or more modifications increase the affinity of the variant Fc region for Fc γ RIIA and decrease the affinity of the variant Fc region for Fc γ RIIB, relative to a comparable molecule comprising a wild-type Fc region which binds Fc γ RIIA and Fc γ RIIB with wild-type affinity. In a certain embodiment, the one or more amino acid modifications is not a substitution with arginine at position 320.

[0043] In most preferred embodiments, the molecules of the invention with altered affinities for activating and/or inhibitory receptors having variant Fc regions, have one or more amino acid modifications, wherein said one or more amino acid modification is a substitution at position 288 with asparagine, at position 330 with serine and at position 396 with leucine (MgFc10) (See Table 5); or a substitution at position 334 with glutamic acid, at position 359 with asparagine, and at position 366 with serine (MgFc13); or a substitution at position 316 with aspartic acid, at position 378 with valine, and at position 399 with glutamic acid (MgFc27); or a substitution at position 392 with threonine, and at position 396 with leucine (MgFc38); or a substitution at position 221 with glutamic acid, at position 270 with glutamic acid, at position 308 with alanine, at position 311 with histidine, at position 396 with leucine, and at position 402 with aspartic acid (MgFc42); or a substitution at position 240 with alanine, and at position 396 with leucine (MgFc52); or a substitution at position 410 with histidine, and at position 396 with leucine (MgFc53); or a substitution at position 243 with leucine, at position 305 with isoleucine, at position 378 with aspartic acid, at position 404 with serine, and at position 396 with leucine (MgFc54); or a substitution at position 255 with isoleucine, and at position 396 with leucine (MgFc55); or a substitution at position 370 with glutamic acid and at position 396 with leucine (MgFc59).

[0044] The preferred method for screening and identifying molecules comprising variant Fc regions with altered Fc γ R affinities (e.g., enhanced Fc γ RIIA affinity) is yeast surface display technology (for review see Boder and Wittrup, 2000, *Methods in Enzymology*, 328: 430-444, which is incorporated herein by reference in its entirety). Specifically, the yeast surface display is a genetic method whereby polypeptides comprising Fc mutants are expressed on the yeast cell wall in a form accessible for interacting with Fc γ R. Yeast surface display of the mutant Fc containing polypeptides of the invention may

be performed in accordance with any of the techniques known to those skilled in the art or the specific methods described herein.

[0045] One aspect of the invention provides a method for selecting mutant Fc fusion proteins with a desirable binding property, *e.g.*, the ability of the mutant Fc fusion protein to bind Fc γ RIIIA with a greater affinity than a comparable polypeptide comprising a wild-type Fc region binds Fc γ RIIIA. Yeast cells displaying the mutant Fc fusion proteins can be screened and characterized by any biochemical or immunological based assays known to those skilled in the art for assessing binding interactions. In a specific embodiment, screening of mutant Fc fusion proteins is done using one or more biochemical based assays, *e.g.*, an ELISA assay.

[0046] In preferred embodiments, screening and identifying molecules comprising variant Fc regions with altered Fc γ R affinities (*e.g.*, enhanced Fc γ RIIIA affinity) are done using the yeast display technology as described herein in combination with one or more biochemical based assays, preferably in a high throughput manner. The one or more biochemical assays can be any assay known in the art for identifying Fc-Fc γ R interaction, *i.e.*, specific binding of an Fc region to an Fc γ R, including, but not limited to, an ELISA assay, surface plasmon resonance assays, immunoprecipitation assay, affinity chromatography, and equilibrium dialysis. In some embodiments, screening and identifying molecules comprising variant Fc regions with altered Fc γ R affinities (*e.g.*, enhanced Fc γ RIIIA affinity) are done using the yeast display technology as described herein in combination with one or more functional based assays, preferably in a high throughput manner. The functional based assays can be any assay known in the art for characterizing one or more Fc γ R mediated effector cell function such as those described herein in Section 5.2.6. Non-limiting examples of effector cell functions that can be used in accordance with the methods of the invention, include but are not limited to, antibody-dependent cell mediated cytotoxicity (ADCC), antibody-dependent phagocytosis, phagocytosis, opsonization, opsonophagocytosis, cell binding, rosetting, C1q binding, and complement dependent cell mediated cytotoxicity. In some embodiments, screening and identifying molecules comprising variant Fc regions with altered Fc γ R affinities (*e.g.*, enhanced Fc γ RIIIA affinity) are done using the yeast display technology as described herein in combination with one or more biochemical based assays in combination or in parallel with one or more functional based assays, preferably in a high throughput manner.

[0047] A preferred method for measuring the Fc γ R-Fc interaction in accordance with the invention is an assay developed by the inventors, which allows detection and

quantitation of the interaction, despite the inherently weak affinity of the receptor for its ligand, *e.g.*, in the micromolar range for Fc γ R IIB and Fc γ R IIIA. The method involves the formation of an Fc γ R complex (*e.g.*, Fc γ R IIIA, Fc γ R IIB) that has an improved avidity for an Fc region, relative to an uncomplexed Fc γ R. In a specific embodiment, the invention encompasses a method for producing a tetrameric Fc γ R complex, wherein said tetrameric complex has an enhanced affinity for an Fc region, relative to the affinity of a monomeric Fc γ R for the Fc region, said method comprising: (i) producing a fusion protein, such that a 15 amino acid AVITAG sequence operably linked to the soluble region of Fc γ R; (ii) biotinylating the protein produced using an *E. coli* BirA enzyme; (iii) mixing the biotinylated protein produced with streptavidin-phycoerythrin in an appropriate molar ratio, such that a tetrameric Fc γ R complex is formed.

[0048] In a preferred embodiment of the invention, polypeptides comprising Fc regions bind the tetrameric Fc γ R complexes, formed according to the methods of the invention, with at least an 8-fold higher affinity than they bind the monomeric uncomplexed Fc γ R. The binding of polypeptides comprising Fc regions to the tetrameric Fc γ R complexes may be determined using standard techniques known to those skilled in the art, such as for example, fluorescence activated cell sorting (FACS), radioimmunoassays, ELISA assays, *etc.*

[0049] The invention encompasses the use of the immune complexes formed according to the methods described above for determining the functionality of molecules comprising an Fc region in cell-based or cell-free assays.

[0050] In a specific embodiment, the invention provides modified immunoglobulins comprising a variant Fc region with an enhanced affinity for Fc γ R IIIA and/or Fc γ R IIA. Such immunoglobulins include IgG molecules that naturally contain Fc γ R binding regions (*e.g.*, Fc γ R IIIA and/or Fc γ R IIB binding regions), or immunoglobulin derivatives that have been engineered to contain an Fc γ R binding region (*e.g.*, Fc γ R IIIA and/or Fc γ R IIB binding regions). The modified immunoglobulins of the invention include any immunoglobulin molecule that binds, preferably, immunospecifically, *i.e.*, competes off non-specific binding as determined by immunoassays well known in the art for assaying specific antigen-antibody binding, an antigen and contains an Fc γ R binding region (*e.g.*, a Fc γ R IIIA and/or Fc γ R IIB binding region). Such antibodies include, but are not limited to, polyclonal, monoclonal, bi-specific, multi-specific, human, humanized, chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, disulfide-linked Fvs, and fragments containing either a VL or VH domain or even a complementary determining region (CDR)

that specifically binds an antigen, in certain cases, engineered to contain or fused to an Fc γ R binding region.

[0051] In certain embodiment, the invention encompasses immunoglobulins comprising a variant Fc region with an enhanced affinity for Fc γ RIIA and/or Fc γ RIIA such that the immunoglobulin has an enhanced effector function, *e.g.*, antibody dependent cell mediated cytotoxicity. The effector function of the molecules of the invention can be assayed using any assay described herein or known to those skilled in the art. In some embodiments, immunoglobulins comprising a variant Fc region with an enhanced affinity for Fc γ RIIA and/or Fc γ RIIA have an enhanced ADCC activity relative to wild-type by at least 2-fold, at least 4-fold, at least 8-fold, at least 10-fold, at least 50-fold, or at least 100-fold.

[0052] The invention encompasses engineering human or humanized therapeutic antibodies (*e.g.*, tumor specific monoclonal antibodies) in the Fc region by modification (*e.g.*, substitution, insertion, deletion) of one or more amino acid residues, which modifications modulate the affinity of the therapeutic antibody for an Fc γ R activating receptor and/or an Fc γ R inhibitory receptor. In one embodiment, the invention relates to engineering human or humanized therapeutic antibodies (*e.g.*, tumor specific monoclonal antibodies) in the Fc region by modification of one or more amino acid residues, which modifications increase the affinity of the Fc region for Fc γ RIIA and/or Fc γ RIIA. In another embodiment, the invention relates to engineering human or humanized therapeutic antibodies (*e.g.*, tumor specific monoclonal antibodies) in the Fc region by modification of one or more amino acid residues, which modification increases the affinity of the Fc region for Fc γ RIIA and/or Fc γ RIIA and further decreases the affinity of the Fc region for Fc γ RIIB. The engineered therapeutic antibodies may further have an enhanced effector function, *e.g.*, enhanced ADCC activity, phagocytosis activity, *etc.*, as determined by standard assays known to those skilled in the art.

[0053] In a specific embodiment, the invention encompasses engineering a humanized monoclonal antibody specific for Her2/neu protooncogene (*e.g.*, Ab4D5 humanized antibody as disclosed in Carter *et al.*, 1992, *Proc. Natl. Acad. Sci. USA* 89:4285-9) by modification (*e.g.*, substitution, insertion, deletion) of at least one amino acid residue which modification increases the affinity of the Fc region for Fc γ RIIA and/or Fc γ RIIA. In another specific embodiment, modification of the humanized Her2/neu monoclonal antibody may also further decrease the affinity of the Fc region for Fc γ RIIB. In yet another specific embodiment, the engineered humanized monoclonal antibodies specific for

Her2/neu may further have an enhanced effector function as determined by standard assays known in the art and disclosed and exemplified herein.

[0054] In another specific embodiment, the invention encompasses engineering a mouse human chimeric anti-CD20 monoclonal antibody, 2H7 by modification (e.g., substitution, insertion, deletion) of at least one amino acid residue which modification increases the affinity of the Fc region for Fc γ RIIIA and/or Fc γ RIIA. In another specific embodiment, modification of the anti-CD20 monoclonal antibody, 2H7 may also further decrease the affinity of the Fc region for Fc γ RIIB. In yet another specific embodiment, the engineered anti-CD20 monoclonal antibody, 2H7 may further have an enhanced effector function as determined by standard assays known in the art and disclosed and exemplified herein.

[0055] In another specific embodiment, the invention encompasses engineering an anti-Fc γ RIIB antibody including but not limited to any of the antibodies disclosed in U.S. Provisional Application No. 60/403,266 filed on August 12, 2002 and U.S. Application No. 10/643,857 filed on August 14, 2003, having Attorney Docket No. 011183-010-999, by modification (e.g., substitution, insertion, deletion) of at least one amino acid residue which modification increases the affinity of the Fc region for Fc γ RIIIA and/or Fc γ RIIA. Examples of anti-Fc γ RIIB antibodies that may be engineered in accordance with the methods of the invention are 2B6 monoclonal antibody having ATCC accession number PTA-4591 and 3H7 having ATCC accession number PTA-4592 (deposited at ATCC, 10801 University Boulevard, Manassas, VA 02209-2011, which are incorporated herein by reference. In another specific embodiment, modification of the anti-Fc γ RIIB antibody may also further decrease the affinity of the Fc region for Fc γ RIIB. In yet another specific embodiment, the engineered anti-Fc γ RIIB antibody may further have an enhanced effector function as determined by standard assays known in the art and disclosed and exemplified herein. In a specific embodiment, the 2B6 monoclonal antibody comprises a modification at position 334 with glutamic acid, at position 359 with asparagine, and at position 366 with serine (MgFc13); or a substitution at position 316 with aspartic acid, at position 378 with valine, and at position 399 with glutamic acid (MgFc27); or a substitution at position 243 with isoleucine, at position 379 with leucine, and at position 420 with valine (MgFc29); or a substitution at position 392 with threonine and at position 396 with leucine (MgFc38); or a substitution at position 221 with glutamic acid, at position 270 with glutamic acid, at position 308 with alanine, at position 311 with histidine, at position 396 with leucine, and at position 402 with aspartic (MgFc42); or a substitution at position 410 with histidine, and at position

396 with leucine (MgFc53); or a substitution at position 243 with leucine, at position 305 with isoleucine, at position 378 with aspartic acid, at position 404 with serine, and at position 396 with leucine (MgFc54); or a substitution at position 255 with isoleucine, and at position 396 with leucine (MgFc55); or a substitution at position 370 with glutamic acid, and at position 396 with leucine (MgFc59).

[0056] The present invention also includes polynucleotides that encode a molecule of the invention, including polypeptides and antibodies, identified by the methods of the invention. The polynucleotides encoding the molecules of the invention may be obtained, and the nucleotide sequence of the polynucleotides determined, by any method known in the art. The invention relates to an isolated nucleic acid encoding a molecule of the invention. The invention also provides a vector comprising said nucleic acid. The invention further provides host cells containing the vectors or polynucleotides of the invention.

[0057] The invention further provides methods for the production of the molecules of the invention. The molecules of the invention, including polypeptides and antibodies, can be produced by any method known to those skilled in the art, in particular, by recombinant expression. In a specific embodiment, the invention relates to a method for recombinantly producing a molecule of the invention, said method comprising: (i) culturing in a medium a host cell comprising a nucleic acid encoding said molecule, under conditions suitable for the expression of said molecule; and (ii) recovery of said molecule from said medium.

[0058] The molecules identified in accordance with the methods of the invention are useful in preventing, treating, or ameliorating one or more symptoms associated with a disease, disorder, or infection. The molecules of the invention are particularly useful for the treatment or prevention of a disease or disorder where an enhanced efficacy of effector cell function (*e.g.*, ADCC) mediated by Fc γ R is desired, *e.g.*, cancer, infectious disease, and in enhancing the therapeutic efficacy of therapeutic antibodies the effect of which is mediated by ADCC.

[0059] In one embodiment, the invention encompasses a method of treating cancer in a patient having a cancer characterized by a cancer antigen, said method comprising administering a therapeutically effective amount of a therapeutic antibody that binds the cancer antigen, which has been engineered in accordance with the methods of the invention. In a specific embodiment, the invention encompasses a method for treating cancer in a patient having a cancer characterized by a cancer antigen, said method comprising administering a therapeutically effective amount of a therapeutic antibody that specifically

binds said cancer antigen, said therapeutic antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said therapeutic antibody specifically binds Fc γ RIIIA with a greater affinity than the therapeutic antibody comprising the wild-type Fc region binds Fc γ RIIIA, provided that said variant Fc region does not have a substitution at positions 329, 331, or 332, and does not have an alanine at any of positions 256, 290, 298, 312, 333, 334, 359, 360, or 430; a lysine at position 330; a threonine at position 339; a methionine at position 320; a serine at position 326; an asparagine at position 326; an aspartic acid at position 326; a glutamic acid at position 326; a glutamine at position 334; a glutamic acid at position 334; a methionine at position 334; a histidine at position 334; a valine at position 334; or a leucine at position 334. In another specific embodiment, the invention encompasses a method for treating cancer in a patient having a cancer characterized by a cancer antigen, said method comprising administering a therapeutically effective amount of a therapeutic antibody that specifically binds a cancer antigen, said therapeutic antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region such that said therapeutic antibody specifically binds Fc γ RIIIA with a greater affinity than a therapeutic antibody comprising the wild-type Fc region binds Fc γ RIIIA, and said therapeutic antibody further specifically binds Fc γ RIIB with a lower affinity than a therapeutic antibody comprising the wild-type Fc region binds Fc γ RIIB, provided that said variant Fc region does not have an alanine at any of positions 256, 298, 333, or 334. The invention encompasses a method for treating cancer in a patient characterized by a cancer antigen, said method comprising administering a therapeutically effective amount of a therapeutic antibody that specifically binds said cancer antigen and said therapeutic antibody comprises a variant Fc region so that the antibody has an enhanced ADCC activity.

[0060] The invention encompasses a method of treating an autoimmune disorder and/or inflammatory disorder in a patient in need thereof, said method comprising administering to said patient a therapeutically effective amount of a molecule comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild type Fc region, such that said molecule specifically binds Fc γ RIIB with a greater affinity than a comparable molecule comprising the wild type Fc region, and said molecule further specifically binds Fc γ RIIIA with a lower affinity than a comparable molecule comprising the wild type Fc region., and said molecule binds an immune complex (e.g., an antigen/antibody complex). The invention encompasses a method of treating an autoimmune disorder and/or inflammatory disorder further

comprising administering one or more additional prophylactic or therapeutic agents, *e.g.*, immunomodulatory agents, anti-inflammatory agents, used for the treatment and/or prevention of such diseases.

5 [0061] The invention also encompasses methods for treating or preventing an infectious disease in a subject comprising administering a therapeutically or prophylactically effective amount of one or more molecules of the invention that bind an infectious agent or cellular receptor therefor. Infectious diseases that can be treated or prevented by the molecules of the invention are caused by infectious agents including but not limited to viruses, bacteria, fungi, protozoa, and viruses.

10 [0062] According to one aspect of the invention, molecules of the invention comprising variant Fc regions have an enhanced antibody effector function towards an infectious agent, *e.g.*, a pathogenic protein, relative to a comparable molecule comprising a wild-type Fc region. In a specific embodiment, molecules of the invention enhance the efficacy of treatment of an infectious disease by enhancing phagocytosis and/or
15 opsonization of the infectious agent causing the infectious disease. In another specific embodiment, molecules of the invention enhance the efficacy of treatment of an infectious disease by enhancing ADCC of infected cells causing the infectious disease.

[0063] In some embodiments, the molecules of the invention may be administered in combination with a therapeutically or prophylactically effective amount of one or
20 additional therapeutic agents known to those skilled in the art for the treatment and/or prevention of an infectious disease. The invention contemplates the use of the molecules of the invention in combination with antibiotics known to those skilled in the art for the treatment and or prevention of an infectious disease.

[0064] The invention provides pharmaceutical compositions comprising a molecule
25 of the invention, *e.g.*, a polypeptide comprising a variant Fc region, an immunoglobulin comprising a variant Fc region, a therapeutic antibody engineered in accordance with the invention, and a pharmaceutically acceptable carrier. The invention additionally provides pharmaceutical compositions further comprising one or more additional therapeutic agents, including but not limited to anti-cancer agents, anti-inflammatory agents, immunomodulatory
30 agents.

3.1 DEFINITIONS

[0065] As used herein, the term "Fc region" is used to define a C-terminal region of an IgG heavy chain. Although the boundaries may vary slightly, the human IgG heavy chain Fc region is defined to stretch from Cys226 to the carboxy terminus. The Fc region of

an IgG comprises two constant domains, CH2 and CH3. The CH2 domain of a human IgG Fc region usually extends from amino acids 231 to amino acid 341. The CH3 domain of a human IgG Fc region usually extends from amino acids 342 to 447. The Fc region of an IgG comprises two constant domains, CH2 and CH3. The CH2 domain of a human IgG Fc region (also referred to as "C γ 2" domain) usually extends from amino acid 231-340. The CH2 domain is unique in that it is not closely paired with another domain. Rather, two N-linked branched carbohydrate chains are interposed between the two CH2 domains of an intact native IgG.

[0066] Throughout the present specification, the numbering of the residues in an IgG heavy chain is that of the EU index as in Kabat *et al.*, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, NH1, MD (1991), expressly incorporated herein by references. The "EU index as in Kabat" refers to the numbering of the human IgG1 EU antibody.

[0067] The "hinge region" is generally defined as stretching from Glu216 to Pro230 of human IgG1. Hinge regions of other IgG isotypes may be aligned with the IgG1 sequence by placing the first and last cysteine residues forming inter-heavy chain S-S binds in the same positions.

[0068] As used herein, the terms "antibody" and "antibodies" refer to monoclonal antibodies, multispecific antibodies, human antibodies, humanized antibodies, synthetic antibodies, chimeric antibodies, polyclonal antibodies, camelized antibodies, single-chain Fvs (scFv), single chain antibodies, Fab fragments, F(ab') fragments, disulfide-linked bispecific Fvs (sdFv), intrabodies, and anti-idiotypic (anti-Id) antibodies (including, *e.g.*, anti-Id and anti-anti-Id antibodies to antibodies of the invention), and epitope-binding fragments of any of the above. In particular, antibodies include immunoglobulin molecules and immunologically active fragments of immunoglobulin molecules, *i.e.*, molecules that contain an antigen binding site. Immunoglobulin molecules can be of any type (*e.g.*, IgG, IgE, IgM, IgD, IgA and IgY), class (*e.g.*, IgG₁, IgG₂, IgG₃, IgG₄, IgA₁ and IgA₂) or subclass.

[0069] As used herein, the term "derivative" in the context of polypeptides or proteins refers to a polypeptide or protein that comprises an amino acid sequence which has been altered by the introduction of amino acid residue substitutions, deletions or additions. The term "derivative" as used herein also refers to a polypeptide or protein which has been modified, *i.e.*, by the covalent attachment of any type of molecule to the polypeptide or protein. For example, but not by way of limitation, an antibody may be modified, *e.g.*, by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known

protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, *etc.* A derivative polypeptide or protein may be produced by chemical modifications using techniques known to those of skill in the art, including, but not limited to specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, *etc.* Further, a derivative polypeptide or protein derivative possesses a similar or identical function as the polypeptide or protein from which it was derived.

[0070] As used herein, the term "derivative" in the context of a non-proteinaceous derivative refers to a second organic or inorganic molecule that is formed based upon the structure of a first organic or inorganic molecule. A derivative of an organic molecule includes, but is not limited to, a molecule modified, *e.g.*, by the addition or deletion of a hydroxyl, methyl, ethyl, carboxyl or amine group. An organic molecule may also be esterified, alkylated and/or phosphorylated.

[0071] As used herein, the terms "disorder" and "disease" are used interchangeably to refer to a condition in a subject. In particular, the term "autoimmune disease" is used interchangeably with the term "autoimmune disorder" to refer to a condition in a subject characterized by cellular, tissue and/or organ injury caused by an immunologic reaction of the subject to its own cells, tissues and/or organs. The term "inflammatory disease" is used interchangeably with the term "inflammatory disorder" to refer to a condition in a subject characterized by inflammation, preferably chronic inflammation. Autoimmune disorders may or may not be associated with inflammation. Moreover, inflammation may or may not be caused by an autoimmune disorder. Thus, certain disorders may be characterized as both autoimmune and inflammatory disorders.

[0072] As used herein, the term "cancer" refers to a neoplasm or tumor resulting from abnormal uncontrolled growth of cells. As used herein, cancer explicitly includes, leukemias and lymphomas. In some embodiments, cancer refers to a benign tumor, which has remained localized. In other embodiments, cancer refers to a malignant tumor, which has invaded and destroyed neighboring body structures and spread to distant sites. In some embodiments, the cancer is associated with a specific cancer antigen.

[0073] As used herein, the term "immunomodulatory agent" and variations thereof refer to an agent that modulates a host's immune system. In certain embodiments, an immunomodulatory agent is an immunosuppressant agent. In certain other embodiments, an immunomodulatory agent is an immunostimulatory agent. Immunomodulatory agents include, but are not limited to, small molecules, peptides, polypeptides, fusion proteins, antibodies, inorganic molecules, mimetic agents, and organic molecules.

[0074] As used herein, the term "epitope" refers to a fragment of a polypeptide or protein or a non-protein molecule having antigenic or immunogenic activity in an animal, preferably in a mammal, and most preferably in a human. An epitope having immunogenic activity is a fragment of a polypeptide or protein that elicits an antibody response in an animal. An epitope having antigenic activity is a fragment of a polypeptide or protein to which an antibody immunospecifically binds as determined by any method well-known to one of skill in the art, for example by immunoassays. Antigenic epitopes need not necessarily be immunogenic.

[0075] As used herein, the term "fragment" refers to a peptide or polypeptide comprising an amino acid sequence of at least 5 contiguous amino acid residues, at least 10 contiguous amino acid residues, at least 15 contiguous amino acid residues, at least 20 contiguous amino acid residues, at least 25 contiguous amino acid residues, at least 40 contiguous amino acid residues, at least 50 contiguous amino acid residues, at least 60 contiguous amino acid residues, at least 70 contiguous amino acid residues, at least 80 amino acid residues, at least 90 amino acid residues, at least 100 amino acid residues, at least 125 amino acid residues, at least 150 contiguous amino acid residues, at least 175 amino acid residues, at least 200 amino acid residues, or at least 250 amino acid residues of the amino acid sequence of another polypeptide. In a specific embodiment, a fragment of a polypeptide retains at least one function of the polypeptide.

[0076] As used herein, the terms "nucleic acids" and "nucleotide sequences" include DNA molecules (*e.g.*, cDNA or genomic DNA), RNA molecules (*e.g.*, mRNA), combinations of DNA and RNA molecules or hybrid DNA/RNA molecules, and analogs of DNA or RNA molecules. Such analogs can be generated using, for example, nucleotide analogs, which include, but are not limited to, inosine or tritylated bases. Such analogs can also comprise DNA or RNA molecules comprising modified backbones that lend beneficial attributes to the molecules such as, for example, nuclease resistance or an increased ability to cross cellular membranes. The nucleic acids or nucleotide sequences can be single-stranded, double-stranded, may contain both single-stranded and double-stranded portions, and may contain triple-stranded portions, but preferably is double-stranded DNA.

[0077] As used herein, a "therapeutically effective amount" refers to that amount of the therapeutic agent sufficient to treat or manage a disease or disorder. A therapeutically effective amount may refer to the amount of therapeutic agent sufficient to delay or minimize the onset of disease, *e.g.*, delay or minimize the spread of cancer. A therapeutically effective amount may also refer to the amount of the therapeutic agent that

provides a therapeutic benefit in the treatment or management of a disease. Further, a therapeutically effective amount with respect to a therapeutic agent of the invention means the amount of therapeutic agent alone, or in combination with other therapies, that provides a therapeutic benefit in the treatment or management of a disease.

5 [0078] As used herein, the terms "prophylactic agent" and "prophylactic agents" refer to any agent(s) which can be used in the prevention of a disorder, or prevention of recurrence or spread of a disorder. A prophylactically effective amount may refer to the amount of prophylactic agent sufficient to prevent the recurrence or spread of hyperproliferative disease, particularly cancer, or the occurrence of such in a patient, including but not limited to those predisposed to hyperproliferative disease, for example those genetically predisposed to cancer or previously exposed to carcinogens. A prophylactically effective amount may also refer to the amount of the prophylactic agent that provides a prophylactic benefit in the prevention of disease. Further, a prophylactically effective amount with respect to a prophylactic agent of the invention means that amount of prophylactic agent alone, or in combination with other agents, that provides a prophylactic benefit in the prevention of disease.

[0079] As used herein, the terms "prevent", "preventing" and "prevention" refer to the prevention of the recurrence or onset of one or more symptoms of a disorder in a subject resulting from the administration of a prophylactic or therapeutic agent.

20 [0080] As used herein, the term "in combination" refers to the use of more than one prophylactic and/or therapeutic agents. The use of the term "in combination" does not restrict the order in which prophylactic and/or therapeutic agents are administered to a subject with a disorder. A first prophylactic or therapeutic agent can be administered prior to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concomitantly with, or subsequent to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of a second prophylactic or therapeutic agent to a subject with a disorder.

4. BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 SDS-PAGE ANALYSIS OF RECOMBINANT SOLUBLE Fc γ R

[0081] The purity of recombinant soluble Fc γ R proteins was assessed by 10% polyacrylamide gel electrophoresis. The gels were stained with Coomassie blue. Lane 1: purified recombinant soluble Fc γ R111A; Lane 2: molecular weight marker; Lane 3: molecular weight marker; Lane 4: purified recombinant soluble Fc γ R111B. The dashes refer to the molecular weight of the markers, from top to bottom, they correspond to a molecular weight of 98, 50, 36, and 22 KDa respectively.

FIG. 2 ELISA ASSAY OF RECOMBINANT SOLUBLE Fc γ R

[0082] The direct binding of purified recombinant soluble Fc γ R111A to aggregated and monomeric IgG was determined using an ELISA assay. Binding of aggregated biotinylated IgG (■) and biotinylated monomeric IgG (♦) is compared. 3G8 antibody blocks the binding of aggregated biotinylated IgG to Fc γ R111A (X).

FIGS. 3 A and B CHARACTERIZATION OF Fc γ R111A TETRAMERIC COMPLEX USING AN ELISA ASSAY

- A. Soluble tetrameric Fc γ R111A complex binds soluble monomeric human IgG specifically. Binding of soluble tetrameric Fc γ R111A to human IgG is blocked by 3G8 (♦), a mouse anti-Fc γ R111A monoclonal antibody; the 4-4-20 monoclonal antibody harboring the D265A mutation was not able to block the binding of soluble tetrameric Fc γ R111A to aggregated human IgG (Δ).
- B. Binding of soluble tetrameric Fc γ R111A complex to soluble monomeric human IgG (■) is compared to the binding of monomeric soluble Fc γ R111A human IgG (□).

FIGS. 4 A and B CHARACTERIZATION OF Fc γ R111A TETRAMERIC COMPLEX USING A MAGNETIC BEAD ASSAY

- A. Fc γ R111A Complex: two Fc γ R111A (filled shape) are joined by a monoclonal antibody DJ130c (1st Ab); the anti-mouse F(ab)₂ is conjugated to PE (circle).
- B. FACS analysis of Fc γ R111A bound to Fc coated beads: (a) beads alone; (b) complex without Fc γ R111A; (c) complex with Fc γ R111A; (d) complex with Fc γ R111A and LNK16.

FIG. 5 SCHEMATIC PRESENTATION OF Fc CONTAINING CONSTRUCTS

[0083] A schematic diagram of the IgG1 Fc domains cloned into pYD1 is presented. The open box represents the hinge-CH2-CH3 domains; parallel vertical lines represent the CH1 domain. In the case of the GIF206 and 227 constructs; the N-terminal amino acids are shown. The underlined residues correspond to the hinge region; the * represents the Xpress epitope tag; hatched boxes represent the Gly4-Ser linker, and the stippled boxes represent the Aga2p gene.

FIGS. 6A-H FACS ANALYSIS OF THE Fc FUSION PROTEINS ON THE YEAST CELL WALL

[0084] Cells were incubated with either a PE-conjugated polyclonal goat anti-human Fc antibody (FIGS. 6A-D) or with HP6017 (Sigma), a mouse anti-human IgG1 Fc (CH3) specific monoclonal antibody (FIGS. 6E-H). A and E represent vector alone; Panels B and F represent the CH1-CH3 construct; Panels C and G represent the GIF227; and Panels D and H represent the GIF 206 construct.

FIGS. 7A-C BINDING OF SOLUBLE TETRAMERIC Fc γ R11A TO THE SURFACE DISPLAYED Fc FUSION PROTEINS

[0085] Cells containing pYD1-CH1 (A); pYD-CH1-D265A (B); and pYD vector (C) were grown under conditions to express Aga2p fusion proteins on the cell surface. Cells were incubated with Fc γ R11A at 0.15 mM, 7.5 mM, and 7.5 mM, respectively, and analyzed by FACS.

FIG. 8 CHARACTERIZATION OF THE BINDING OF SOLUBLE TETRAMERIC Fc γ R11A TO THE SURFACE DISPLAYED Fc FUSION PROTEINS

[0086] Binding of Fc γ R11A tetrameric complex to Fc fusion proteins on the yeast cell surface was analyzed. PE-conjugated Fc γ R11A tetrameric complexes were pre-incubated with different concentrations of 3G8 (\diamond), LNK (\blacktriangle) or an irrelevant isotype control (\blacksquare), and subsequently incubated with the yeast cells. Cells were analyzed by FACS for PE fluorescence. The percent cells that bound the Fc γ R11A tetrameric complex were plotted on the y-axis.

FIG. 9 EXAMPLE OF SORT GATE FOR SELECTING Fc MUTANTS WITH INCREASED BINDING TO Fc γ R11A

[0087] Cells were stained with PE-conjugated Fc γ R11A tetrameric complexes (y-axis) and anti-Fc-FITC conjugated antibody (x-axis). Boxed area represents sort gate set to select ~1.0% of the cell population.

FIGs. 10A-N FACS ANALYSIS OF SOME OF THE Fc MUTANTS IDENTIFIED HAVING AN INCREASED AFFINITY FOR Fc γ RIIIA TETRAMERIC COMPLEXES

[0088] Individual clones harboring the pYD-CH1 plasmid containing independent
 5 Fc mutations were amplified in selective media containing glucose, induced for Fc expression in selective media containing galactose, and subsequently analyzed by FACs. FIGS. 10A and B represent cells harboring wild-type Fc; FIGS. 10C and D represent mutant # 5; FIGS. 10E and F represent mutant # 20; FIGS. 10G and H represent mutant # 21; FIGS. 10 I and J represent mutant # 24; FIGS. 10K and L represent mutant # 25; FIGS. 10M and
 10 N represent mutant # 27. Cells were stained with Fc γ RIIIA tetrameric complex (FIGS. 10 A, C, E, G, I, K, and M) or Fc γ RIIB tetrameric complex (FIGS. 10B, D, F, H, J, L, and N).

FIGs. 11A-B CHARACTERIZATION OF Fc MUTANTS IN THE 4-4-20 MONOCLONAL ANTIBODY BY ELISA

[0089] Fc domains from the pYD-CH1 plasmids were cloned into the heavy chain
 15 of the chimeric 4-4-20 monoclonal antibody. The 4-4-20 monoclonal antibody was expressed in 293 cells and supernatants were collected. ELISA plates were coated with fluoresceine conjugated BSA to capture the chimeric 4-4-20 mutant antibodies. Fc γ RIIIA (Panel A) and Fc γ RIIB (Panel B) receptors were then coated onto the ELISA plates to
 20 which the 4-4-20 monoclonal antibodies had been absorbed in order to determine the relative affinities of the variant receptors to the Fc domains. Mutants # 15 and # 29 were non-binding isolates included as controls.

FIG. 12 ADCC ACTIVITY OF MUTANTS IN THE 4-4-20 MONOCLONAL ANTIBODY

[0090] 4-4-20 antibodies containing mutant Fc regions were assessed for their
 25 ADCC activity, and compared to the ADCC activity of a wild type 4-4-20 antibody. The mutants analyzed are as follows: MGFc-10 (K288N, A330S, P396L), MGFc-26 (D265A), MGFc-27 (G316D, A378V, D399E), MGFc28 (N315I, A379M, D399E), MGFc29 (F243I, V379L, G420V), MGFc30 (F275V), MGFc-31 (P247L, N421K), MGFc-32 (D280E, S354F, A431D, L441I), MGFc-33 (K317N, F423 deleted), MGFc-34 (F241I, E258G),
 30 MGFc-35 (R255Q, K326E), MGFc-36 (K218R, G281D, G385R)

FIGs. 13 A and B ADCC ACTIVITY OF MUTANTS IN THE HER2/NEU HUMANIZED MONOCLONAL ANTIBODY

A. Humanized HER2/neu monoclonal antibodies containing mutant Fc regions were assessed for their ADCC activity and compared to the ADCC activity of a wild type Her2/neu antibody. The mutants analyzed are as follows: MGFc-5 (V379M), MGFc-9 (F243L, V379L), MGFc-10 (K288N, A330S, P396L), MGFc-13 (K334E, T359N, T366S),
 5 MGFc-27 (G316D, A378V, D399E).

B. ADCC activity of additional mutants in the context of the humanized Her2/neu monoclonal antibody MGFc-37 (K248M), MGFc-39 (E293V Q295E, A327T), MGFc-38 (K392T, P396L), MGFc-41 (H268N, P396L), MGFc-23 (K334E, R292L), MGFc-44, MGFc-45.

10 **FIG. 14 CAPTURE OF CH 4-4-20 ANTIBODY ON BSA-FITC SURFACE**

[0091] 6 μ L of antibody at a concentration of approximately 20 μ g/mL was injected at 5 μ L/min over a BSA- fluorescein isothiocyanate (FITC) surface. BIAcore sensogram of the binding of ch 4-4-20 antibodies with mutant Fc regions on the surface of the BSA-FITC
 15 immobilized sensor ship is shown. The marker was set on wild-type captured antibody response.

FIG. 15 SENSOGRAM OF REAL TIME BIDDING OF Fc γ RIIA TO CH 4-4-20 ANTIBODIES CARRYING VARIANT Fc REGIONS

[0092] Binding of Fc γ RIIA to ch-4-4-20 antibodies carrying variant Fc regions was
 20 analyzed at 200 nM concentration. Responses were normalized at the level of ch-4-4-20 antibody obtained for wild-type.

FIG. 16 ANALYSIS OF KINETIC PARAMETERS OF Fc γ RIIA BINDING TO ANTIBODIES CARRYING VARIANT Fc REGIONS

[0093] Kinetic parameters for Fc γ RIIA binding to antibodies carrying variant Fc
 25 regions were obtained by generating separate best fit curves for 200 nM and 800 nM. Solid line indicates an association fit which was obtained based on the k_{off} values calculated for the dissociation curves in the 32-34 sec interval. K_d and k_{off} values represent the average from two concentrations.

30 **FIG. 17 SENSOGRAM OF REAL TIME BINDING OF Fc γ RIIB-Fc FUSION PROTEINS TO ANTIBODIES CARRYING VARIANT Fc REGIONS**

[0094] Binding of Fc γ RIIB-Fc fusion proteins to ch-4-4-20 antibodies carrying variant Fc regions was analyzed at 200 nM concentration. Responses were normalized at the level of ch-4-4-20 antibody obtained for wild type.

FIG. 18 ANALYSIS OF KINETIC PARAMETERS Fc γ RIIB-Fc FUSION PROTEINS TO ANTIBODIES CARRYING VARIANT Fc REGIONS

[0095] Kinetic parameters for Fc γ RIIB-Fc binding to antibodies carrying variant Fc regions were obtained by generating separate best fit curves for 200 nM and 800 nM. Solid line indicates an association fit which was obtained based on the k_{off} values calculated for the dissociation curves in the 32-34 sec. interval. K_d and K_{off} values represent the average from two concentrations.

FIG. 19 RATIOS OF K_{off} (WT)/ K_{off} (MUT) FOR Fc γ RIIA-Fc PLOTTED AGAINST ADCC DATA

[0096] Numbers higher than one show a decreased dissociation rate for Fc γ RIIA binding and increased dissociation rate for Fc γ RIIB-Fc binding relative to wild-type.

Mutants in the box have lower off rate for Fc γ RIIA binding and higher off rate for Fc γ RIIB-Fc binding.

FIG. 20 COMPETITION WITH UNLABELED Fc γ RIIA

[0097] A kinetic screen was implemented to identify Fc region mutants with improved K_{off} rates for binding Fc γ RIIA. A library of Fc region variants containing P396L mutation was incubated with 0.1 μ M biotinylated Fc γ RIIA-Linker-Avitag for one hour and then washed. Subsequently 0.8 μ M unlabeled Fc γ RIIA was incubated with the labeled yeast for different time points. Yeast was spun down and unlabeled Fc γ RIIA was removed, Receptor bound yeast was stained with SA (streptavidin):PE (phycoerythrin) for FACS analysis.

FIG. 21 FACS ANALYSIS BASED ON THE KINETIC SCREEN

[0098] Based on the calculated K_{off} from the data presented in FIG. 20, a one minute time point selection was chosen. A 10-fold excess of library was incubated with 0.1 μ M biotinylated Fc γ RIIA-Linker-Avitag monomer; cells were washed and incubated with unlabeled ligand for one minute; then washed and labeled with SA:PE. The cells were then sorted by FACS, selecting the top 0.3% binders. The nonselected P396L library was compared to the yeast cells selected for improved binding by FACS. The histograms show

the percentage of cells that are costained with both Fc γ RIIIA /PE and goat anti-human Fc/FITC (upper right).

FIG. 22 SELECTION BASED ON SOLID PHASE DEPLETION OF Fc γ RIIB Fc BINDERS

5 [0099] The P396L library was screened based on Fc γ RIIB depletion and Fc γ RIIIA selection using magnetic beads. The Fc γ RIIB depletion by magnetic beads was repeated 5 times. The resulting yeast population was analyzed and found to show greater than 50% cell staining with goat anti-human Fc and a very small percentage of cells stained with Fc γ RIIIA. Subsequently cells were selected twice by FACS using 0.1 μ M biotinylated
10 Fc γ RIIIA linker-avitag. Yeast cells were analyzed for both Fc γ RIIIA and Fc γ RIIB binding after each sort and compared to wild type binding.

FIG. 23 RELATIVE RATES OF SKBR3 TARGET CELL LYSIS MEDIATED BY CHIMERIC 4D5 HARBORING FC MUTANTS

[00100] Relative rate of lysis was calculated for each Fc mutant tested. Lysis rates
15 for 4D5 antibody with Fc mutants were divided by the rate of lysis mediated by wild type 4D5 antibody. Data from at least 2 independent assays were averaged and plotted on the histogram. For each Fc mutant data from two different antibody concentrations are shown. The antibody concentrations were chosen to flank the point along the curve at which lysis was ~50%.

20 **FIG. 24 RELATIVE RATES OF DAUDI CELL LYSIS MEDIATED BY CHIMERIC 2H7 HARBORING FC MUTANTS**

[00101] Relative rate of lysis was calculated for each Fc mutant tested. Lysis rates
for 2H7 antibody with Fc mutants were divided by the rate of lysis mediated by wild type 2H7 antibody. Data from at least 1- 2 independent assays were averaged and plotted on the
25 histogram. For each Fc mutant data from two different antibody concentrations are shown. The antibody concentrations were chosen based on the point along the curve at which lysis was ~50%.

5. DESCRIPTION OF THE PREFERRED EMBODIMENTS

30 [00102] The present invention relates to molecules, preferably polypeptides, and more preferably immunoglobulins (*e.g.*, antibodies), comprising a variant Fc region, having one or more amino acid modifications (*e.g.*, substitutions, but also including insertions or

deletions) in one or more regions, which modifications alter, *e.g.*, increase or decrease, the affinity of the variant Fc region for an Fc γ R. In some embodiments, the invention provides molecules comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, which variant Fc region binds Fc γ RIIA with a greater affinity, relative to a comparable molecule, *i.e.*, being the same as said molecule with a variant Fc region but not having the one or more amino acid modifications, comprising the wild-type Fc region as determined by methods known to one skilled in the art for determining Fc-Fc γ R interactions and methods disclosed herein, for example, an ELISA assay or a surface plasmon resonance assay. In yet other embodiments, the invention encompasses molecules comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, which variant Fc region binds Fc γ RIIA with a reduced affinity relative to a comparable molecule comprising the wild-type Fc region. In a preferred embodiment, the molecules of the invention further specifically bind Fc γ RIIB (via the Fc region) with a lower affinity than a comparable molecule comprising the wild-type Fc region binds Fc γ RIIB. In some embodiments, the invention encompasses molecules comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, which variant Fc region binds Fc γ RIIA and Fc γ RIIB with a greater affinity, relative to a comparable molecule comprising the wild-type Fc region. In other embodiments, the invention encompasses molecules comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, which variant Fc region binds Fc γ RIIB with a greater affinity, relative to a comparable molecule comprising the wild-type Fc region. In other embodiments, the invention encompasses molecules comprising variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, which variant Fc region binds Fc γ RIIB with a reduced affinity, relative to a comparable molecule comprising the wild-type Fc region.

[00103] In some embodiments, the invention encompasses molecules comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild type Fc region, which variant Fc region does not show a detectable binding to any Fc γ R (*e.g.*, does not bind Fc γ RIIA, Fc γ RIIB, or Fc γ RIIA, as determined by, for example, an ELISA assay), relative to a comparable molecule comprising the wild-type Fc region.

[00104] In a specific embodiment, the invention encompasses molecules comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild type Fc region, which variant Fc region only binds one Fc γ R, wherein said Fc γ R is Fc γ IIIA. In another specific embodiment, the invention encompasses molecules comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild type Fc region, which variant Fc region only binds one Fc γ R, wherein said Fc γ R is Fc γ RIIA. In yet another embodiment, the invention encompasses molecules comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild type Fc region, which variant Fc region only binds one Fc γ R, wherein said Fc γ R is Fc γ RIIB. The invention particularly relates to the modification of human or humanized therapeutic antibodies (e.g., tumor specific anti-angiogenic or anti-inflammatory monoclonal antibodies) for enhancing the efficacy of therapeutic antibodies by enhancing, for example, the effector function of the therapeutic antibodies, e.g., enhancing ADCC.

[00105] The affinities and binding properties of the molecules of the invention for an Fc γ R are initially determined using *in vitro* assays (biochemical or immunological based assays) known in the art for determining Fc-Fc γ R interactions, i.e., specific binding of an Fc region to an Fc γ R including but not limited to ELISA assay, surface plasmon resonance assay, immunoprecipitation assays (See Section 5.2.1). Preferably, the binding properties of the molecules of the invention are also characterized by *in vitro* functional assays for determining one or more Fc γ R mediator effector cell functions (See Section 5.2.6). In most preferred embodiments, the molecules of the invention have similar binding properties in *in vivo* models (such as those described and disclosed herein) as those in *in vitro* based assays. However, the present invention does not exclude molecules of the invention that do not exhibit the desired phenotype in *in vitro* based assays but do exhibit the desired phenotype *in vivo*.

[00106] In some embodiments, the molecules of the invention comprising a variant Fc region comprise at least one amino acid modification in the CH3 domain of the Fc region, which is defined as extending from amino acids 342-447. In other embodiments, the molecules of the invention comprising a variant Fc region comprise at least one amino acid modification in the CH2 domain of the Fc region, which is defined as extending from amino acids 231-341. In some embodiments, the molecules of the invention comprise at least two amino acid modifications, wherein one modification is in the CH3 region and one modification is in the CH2 region. The invention further encompasses amino acid modification in the hinge region. Molecules of the invention with one or more amino acid

modifications in the CH2 and/or CH3 domains have altered affinities for an FcγR as determined using methods described herein or known to one skilled in the art.

[00107] In a preferred specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said molecule has an altered affinity for an FcγR, provided that said variant Fc region does not have a substitution at positions that make a direct contact with FcγR based on crystallographic and structural analysis of Fc-FcγR interactions such as those disclosed by Sondermann *et al.*, 2000 (*Nature*, 406: 267-273 which is incorporated herein by reference in its entirety). Examples of positions within the Fc region that make a direct contact with FcγR are amino acids 234-239 (hinge region), amino acids 265-269 (B/C loop), amino acids 297-299 (C'/E loop), and amino acids 327-332 (F/G) loop. In some embodiments, the molecules of the invention comprising variant Fc regions comprise modification of at least one residue that makes a direct contact with an FcγR based on structural and crystallographic analysis.

[00108] In another preferred embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said molecule binds an FcγR with an altered affinity relative to a molecule comprising a wild-type Fc region, provided that said variant Fc region does not have or are not solely a substitution at any of positions 255, 256, 258, 267, 268, 269, 270, 272, 276, 278, 280, 283, 285, 286, 289, 290, 292, 293, 294, 295, 296, 298, 300, 301, 303, 305, 307, 309, 312, 320, 322, 326, 329, 330, 332, 331, 333, 334, 335, 337, 338, 339, 340, 359, 360, 373, 376, 416, 419, 430, 434, 435, 437, 438, 439. In a specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said molecule binds an FcγR with an altered affinity relative to a molecule comprising a wild-type Fc region, provided that said variant Fc region does not have or are not solely a substitution at any of positions 255, 258, 267, 269, 270, 276, 278, 280, 283, 285, 289, 292, 293, 294, 295, 296, 300, 303, 305, 307, 309, 322, 329, 332, 331, 337, 338, 340, 373, 376, 416, 419, 434, 435, 437, 438, 439 and does not have an alanine at any of positions 256, 290, 298, 312, 333, 334, 359, 360, 326, or 430; a lysine at position 330; a threonine at position 339; a methionine at position 320; a serine at position 326; an asparagine at position 326; an aspartic acid at position 326; a glutamic acid at position 326; a glutamine at position 334; a glutamic acid at position 334; a methionine at position 334; a histidine at position 334; a valine at position 334; or a leucine at position

334; a lysine at position 335 an asparagine at position 268; a glutamine at position 272; a glutamine, serine, or aspartic acid at position 286; a serine at position 290; a methionine, glutamine, glutamic acid, or arginine at position 320; a glutamic acid at position 322; a serine, glutamic acid, or aspartic acid at position 326; a lysine at position 330; a glutamine
5 at position 335; or a methionine at position 301.

[00109] In a specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region does not have or are not solely a substitution at any of positions 268, 269, 270, 272, 276, 278, 283, 285, 286, 289, 292, 293, 301, 303, 305, 307, 309, 331, 333, 334, 335, 337, 338, 340, 360, 373, 376, 416, 419, 430,
10 434, 435, 437, 438 or 439 and does not have a histidine, glutamine, or tyrosine at position 280; a serine, glycine, threonine or tyrosine at position 290, a leucine or isoleucine at position 300; an asparagine at position 294, a proline at position 296; a proline, asparagine, aspartic acid, or valine at position 298; a lysine at position 295. In yet another preferred embodiment, the invention encompasses a molecule comprising a variant Fc region,
15 wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said molecule binds an FcγR with a reduced affinity relative to a molecule comprising a wild-type Fc region provided that said variant Fc region does not have or are not solely a substitution at any of positions 252, 254, 265, 268, 269, 270, 278, 289, 292, 293, 294, 295, 296, 298, 300, 301, 303, 322, 324, 327, 329, 333, 335, 338, 340,
20 373, 376, 382, 388, 389, 414, 416, 419, 434, 435, 437, 438, or 439. In yet another preferred embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said molecule binds an FcγR with an enhanced affinity relative to a molecule comprising a wild-type Fc region provided that said variant Fc region
25 does not have or are not solely a substitution at any of positions 280, 283, 285, 286, 290, 294, 295, 298, 300, 301, 305, 307, 309, 312, 315, 331, 333, 334, 337, 340, 360, 378, 398, or 430.

[00110] In most preferred embodiments, the molecules of the invention with altered affinities for activating and/or inhibitory receptors having variant Fc regions, have one or
30 more amino acid modifications, wherein said one or more amino acid modification is a substitution at position 288 with asparagine, at position 330 with serine and at position 396 with leucine (MgFc10)(See Table 5); or a substitution at position 334 with glutamic acid, at position 359 with asparagine, and at position 366 with serine (MgFc13); or a substitution at position 316 with aspartic acid, at position 378 with valine, and at position 399 with
35 glutamic acid (MgFc27); or a substitution at position 392 with threonine, and at position

396 with leucine (MgFc38); or a substitution at position 221 with glutamic acid, at position 270 with glutamic acid, at position 308 with alanine, at position 311 with histidine, at position 396 with leucine, and at position 402 with aspartic acid (MgFc42); or a substitution at position 240 with alanine, and at position 396 with leucine (MgFc52); or a substitution at position 410 with histidine, and at position 396 with leucine (MgFc53); or a substitution at position 243 with leucine, at position 305 with isoleucine, at position 378 with aspartic acid, at position 404 with serine, and at position 396 with leucine (MgFc54); or a substitution at position 255 with isoleucine, and at position 396 with leucine (MgFc55); or a substitution at position 370 with glutamic acid and at position 396 with leucine (MgFc59).

- 10 **[00111]** In some embodiments, the invention encompasses molecules comprising a variant Fc region having an amino acid modification at one or more of the following positions: 185, 142, 192, 141, 132, 149, 133, 125, 162, 147, 119, 166, 251, 292, 290, 291, 252, 288, 268, 256, 262, 218, 214, 205, 215, 247, 275, 202, 289, 258, 219, 279, 222, 246, 233, 246, 268, 244, 217, 253, 246, 224, 298, 280, 255, 218, 281, 284, 216, 223, 235, 221, 15 252, 241, 258, 227, 231, 215, 274, 287, 244, 229, 287, 291, 240, 281, 232, 269, 225, 246, 246, 293, 295, 248, 276, 268, 210, 288, 227, 221, 217, 261, 210, 242, 255, 240, 250, 247, 258, 246, 282, 219, 225, 270, 263, 272, 292, 233, 247, 254, 243, 347, 339, 392, 399, 301, 315, 383, 396, 385, 348, 333, 334, 310, 337, 371, 359, 366, 359, 379, 330, 318, 395, 319, 380, 305, 309, 335, 370, 378, 394, 386, 377, 358, 384, 397, 372, 326, 320, 375, 327, 381, 20 354, 385, 335, 387, 353, 375, 383, 397, 345, 375, 389, 335, 394, 316, 399, 315, 394, 382, 390, 369, 377, 304, 323, 313, 388, 339, 317, 365, 367, 340, 311, 312, 398, 343, 352, 362, 303, 308, 327, 307, 344, 328, 393, 355, 360, 306, 361, 355, 415, 408, 409, 407, 424, 401, 402, 435, 421, 431, 441, 440, 435, 431, 442, 400, 422, 406, 411, 422, 433, 406, 423, 420, 412, 447, 443, 414, 433, 428, 446, 402, 419, 410, 404, 427, 417, 433, 436, 438, 416.
- 25 Preferably such mutations result in molecules that have an altered affinity for an FcγR and/or has an altered effector cell mediated function as determined using methods disclosed and exemplified herein and known to one skilled in the art.

[00112] The invention encompasses molecules comprising variant Fc regions consisting of or comprising any of the mutations listed in the table below in Table 2.

30

TABLE 2. EXEMPLARY MUTATIONS

SINGLE SITE MUTANTS	DOUBLE SITE MUTANTS
K392R	Q347H, A339V
N315I	S415I, L251F
S132I	K290E, L142P
P396L	G285E, P247H
P396H	K409R, S166N

A162V	E334A, K334A
R292L	R292L, K334E
T359N	K288N, A330S
T366S	R255L, E318K
V379L	F243L, E318K
K288N	V279L, P395S
A330S	K246T, Y319F
F243L	F243I, V379L
E318K	K288M, K334E
V379M	K334E, E308D
S219Y	E233D, K334E
V282M	K246T, P396H
D401V	H268D, E318D
K222N	K246I, K334N
K334I	K320E, K326E
K334E	S375C, P396L
I377F	K288N, K326N
P247L	P247L, N421K
F372Y	S298N, W381R
K326E	R255Q, K326E
H224L	V284A, F372L
F275Y	T394M, V397M
L398V	P247L, E389G
K334N	K290T, G371D
S400P	P247L, L398Q
S407I	P247L, I377F
F372Y	K326E, G385E
T366N	S298N, S407R
K414N	E258D, N384K
M352L	F241L, E258G
T225S	K370N, S440N
I377N	K317N, F423-DELETED
K248M	P227S, K290E
R292G	K334E, E380D
S298N	P291S, P353Q
D270E	V240I, V281M
E233G	P232S, S304G
	P247L, L406F
	D399E, M428L
	L251F, F372L
	D399E, G402D
	D399E, M428L
	K392T, P396L
	H268N, P396L
	K326I, P396L
	H268D, P396L
	K210M, P396L
	L358P, P396L
	K334N, P396L
	V379M, P396L

	P227S, P396L
	P217S, P396L
	Q419H, P396L
	K370E, P396L
	L242F, P396L
	R255L, P396L
	V240A, P396L
	T250A, P396L
	P247S, P396L
	L410H, P396L
	Q419L, P396L
	V427A, P396L
	E258D, P396L
	N384K, P396L
	V323I, P396L
	P244H, P396L
	V305L, P396L
	S400F, P396L
	V303I, P396L
	A330V, Q419H
	V263Q, E272D
	K326E, A330T

- [00113] In yet other embodiments, the invention encompasses molecules comprising variant Fc regions having more than two amino acid modifications. A non-limiting example of such variants is listed in the table below (Table 3). The invention encompasses mutations listed in Table 3 which further comprise one or more amino acid modifications such as
- 5 those disclosed herein.

TABLE 3. EXEMPLARY COMBINATION VARIANTS

D399E, R292L, V185M
R301C, M252L, S192T
P291S, K288E, H268L, A141V
S383N, N384K, T256N, V262L, K218E, R214I, K205E, F149Y, K133M
S408I, V215I, V125L
G385E, P247H
V348M, K334N, F275I, Y202M, K147T
H310Y, T289A, Y407V, E258D
R292L, P396L, T359N
F275I, K334N, V348M
F243L, R255L, E318K
K334E, T359N, T366S
T256S, V305I, K334E, N390S
T335N, K370E, A378V, T394M, S424L
K334E, T359N, T366S, Q386R
K288N, A330S, P396L
P244H, L358M, V379M, N384K, V397M
P217S, A378V, S408R
P247L, I253N, K334N

D312E, K327N, I378S
D280E, S354F, A431D, L441I
K218R, G281D, G385R
P247L, A330T, S440G
T355N, P387S, H435Q
P247L, A431V, S442F
P343S, P353L, S375I, S383N
E216D, E345K, S375I
K288N, A330S, P396L
K222N, T335N, K370E, A378V, T394M
G316D, A378V, D399E
N315L, V379M, T394M
K326Q, K334E, T359N, T366S
A378V, N390I, V422I
V282E, V369I, L406F
V397M, T411A, S415N
T223I, T256S, L406F
L235P, V382M, S304G, V305I, V323I
P247L, W313R, E388G
D221Y, M252I, A330G, A339T, T359N, V422I, H433L
F243I, V379L, G420V
A231V, Q386H, V412M
T215P, K274N, A287G, K334N, L365V, P396L
P244A, K326I, C367R, S375I, K447T
R301H, K340E, D399E
C229Y, A287T, V379M, P396L, L443V
E269K, K290N, Q311R, H433Y
E216D, K334R, S375I
T335N, P387S, H435Q
K246I, Q362H, K370E
K334E, E380D, G446V
V303I, V369F, M428L
K246E, V284M, V308A
E293V, Q295E, A327T
Y319F, P352L, P396L
D221E, D270E, V308A, Q311H, P396L, G402D
K290T, N390I, P396L
K288R, T307A, K344E, P396L
V273I, K326E, L328I, P396L
K326I, S408N, P396L
K261N, K210M, P396L
F243L, V305I, A378D, F404S, P396L
K290E, V369A, T393A, P396L
K210N, K222I, K320M, P396L
P217S, V305I, I309L, N390H, P396L
K246N, Q419R, P396L
P217A, T359A, P396L
V215I, K290V, P396L
F275L, Q362H, N384K, P396L
A330V, H433Q, V427M

V263Q, E272D, Q419H
N276Y, T393N, W417R
V282L, A330V, H433Y, T436R
V284M, S298N, K334E, R355W
A330V, G427M, K438R
S219T, T225K, D270E, K360R
K222E, V263Q, S298N
E233G, P247S, L306P
S219T, T225K, D270E
S254T, A330V, N361D, P243L
V284M, S298N, K334E, R355W R416T

[00114] In some embodiments, the molecules, preferably the immunoglobulins of the invention further comprise one or more glycosylation sites, so that one or more carbohydrate moieties are covalently attached to the molecule. Preferably, the antibodies of the invention with one or more glycosylation sites and/or one or more modifications in the Fc region have an enhanced antibody mediated effector function, *e.g.*, enhanced ADCC activity. In some embodiments, the invention further comprises antibodies comprising one or more modifications of amino acids that are directly or indirectly known to interact with a carbohydrate moiety of the antibody, including but not limited to amino acids at positions 241, 243, 244, 245, 249, 256, 258, 260, 262, 264, 265, 296, 299, and 301. Amino acids that directly or indirectly interact with a carbohydrate moiety of an antibody are known in the art, *see, e.g.*, Jefferis *et al.*, 1995 *Immunology Letters*, 44: 111-7, which is incorporated herein by reference in its entirety.

[00115] The invention encompasses antibodies that have been modified by introducing one or more glycosylation sites into one or more sites of the antibodies, preferably without altering the functionality of the antibody, *e.g.*, binding activity to FcγR. Glycosylation sites may be introduced into the variable and/or constant region of the antibodies of the invention. As used herein, "glycosylation sites" include any specific amino acid sequence in an antibody to which an oligosaccharide (*i.e.*, carbohydrates containing two or more simple sugars linked together) will specifically and covalently attach. Oligosaccharide side chains are typically linked to the backbone of an antibody via either N- or O-linkages. N-linked glycosylation refers to the attachment of an oligosaccharide moiety to the side chain of an asparagine residue. O-linked glycosylation refers to the attachment of an oligosaccharide moiety to a hydroxyamino acid, *e.g.*, serine, threonine. The antibodies of the invention may comprise one or more glycosylation sites, including N-linked and O-linked glycosylation sites. Any glycosylation site for N-linked or O-linked glycosylation known in the art may be used in accordance with the instant

invention. An exemplary N-linked glycosylation site that is useful in accordance with the methods of the present invention, is the amino acid sequence: Asn-X-Thr/Ser, wherein X may be any amino acid and Thr/Ser indicates a threonine or a serine. Such a site or sites may be introduced into an antibody of the invention using methods well known in the art to which this invention pertains. *See*, for example, "In Vitro Mutagenesis," Recombinant DNA: A Short Course, J. D. Watson, *et al.* W.H. Freeman and Company, New York, 1983, chapter 8, pp. 106-116, which is incorporated herein by reference in its entirety. An exemplary method for introducing a glycosylation site into an antibody of the invention may comprise: modifying or mutating an amino acid sequence of the antibody so that the desired Asn-X-Thr/Ser sequence is obtained.

[00116] In some embodiments, the invention encompasses methods of modifying the carbohydrate content of an antibody of the invention by adding or deleting a glycosylation site. Methods for modifying the carbohydrate content of antibodies are well known in the art and encompassed within the invention, *see, e.g.*, U.S. Patent No. 6,218,149; EP 0 359 096 B1; U.S. Publication No. US 2002/0028486; WO 03/035835; U.S. Publication No. 2003/0115614; U.S. Patent No. 6,218,149; U.S. Patent No. 6,472,511; all of which are incorporated herein by reference in their entirety. In other embodiments, the invention encompasses methods of modifying the carbohydrate content of an antibody of the invention by deleting one or more endogenous carbohydrate moieties of the antibody. In a specific embodiment, the invention encompasses shifting the glycosylation site of the Fc region of an antibody, by modifying positions adjacent to 297. In a specific embodiment, the invention encompasses modifying position 296 so that position 296 and not position 297 is glycosylated.

5.1 POLYPEPTIDES AND ANTIBODIES WITH VARIANT Fc REGIONS

[00117] The present invention is based, in part, on the identification of mutant human IgG1 heavy chain Fc regions, with altered affinities for different Fc γ R receptors, using a yeast display system. Accordingly, the invention relates to molecules, preferably polypeptides, and more preferably immunoglobulins (*e.g.*, antibodies), comprising a variant Fc region, having one or more amino acid modifications (*e.g.*, substitutions, but also including insertions or deletions) in one or more regions, which modifications alter the affinity of the variant Fc region for an Fc γ R. The affinities and binding properties of the molecules of the invention for an Fc γ R are initially determined using *in vitro* assays (biochemical or immunological based assays) known in the art for determining Fc-Fc γ R interactions, *i.e.*, specific binding of an Fc region to an Fc γ R including but not limited to

ELISA assay, surface plasmon resonance assay, immunoprecipitation assays (*See* Section 5.2.1). Preferably, the binding properties of the molecules of the invention are also characterized by in vitro functional assays for determining one or more Fc γ R mediator effector cell functions (*See* Section 5.2.6). In most preferred embodiments, the molecules of the invention have similar binding properties in in vivo models (such as those described and disclosed herein) as those in in vitro based assays. However, the present invention does not exclude molecules of the invention that do not exhibit the desired phenotype in in vitro based assays but do exhibit the desired phenotype in vivo.

A. MUTANTS WITH ENHANCED ALTERED AFFINITIES FOR

Fc γ RIIA and/or Fc γ RIIA

[00118] The invention encompasses molecules comprising a variant Fc region, having one or more amino acid modifications (*e.g.*, substitutions) in one or more regions, wherein such modifications alter the affinity of the variant Fc region for an activating Fc γ R. In some embodiments, molecules of the invention comprise a variant Fc region, having one or more amino acid modifications (*e.g.*, substitutions) in one or more regions, which modifications increase the affinity of the variant Fc region for Fc γ RIIA and/or Fc γ RIIA by at least 2-fold, relative to a comparable molecule comprising a wild-type Fc region. In another specific embodiment, molecules of the invention comprise a variant Fc region, having one or more amino acid modifications (*e.g.*, substitutions) in one or more regions, which modifications increase the affinity of the variant Fc region for Fc γ RIIA and/or Fc γ RIIA by greater than 2 fold, relative to a comparable molecule comprising a wild-type Fc region. In other embodiments of the invention the one or more amino acid modifications increase the affinity of the variant Fc region for Fc γ RIIA and/or Fc γ RIIA by at least 3-fold, 4-fold, 5-fold, 6-fold, 8-fold, or 10-fold relative to a comparable molecule comprising a wild-type Fc region. In yet other embodiments of the invention the one or more amino acid modifications decrease the affinity of the variant Fc region for Fc γ RIIA and/or Fc γ RIIA by at least 3-fold, 4-fold, 5-fold, 6-fold, 8-fold, or 10-fold relative to a comparable molecule comprising a wild-type Fc region. Such fold increases are preferably determined by an ELISA or surface plasmon resonance assays. In a specific embodiment, the one or more amino acid modifications do not include or are not solely a substitution at any one of positions 329, 331, or 322 with any amino acid. In certain embodiments, the one or more amino acid modification do not include or are not solely a substitution with any one of alanine at positions 256, 290, 298, 312, 333, 334, 359, 360, or 430; with lysine at position 330; with threonine at position 339; with methionine at position 320; with serine, asparagine, aspartic acid, or glutamic acid at position 326 with glutamine, glutamic acid,

methionine, histidine, valine, or leucine at position 334. In another specific embodiment, the one or more amino acid modifications do not include or are not solely a substitution at any of positions 280, 290, 300, 294, or 295. In another more specific embodiment, the one or more amino acid modification does not include or are not solely a substitution at position 5 300 with leucine or isoleucine; at position 295 with lysine; at position 294 with asparagine; at position 298 with valine, aspartic acid, proline, asparagine, or valine; at position 280 with histidine, glutamine or tyrosine; at position 290 with serine, glycine, theonine or tyrosine.

[00119] In another specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically 10 binds Fc γ R1IA with a greater affinity than a comparable molecule comprising the wild-type Fc region binds Fc γ R1IA, provided that said variant Fc region does not have an alanine at any of positions 256, 290, 326, 255, 258, 267, 272, 276, 280, 283, 285, 286, 331, 337, 268, 272, or 430; an asparagine at position 268; a glutamine at position 272; a glutamine, serine, 15 or aspartic acid at position 286; a serine at position 290; a methionine, glutamine, glutamic acid, or arginine at position 320; a glutamic acid at position 322; a serine, glutamic acid, or aspartic acid at position 326; a lysine at position 330; a glutamine at position 335; or a methionine at position 301. In a specific embodiment, molecules of the invention comprise a variant Fc region, having one or more amino acid modifications (*e.g.*, substitutions) in one 20 or more regions, which modifications increase the affinity of the variant Fc region for Fc γ R1IA by at least 2-fold, relative to a comparable molecule comprising a wild-type Fc region. In another specific embodiment, molecules of the invention comprise a variant Fc region, having one or more amino acid modifications (*e.g.*, substitutions) in one or more regions, which modifications increase the affinity of the variant Fc region for Fc γ R1IA by 25 greater than 2 fold, relative to a comparable molecule comprising a wild-type Fc region. In other embodiments of the invention the one or more amino acid modifications increase the affinity of the variant Fc region for Fc γ R1IA by at least 3-fold, 4-fold, 5-fold, 6-fold, 8-fold, or 10-fold relative to a comparable molecule comprising a wild-type Fc region

[00120] In a specific embodiment, the invention encompasses molecules, preferably 30 polypeptides, and more preferably immunoglobulins (*e.g.*, antibodies), comprising a variant Fc region, having one or more amino acid modifications (*e.g.*, substitutions but also include insertions or deletions), which modifications increase the affinity of the variant Fc region for Fc γ R1IA and/or Fc γ R1IA by at least 65%, at least 70%, at least 75%, at least 85%, at least 90%, at least 95%, at least 99%, at least 100%, at least 150%, and at least 200%, 35 relative to a comparable molecule comprising a wild-type Fc region.

[00121] In a specific embodiment, the one or more amino acid modifications which increase the affinity of the variant Fc region comprise a substitution at position 347 with histidine, and at position 339 with valine; or a substitution at position 425 with isoleucine and at position 215 with phenylalanine; or a substitution at position 408 with isoleucine, at position 215 with isoleucine, and at position 125 with leucine; or a substitution at position 385 with glutamic acid and at position 247 with histidine; or a substitution at position 348 with methionine, at position 334 with asparagine, at position 275 with isoleucine, at position 202 with methionine, and at position 147 with threonine; or a substitution at position 275 with isoleucine, at position 334 with asparagine, and at position 348 with methionine; or a substitution at position 279 with leucine and at position 395 with serine; or a substitution at position 246 with threonine and at position 319 with phenylalanine; or a substitution at position 243 with isoleucine and at position 379 with leucine; or a substitution at position 243 with leucine, at position 255 with leucine and at position 318 with lysine; or a substitution at position 334 with glutamic acid, at position 359 with asparagine, and at position 366 with serine; or a substitution at position 288 with methionine and at position 334 with glutamic acid; or a substitution at position 334 with glutamic acid and at position 380 with aspartic acid; or a substitution at position 256 with serine, at position 305 with isoleucine, at position 334 with glutamic acid and at position 390 with serine; or a substitution at position 335 with asparagine, at position 370 with glutamic acid, at position 378 with valine, at position 394 with methionine, and at position 424 with leucine; or a substitution at position 233 with aspartic acid and at position 334 with glutamic acid; or a substitution at position 334 with glutamic acid, at position 359 with asparagine, at position 366 with serine, and at position 386 with arginine; or a substitution at position 246 with threonine and at position 396 with histidine; or a substitution at position 268 with aspartic acid and at position 318 with aspartic acid; or a substitution at position 288 with asparagine, at position 330 with serine, and at position 396 with leucine; or a substitution at position 244 with histidine, at position 358 with methionine, at position 379 with methionine, at position 384 with lysine and at position 397 with methionine; or a substitution at position 217 with serine, at position 378 with valine, and at position 408 with arginine; or a substitution at position 247 with leucine, at position 253 with asparagine, and at position 334 with asparagine; or a substitution at position 246 with isoleucine, and at position 334 with asparagine; or a substitution at position 320 with glutamic acid and at position 326 with glutamic acid; or a substitution at position 375 with cysteine and at position 396 with leucine. Examples of other amino acid substitutions that results in an enhanced affinity for FcγRIIIA in vitro are disclosed below and summarized in Table 4.

[00122] The invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises a substitution at position 243 with isoleucine and at position 379 with leucine, such that said molecule binds FcγRIIIA with about a 1.5 fold higher affinity than a comparable molecule comprising the wild type Fc region binds FcγRIIIA, as determined by an ELISA assay. In a specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises a substitution at position 288 with asparagine, at position 330 with serine, and at position 396 with leucine, such that said molecule binds FcγRIIIA with about a 5 fold higher affinity than a comparable molecule comprising the wild type Fc region binds FcγRIIIA, as determined by an ELISA assay. In a specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises a substitution at position 243 with leucine and at position 255 with leucine such that said molecule binds FcγRIIIA with about a 1 fold higher affinity than a comparable molecule comprising the wild type Fc region binds FcγRIIIA, as determined by an ELISA assay. In a specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises a substitution at position 334 with glutamic acid, at position 359 with asparagine, and at position 366 with serine, such that said molecule binds FcγRIIIA with about a 1.5 fold higher affinity than a comparable molecule comprising the wild type Fc region binds FcγRIIIA, as determined by an ELISA assay. In a specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises a substitution at position 288 with methionine and at position 334 with glutamic acid, such that said molecule binds FcγRIIIA with about a 3 fold higher affinity than a comparable molecule comprising the wild type Fc region binds FcγRIIIA, as determined by an ELISA assay. In a specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises a substitution at position 316 with aspartic acid, at position 378 with valine, and at position 399 with glutamic acid, such that said molecule binds FcγRIIIA with about a 1.5 fold higher affinity than a comparable molecule comprising the wild type Fc region binds FcγRIIIA, as determined by an ELISA assay. In a specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises a substitution at position 315 with isoleucine, at position 379 with methionine, and at position 399 with glutamic acid, such that said molecule binds FcγRIIIA with about a 1 fold higher affinity than a comparable molecule comprising the wild type Fc region binds FcγRIIIA, as determined by an ELISA assay. In a

- specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises a substitution at position 243 with isoleucine, at position 379 with leucine, and at position 420 with valine, such that said molecule binds FcγRIIIA with about a 2.5 fold higher affinity than a comparable molecule comprising the wild type Fc region binds FcγRIIIA, as determined by an ELISA assay. In a specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises a substitution at position 247 with leucine, and at position 421 with lysine, such that said molecule binds FcγRIIIA with about a 3 fold higher affinity than a comparable molecule comprising the wild type Fc region binds FcγRIIIA, as determined by an ELISA assay. In a specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises a substitution at position 392 with threonine and at position 396 with leucine such that said molecule binds FcγRIIIA with about a 4.5 fold higher affinity than a comparable molecule comprising the wild type Fc region binds FcγRIIIA, as determined by an ELISA assay. In a specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises a substitution at position 293 with valine, at position 295 with glutamic acid, and at position 327 with threonine, such that said molecule binds FcγRIIIA with about a 1.5 fold higher affinity than a comparable molecule comprising the wild type Fc region binds FcγRIIIA, as determined by an ELISA assay. In a specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises a substitution at position 268 with asparagine and at position 396 with leucine, such that said molecule binds FcγRIIIA with about a 2 fold higher affinity than a comparable molecule comprising the wild type Fc region binds FcγRIIIA, as determined by an ELISA assay. In a specific embodiment, the invention encompasses a molecule comprising a variant Fc region, wherein said variant Fc region comprises a substitution at position 319 with phenylalanine, at position 352 with leucine, and at position 396 with leucine, such that said molecule binds FcγRIIIA with about a 2 fold higher affinity than a comparable molecule comprising the wild type Fc region binds FcγRIIIA, as determined by an ELISA assay.
- 30 [00123] In a specific embodiment, the invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds FcγRIIIA with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at

position 396 with histidine. In a specific embodiment, the invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable

5 polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 248 with methionine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a similar affinity than a

10 comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 392 with arginine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a similar affinity than a

15 comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 315 with isoleucine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a similar affinity than a

20 comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 132 with isoleucine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a similar affinity than a

25 comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 162 with valine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a

30 comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 396 with leucine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a

35 comparable polypeptide comprising the wild-type Fc region, wherein said at least one

amino acid modification comprises substitution at position 379 with methionine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 219 with tyrosine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 282 with methionine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 401 with valine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 222 with asparagine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 334 with glutamic acid. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 377 with phenylalanine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one

- amino acid modification comprises substitution at position 334 with isoleucine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a
- 5 comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 247 with leucine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a
- 10 comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 326 with glutamic acid. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a
- 15 comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 372 with tyrosine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a
- 20 comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 224 with leucine.

- [00124] The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A
- 25 with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 275 with tyrosine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A
- 30 with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 398 with valine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A
- 35 with a greater affinity than a comparable polypeptide comprising the wild-type Fc region,

wherein said at least one amino acid modification comprises substitution at position 334 with asparagine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A

5 with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 400 with proline. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A

10 with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 407 with isoleucine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A

15 with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 372 with tyrosine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A

20 with a similar affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 366 with asparagine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A

25 with a reduced affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 414 with asparagine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A

30 with a reduced affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 225 with serine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A

35 with a reduced affinity than a comparable polypeptide comprising the wild-type Fc region,

wherein said at least one amino acid modification comprises substitution at position 377 with asparagine.

[00125] In a specific embodiment, the invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIIA with about a 2 fold greater affinity than a comparable polypeptide comprising the wild-type Fc region as determined by an ELISA assay, wherein said at least one amino acid modification comprises substitution at position 379 with methionine. In another specific embodiment, the invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIIA with about a 1.5 fold greater affinity than a comparable polypeptide comprising the wild-type Fc region as determined by an ELISA assay, wherein said at least one amino acid modification comprises substitution at position 248 with methionine.

[00126] In some embodiments, the molecules of the invention have an altered affinity for Fc γ RIIIA and/or Fc γ RIIA as determined using *in vitro* assays (biochemical or immunological based assays) known in the art for determining Fc-Fc γ R interactions, *i.e.*, specific binding of an Fc region to an Fc γ R including but not limited to ELISA assay, surface plasmon resonance assay, immunoprecipitation assays (*See* Section 5.2.1). Preferably, the binding properties of these molecules with altered affinities for activating Fc γ R receptors are also correlated to their activity as determined by *in vitro* functional assays for determining one or more Fc γ R mediator effector cell functions (*See* Section 5.2.6), *e.g.*, molecules with variant Fc regions with enhanced affinity for Fc γ RIIIA have an enhanced ADCC activity. In most preferred embodiments, the molecules of the invention that have an altered binding property for an activating Fc receptor, *e.g.*, Fc γ RIIIA in an *in vitro* assay also have an altered binding property in *in vivo* models (such as those described and disclosed herein). However, the present invention does not exclude molecules of the invention that do not exhibit an altered Fc γ R binding in *in vitro* based assays but do exhibit the desired phenotype *in vivo*.

30 **B. MUTANTS WITH ENHANCED AFFINITY FOR Fc γ RIIIA AND REDUCED OR NO AFFINITY FOR Fc γ RIIB**

[00127] In a specific embodiment, the molecules of the invention comprise a variant Fc region, having one or more amino acid modifications (*i.e.*, substitutions) in one or more

regions, which one or more modifications increase the affinity of the variant Fc region for Fc γ RIIA and decreases the affinity of the variant Fc region for Fc γ RIIB, relative to a comparable molecule comprising a wild-type Fc region which binds Fc γ RIIA and Fc γ RIIB with wild-type affinity. In a certain embodiment, the one or more amino acid modifications do not include or are not solely a substitution with alanine at any of positions 256, 298, 333, 334, 280, 290, 294, 298, or 296; or a substitution at position 298 with asparagine, valine, aspartic acid, or proline; or a substitution 290 with serine.. In certain amino embodiments, the one or more amino acid modifications increases the affinity of the variant Fc region for Fc γ RIIA by at least 65%, at least 70%, at least 75%, at least 85%, at least 90%, at least 95%, at least 99%, at least 100%, at least 200%, at least 300%, at least 400% and decreases the affinity of the variant Fc region for Fc γ RIIB by at least 65%, at least 70%, at least 75%, at least 85%, at least 90%, at least 95%, at least 99%, at least 100%, at least 200%, at least 300%, at least 400%.

[00128] In a specific embodiment, the molecule of the invention comprising a variant Fc region with an enhanced affinity for Fc γ RIIA and a lowered affinity or no affinity for Fc γ RIIB, as determined based on an ELISA assay and/or an ADCC based assay using ch-4-4-20 antibody carrying the variant Fc region comprises a substitution at position 275 with isoleucine, at position 334 with asparagine, and at position 348 with methionine; or a substitution at position 279 with leucine and at position 395 with serine; or a substitution at position 246 with threonine and at position 319 with phenylalanine; or a substitution at position 243 with leucine, at position 255 with leucine, and at position 318 with lysine; or a substitution at position 334 with glutamic acid, at position 359 with asparagine and at position 366 with serine; or a substitution at position 334 with glutamic acid and at position 380 with aspartic acid; or a substitution at position 256 with serine, at position 305 with isoleucine, at position 334 with glutamic acid, and at position 390 with serine; or a substitution at position 335 with asparagine, at position 370 with glutamic acid, at position 378 with valine, at position 394 with methionine and at position 424 with leucine; or a substitution at position 233 with aspartic acid and at position 334 with glutamic acid; or a substitution at position 334 with glutamic acid, at position 359 with asparagine, at position 366 with serine and at position 386 with arginine; or a substitution at position 312 with glutamic acid, at position 327 with asparagine, and at position 378 with serine; or a substitution at position 288 with asparagine and at position 326 with asparagine; or a substitution at position 247 with leucine and at position 421 with lysine; or a substitution at position 298 with asparagine and at position 381 with arginine; or a substitution at position 280 with glutamic acid, at position 354 with phenylalanine, at position 431 with aspartic

acid, and at position 441 with isoleucine; or a substitution at position 255 with glutamine and at position 326 with glutamic acid; or a substitution at position 218 with arginine, at position 281 with aspartic acid and at position 385 with arginine; or a substitution at position 247 with leucine, at position 330 with threonine and at position 440 with glycine; or a substitution at position 284 with alanine and at position 372 with leucine; or a substitution at position 335 with asparagine, at position 387 with serine and at position 435 with glutamine; or a substitution at position 247 with leucine, at position 431 with valine and at position 442 with phenylalanine.

[00129] In a specific embodiment, the molecule of the invention comprising a variant Fc region with an enhanced affinity for Fc γ RIIA and a lowered affinity or no affinity for Fc γ RIIB as determined based on an ELISA assay and/or an ADCC based assay using ch-4-4-20 antibody carrying the variant Fc region comprises a substitution at position 379 with methionine; at position 219 with tyrosine; at position 282 with methionine; at position 401 with valine; at position 222 with asparagine; at position 334 with isoleucine; at position 334 with glutamic acid; at position 275 with tyrosine; at position 398 with valine.

[00130] The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIB with about a 3 fold lower affinity than a comparable polypeptide comprising the wild-type Fc region as determined by an ELISA assay, wherein said at least one amino acid modification comprises substitution at position 288 with asparagine, at position 330 with serine, and at position 396 with leucine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIB with about a 10-15 fold lower affinity than a comparable polypeptide comprising the wild-type Fc region as determined by an ELISA assay, wherein said at least one amino acid modification comprises substitution at position 316 with aspartic acid, at position 378 with valine, and at position 399 with glutamic acid. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIB with about a 10 fold lower affinity than a comparable polypeptide comprising the wild-type Fc region as determined by an ELISA assay, wherein said at least one amino acid modification comprises substitution at position 315 with isoleucine, at position 379 with methionine, and at position 399 with glutamic acid. The invention encompasses an isolated polypeptide comprising a variant Fc region,

wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIB with about a 7 fold lower affinity than a comparable polypeptide comprising the wild-type Fc region as determined by an ELISA assay, wherein said at least one amino acid modification

5 comprises substitution at position 243 with isoleucine, at position 379 with leucine, and at position 420 with valine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIB with about a 3 fold lower affinity than a comparable polypeptide comprising the

10 wild-type Fc region as determined by an ELISA assay, wherein said at least one amino acid modification comprises substitution at position 392 with threonine and at position 396 with leucine. The invention encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIB with about a 5

15 fold lower affinity than a comparable polypeptide comprising the wild-type Fc region as determined by an ELISA assay, wherein said at least one amino acid modification comprises substitution at position 268 with asparagine and at position 396 with leucine. The invention also encompasses an isolated polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a

20 wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIB with about a 2 fold lower affinity than a comparable polypeptide comprising the wild-type Fc region as determined by an ELISA assay, wherein said at least one amino acid modification comprises substitution at position 319 with phenylalanine, at position 352 with leucine, and at position 396 with leucine.

25 C. MUTANTS WITH ENHANCED AFFINITY TO Fc γ RIIA AND Fc γ RIIB

[00131] The invention encompasses molecules comprising variant Fc regions, having one or more amino acid modifications, which modifications increase the affinity of the variant Fc region for Fc γ RIIA and Fc γ RIIB by at least 65%, at least 70%, at least 75%, at

30 least 85%, at least 90%, at least 95%, at least 99%, at least 100%, at least 200%, at least 300%, at least 400% and decreases the affinity of the variant Fc region for Fc γ RIIB by at least 65%, at least 70%, at least 75%, at least 85%, at least 90%, at least 95%, at least 99%, at least 100%, at least 200%, at least 300%, at least 400%. In a specific embodiment, the molecule of the invention comprising a variant Fc region with an enhanced affinity for

35 Fc γ RIIA and an enhanced affinity for Fc γ RIIB (as determined based on an ELISA assay

and/or an ADCC based assay using ch-4-4-20 antibody carrying the variant Fc region as described herein) comprises a substitution at position 415 with isoleucine and at position 251 with phenylalanine; or a substitution at position 399 with glutamic acid, at position 292 with leucine, and at position 185 with methionine; or a substitution at position 408 with isoleucine, at position 215 with isoleucine, and at position 125 with leucine; or a substitution at position 385 with glutamic acid and at position 247 with histidine; or a substitution at position 348 with methionine, at position 334 with asparagine, at position 275 with isoleucine, at position 202 with methionine and at position 147 with threonine; or a substitution at position 246 with threonine and at position 396 with histidine; or a substitution at position 268 with aspartic acid and at position 318 with aspartic acid; or a substitution at position 288 with asparagine, at position 330 with serine and at position 396 with leucine; or a substitution at position 244 with histidine, at position 358 with methionine, at position 379 with methionine, at position 384 with lysine and at position 397 with methionine; or a substitution at position 217 with serine, at position 378 with valine, and at position 408 with arginine; or a substitution at position 247 with leucine, at position 253 with asparagine, and at position 334 with asparagine; or a substitution at position 246 with isoleucine and at position 334 with asparagine; or a substitution at position 320 with glutamic acid and at position 326 with glutamic acid; or a substitution at position 375 with cysteine and at position 396 with leucine; or a substitution at position 343 with serine, at position 353 with leucine, at position 375 with isoleucine, at position 383 with asparagine; or a substitution at position 394 with methionine and at position 397 with methionine; or a substitution at position 216 with aspartic acid, at position 345 with lysine and at position 375 with isoleucine; or a substitution at position 288 with asparagine, at position 330 with serine, and at position 396 with leucine; or a substitution at position 247 with leucine and at position 389 with glycine; or a substitution at position 222 with asparagine, at position 335 with asparagine, at position 370 with glutamic acid, at position 378 with valine and at position 394 with methionine; or a substitution at position 316 with aspartic acid, at position 378 with valine and at position 399 with glutamic acid; or a substitution at position 315 with isoleucine, at position 379 with methionine, and at position 394 with methionine; or a substitution at position 290 with threonine and at position 371 with aspartic acid; or a substitution at position 247 with leucine and at position 398 with glutamine; or a substitution at position 326 with glutamine; at position 334 with glutamic acid, at position 359 with asparagine, and at position 366 with serine; or a substitution at position 247 with leucine and at position 377 with phenylalanine; or a substitution at position 378 with valine, at position 390 with isoleucine and at position 422 with isoleucine; or a substitution

at position 326 with glutamic acid and at position 385 with glutamic acid; or a substitution at position 282 with glutamic acid, at position 369 with isoleucine and at position 406 with phenylalanine; or a substitution at position 397 with methionine; at position 411 with alanine and at position 415 with asparagine; or a substitution at position 223 with
5 isoleucine, at position 256 with serine and at position 406 with phenylalanine; or a substitution at position 298 with asparagine and at position 407 with arginine; or a substitution at position 246 with arginine, at position 298 with asparagine, and at position 377 with phenylalanine; or a substitution at position 235 with proline, at position 382 with methionine, at position 304 with glycine, at position 305 with isoleucine, and at position 323
10 with isoleucine; or a substitution at position 247 with leucine, at position 313 with arginine, and at position 388 with glycine; or a substitution at position 221 with tyrosine, at position 252 with isoleucine, at position 330 with glycine, at position 339 with threonine, at position 359 with asparagine, at position 422 with isoleucine, and at position 433 with leucine; or a substitution at position 258 with aspartic acid, and at position 384 with lysine; or a
15 substitution at position 241 with leucine and at position 258 with glycine; or a substitution at position 370 with asparagine and at position 440 with asparagine; or a substitution at position 317 with asparagine and a deletion at position 423; or a substitution at position 243 with isoleucine, at position 379 with leucine and at position 420 with valine; or a substitution at position 227 with serine and at position 290 with glutamic acid; or a
20 substitution at position 231 with valine, at position 386 with histidine, and at position 412 with methionine; or a substitution at positions 215 with proline, at position 274 with asparagine, at position 287 with glycine, at position 334 with asparagine, at position 365 with valine and at position 396 with leucine; or a substitution at position 293 with valine, at position 295 with glutamic acid and at position 327 with threonine; or a substitution at
25 position 319 with phenylalanine, at position 352 with leucine, and at position 396 with leucine; or a substitution at position 392 with threonine and at position 396 with leucine; or a substitution at position 268 with asparagine and at position 396 with leucine; or a substitution at position 290 with threonine, at position 390 with isoleucine, and at position 396 with leucine; or a substitution at position 326 with isoleucine and at position 396 with
30 leucine; or a substitution at position 268 with aspartic acid and at position 396 with leucine; or a substitution at position 210 with methionine and at position 396 with leucine; or a substitution at position 358 with proline and at position 396 with leucine; or a substitution at position 288 with arginine, at position 307 with alanine, at position 344 with glutamic acid, and at position 396 with leucine; or a substitution at position 273 with isoleucine, at position
35 326 with glutamic acid, at position 328 with isoleucine and at position 396 with leucine; or

a substitution at position 326 with isoleucine, at position 408 with asparagine and at position 396 with leucine; or a substitution at position 334 with asparagine and at position 396 with leucine; or a substitution at position 379 with methionine and at position 396 with leucine; or a substitution at position 227 with serine and at position 396 with leucine; or a substitution at position 217 with serine and at position 396 with leucine; or a substitution at position 261 with asparagine, at position 210 with methionine and at position 396 with leucine; or a substitution at position 419 with histidine and at position 396 with leucine; or a substitution at position 370 with glutamic acid and at position 396 with leucine; or a substitution at position 242 with phenylalanine and at position 396 with leucine; or a substitution at position 255 with leucine and at position 396 with leucine; or a substitution at position 240 with alanine and at position 396 with leucine; or a substitution at position 250 with serine and at position 396 with leucine; or a substitution at position 247 with serine and at position 396 with leucine; or a substitution at position 410 with histidine and at position 396 with leucine; or a substitution at position 419 with leucine and at position 396 with leucine; or a substitution at position 427 with alanine and at position 396 with leucine; or a substitution at position 258 with aspartic acid and at position 396 with leucine; or a substitution at position 384 with lysine and at position 396 with leucine; or a substitution at position 323 with isoleucine and at position 396 with leucine; or a substitution at position 244 with histidine and at position 396 with leucine; or a substitution at position 305 with leucine and at position 396 with leucine; or a substitution at position 400 with phenylalanine and at position 396 with leucine; or a substitution at position 303 with isoleucine and at position 396 with leucine; or a substitution at position 243 with leucine, at position 305 with isoleucine, at position 378 with aspartic acid, at position 404 with serine and at position 396 with leucine; or a substitution at position 290 with glutamic acid, at position 369 with alanine, at position 393 with alanine and at position 396 with leucine; or a substitution at position 210 with asparagine, at position 222 with isoleucine, at position 320 with methionine and at position 396 with leucine; or a substitution at position 217 with serine, at position 305 with isoleucine, at position 309 with leucine, at position 390 with histidine and at position 396 with leucine; or a substitution at position 246 with asparagine; at position 419 with arginine and at position 396 with leucine; or a substitution at position 217 with alanine, at position 359 with alanine and at position 396 with leucine; or a substitution at position 215 with isoleucine, at position 290 with valine and at position 396 with leucine; or a substitution at position 275 with leucine, at position 362 with histidine, at position 384 with lysine and at position 396 with leucine; or a substitution at position 334 with asparagine; or a substitution at position 400 with proline; or a substitution at position 407

with isoleucine; or a substitution at position 372 with tyrosine; or a substitution at position 366 with asparagine; or a substitution at position 414 with asparagine; or a substitution at position 352 with leucine; or a substitution at position 225 with serine; or a substitution at position 377 with asparagine; or a substitution at position 248 with methionine.

5 **D.. MUTANTS THAT DO NOT BIND ANY FcγR**

[00132] In some embodiments, the invention encompasses molecules comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, which variant Fc region does not bind any FcγR, as determined by standard assays known in the art and disclosed herein, relative to a comparable molecule comprising the wild type Fc region. In a specific embodiment, the one or more amino acid modifications which abolish binding to all FcγRs comprise a substitution at position 232 with serine and at position 304 with glycine; or a substitution at position 269 with lysine, at position 290 with asparagine, at position 311 with arginine, and at position 433 with tyrosine; or a substitution at position 252 with leucine; or a substitution at position 216 with aspartic acid, at position 334 with arginine, and at position 375 with isoleucine; or a substitution at position 247 with leucine and at position 406 with phenylalanine, or a substitution at position 335 with asparagine, at position 387 with serine, and at position 435 with glutamine; or a substitution at position 334 with glutamic acid, at position 380 with aspartic acid, and at position 446 with valine; or a substitution at position 303 with isoleucine, at position 369 with phenylalanine, and at position 428 with leucine; or a substitution at position 251 with phenylalanine and at position 372 with leucine; or a substitution at position 246 with glutamic acid, at position 284 with methionine and at position 308 with alanine; or a substitution at position 399 with glutamic acid and at position 402 with aspartic acid; or a substitution at position 399 with glutamic acid and at position 428 with leucine.

D. MUTANTS WITH ALTERED FcγR-MEDIATED EFFECTOR FUNCTIONS

30 [00133] In a specific embodiment, the immunoglobulins of the invention with enhanced affinity for FcγRIIIA and/or FcγRIIA have an enhanced FcγR-mediated effector function as determined using ADCC activity assays disclosed herein. Examples of effector functions that could be mediated by the molecules of the invention include, but are not limited to, C1q binding, complement-dependent cytotoxicity, antibody-dependent cell mediate cytotoxicity (ADCC), phagocytosis, *etc.* The effector functions of the molecules of

the invention can be assayed using standard methods known in the art, examples of which are disclosed in Section 5.2.6. In a specific embodiment, the immunoglobulins of the invention comprising a variant Fc region with enhanced affinity for Fc γ RIIA and/or Fc γ RIIA mediate antibody dependent cell mediated cytotoxicity (ADCC) 2- fold more effectively, than an immunoglobulin comprising a wild-type Fc region. In other embodiments, the immunoglobulins of the invention comprising a variant Fc region with enhanced affinity for Fc γ RIIA and/or Fc γ RIIA mediate antibody dependent cell mediated cytotoxicity (ADCC) at least 4- fold, at least 8-fold, at least 10-fold, at least 100-fold, at least 1000-fold, at least 10⁴-fold, at least 10⁵-fold more effectively, than an immunoglobulin comprising a wild-type Fc region. In another specific embodiment, the immunoglobulins of the invention with enhanced affinity for Fc γ RIIA and/or Fc γ RIIA have altered C1q binding activity. In some embodiments, the immunoglobulins of the invention with enhanced affinity for Fc γ RIIA and/or Fc γ RIIA have at least 2-fold, at least 4- fold, at least 8-fold, at least 10-fold, at least 100-fold, at least 1000-fold, at least 10⁴-fold, at least 10⁵-fold higher C1q binding activity than an immunoglobulin comprising a wild-type Fc region. In yet another specific embodiment, the immunoglobulins of the invention with enhanced affinity for Fc γ RIIA and/or Fc γ RIIA have altered complement dependent cytotoxicity. In yet another specific embodiment, the immunoglobulins of the invention with enhanced affinity for Fc γ RIIA and/or Fc γ RIIA have an enhanced complement dependent cytotoxicity than an immunoglobulin comprising a wild-type Fc region. In some embodiments, the immunoglobulins of the invention with enhanced affinity for Fc γ RIIA and/or Fc γ RIIA have at least 2-fold, at least 4- fold, at least 8-fold, at least 10-fold, at least 100-fold, at least 1000-fold, at least 10⁴-fold, at least 10⁵-fold higher complement dependent cytotoxicity than an immunoglobulin comprising a wild-type Fc region.

[00134] In other embodiments, immunoglobulins of the invention with enhanced affinity for Fc γ RIIA and/or Fc γ RIIA have enhanced phagocytosis activity relative to an immunoglobulin comprising a wild-type Fc region, as determined by standard assays known to one skilled in the art or disclosed herein. In some embodiments, the immunoglobulins of the invention with enhanced affinity for Fc γ RIIA and/or Fc γ RIIA have at least 2-fold, at least 4- fold, at least 8-fold, at least 10-fold higher phagocytosis activity relative to an immunoglobulin comprising a wild-type Fc region.

[00135] In a specific embodiment, the invention encompasses an immunoglobulin comprising a variant Fc region with one or more amino acid modifications, with an enhanced affinity for Fc γ RIIA and/or Fc γ RIIA such that the immunoglobulin has an enhanced effector function, *e.g.*, antibody dependent cell mediated cytotoxicity, or

- phagocytosis. In a specific embodiment, the one or more amino acid modifications which increase the affinity of the variant Fc region for FcγRIIIA and/or FcγRIIA and increase the ADCC activity of the immunoglobulin comprise a substitution at position 379 with methionine; or a substitution at position 243 with isoleucine and at position 379 with
- 5 leucine; or a substitution at position 288 with asparagine, at position 330 with serine, and at position 396 with leucine; or a substitution at position 243 leucine and at position 255 with leucine; or a substitution at position 334 with glutamic acid, at position 359 with asparagine, and at position 366 with serine; or a substitution at position 288 with methionine and at position 334 with glutamic acid; or a substitution at position 334 with glutamic acid and at
- 10 position 292 with leucine; or a substitution at position 316 with aspartic acid, at position 378 with valine, and at position 399 with glutamic acid; or a substitution at position 315 with isoleucine, at position 379 with methionine, and at position 399 with glutamic acid; or a substitution at position 243 with isoleucine, at position 379 with leucine, and at position 420 with valine; or a substitution at position 247 with leucine and at position 421 with
- 15 lysine; or a substitution at position 248 with methionine; or a substitution at position 392 with threonine and at position 396 with leucine; or a substitution at position 293 with valine, at position 295 with glutamic acid, and at position 327 with threonine; or a substitution at position 268 with asparagine and at position 396 with leucine; or a substitution at position 319 with phenylalanine, at position 352 with leucine, and at position 396 with leucine.
- 20 [00136] In another specific embodiment, the one or more amino acid modifications which increase the ADCC activity of the immunoglobulin is any of the mutations listed below, in table 4.

TABLE 4.

E333A, K334A
R292L, K334E
V379M
S219Y
V282M
K222N
F243I, V379L
F243L, R255L, E318K
K334I
K334E, T359N, T366S
K288M, K334E
K288N, A330S, P396L
K326E
G316D, A378V, D399E
N315I, V379M, T394M
F243I, V379L, G420V

E293V, Q295E, A327T
Y319F, P352L, P396L
K392T, P396L
K248M
H268N, P396L
K290T, N390I, P396L
K326L, P396L
H268D, P396L
K210M, P396L
L358P, P396L
K288R, T307A, K344E, P396L
V273I, K326E, L328I, P396L
K326L, S408N, P396L
K334N, P396L
V379M, P396L
P227S, P396L
P217S, P396L
K261N, K210M, P396L
Q419H, P396L
K370E, P396L
L242F, P396L
F243L, V305I, A378D, F404S, P396L
R255L, P396L
V240A, P396L
T250S, P396L
P247S, P396L
K290E, V369A, T393A, P396L
K210N, K222I, K320M, P396L
L410H, P396L
Q419L, P396L
V427A, P396L
P217S, V305I, I309L, N390H, P396L
E258D, P396L
N384K, P396L
V323I, P396L
K246N, Q419R, P396L
P217A, T359A, P396L
P244H, P396L
V215I, K290V, P396L
F275L, Q362H, N384K, P396L
V305L, P396L
S400F, P396L
V303I, P396L

[00137] The invention encompasses specific variants of the Fc region that have been identified using the methods of the invention from a yeast library of mutants after 2nd-4th-round of sorting are listed in Table 5. Table 5 summarizes the various mutants that were identified using the methods of the invention. The mutants were assayed using an ELISA

assay for determining binding to FcγRIIIA and FcγRIIB. The mutants were also tested in an ADCC assay, by cloning the Fc variants into a ch 4-4-20 antibody using methods disclosed and exemplified herein. Bolded items refer to experiments, in which the ch4-4-20 were purified prior the ADCC assay. The antibody concentration used was in the range 0.5
5 μg/mL - 1.0 μg/mL.

TABLE 5: MUTATIONS IDENTIFIED IN THE Fc REGION

Mutations	Domain	Binding to FcγRIIA (ELISA)	Binding to FcγRIIB (ELISA)	4-4-20 ADCC
pYD-CH1 library FACS screen with 3A tetramer				
Q347H; A339V	CH3	↑0.5x	NT	
S415I; L251F	CH2, CH3	↑0.5x	↑.75x	0.82
K392R	CH3	N/C	NT	
D399E; R292L; V185M	CH1, CH2, CH3	N/C	↑0.5x	0.65
K290E; L142P	CH1, CH2	N/C	NT	0.9
R301C; M252L; S192T	CH1, CH2	↓.5x	NT	
P291S; K288E; H268L; A141V	CH1, CH2	↓.5x	NT	
N315I	CH2	N/C	↑.75x	
S132I	CH1	N/C	NT	
S383N; N384K; T256N; V262L; K218E; R214I; K205E; F149Y; K133M	All	↑0.5x	NT	
S408I; V215I; V125L	CH1, CH2, CH3	↑0.5x	↑.75x	0.62
P396L	CH3	↑1x	↑1x	0.55
G385E; P247H;	CH2, CH3	↑1x	↑.75x	0.44
P396H	CH3	↑1x	↑1x	0.58
A162V	CH1	N/C	NT	
V348M; K334N; F275I; Y202M; K147T	CH1, CH2, CH3	↑0.5x	↑.75x	0.33
H310Y; T289A; G337E	CH2	↑.5x	NT	
S119F; G371S; Y407V; E258D	CH1, CH2, CH3	N/C	N/C	0.29
K409R; S166N	CH1, CH3	N/C	NT	
In vitro Site Directed mutants				
R292L	CH2	NT	NT	0.82
T359N	CH3	NT	NT	1.06

Mutations	Domain	Binding to FcγRIIIA (ELISA)	Binding to FcγRIIB (ELISA)	4-4-20 ADCC
T366S	CH3	NT	NT	0.93
E333A, K334A	CH2	NT	NT	1.41
R292L, K334E	CH2	NT	NT	1.41; 1.64
R292L, P396L, T359N	CH2, CH3	NT	NT	0.89; 1.15
V379L	CH3	NT	NT	0.83
K288N	CH2	NT	NT	0.78
A330S	CH2	NT	NT	0.52
F243L	CH2	NT	NT	0.38
E318K	CH2	NT	NT	0.86
K288N, A330S	CH2	NT	NT	0.08
R255L, E318K	CH2	NT	NT	0.82
F243L, E318K	CH2	NT	NT	0.07
Mutants in 4-4-20 mini-library				
Increased FcγRIIIA binding, decreased or no change to FcγRIIB binding				
V379M	CH3	↑2x	N/C	1.47
S219Y	Hinge	↑1x	↓ or N/B	1.28
V282M	CH2	↑1x	↓ or N/B	1.25; 1
F275L, K334N, V348M	CH2	↑0.5x	N/C	
D401V	CH3	↑0.5x	N/C	
V279L, P395S	CH2	↑1x	N/C	
K222N	Hinge	↑1x	↓ or N/B	1.33; 0.63
K246T, Y319F	CH2	↑1x	N/C	
F243L, V379L	CH2, CH3	↑1.5x	↓ or N/B	1.86; 1.35
F243L, R255L, E318K	CH2	↑1x	↓ or N/B	1.81; 1.45
K334I	CH2	↑1x	N/C	2.1; 1.97
K334E, T359N, T366S	CH2, CH3	↑1.5x	N/C	1.49; 1.45

Mutations	Domain	Binding to FcγRIIA (ELISA)	Binding to FcγRIIB (ELISA)	4-4-20 ADCC
K288M, K334E	CH2	↑ 3x	↓ or N/B	1.61; 1.69
K334E, E380D	CH2, CH3	↑ 1.5x	N/C	
T256S, V305L, K334E, N390S	CH2, CH3	↑ 1.5x	N/C	
K334E	CH2	↑ 2.5x	N/C	1.75; 2.18
T335N, K370E, A378V, T394M, S424L	CH2, CH3	↑ 0.5x	N/C	
E233D, K334E	CH2	↑ 1.5x	N/C	0.94; 1.02
K334E, T359N, T366S, Q386R	CH2	↑ 1x	N/C	
Increased Binding to FcγRIIA and FcγRIIB				
K246T, P396H	CH2, CH3	↑ 1x	↑ 2.5x	
H268D, E318D	CH2	↑ 1.5x	↑ 5x	
K288N, A330S, P396L	CH2, CH3	↑ 5x	↑ 3x	2.34; 1.66; 2.54
I377F	CH3	↑ 1.5x	↑ 0.5x	
P244H, I358M, V379M, N384K, V397M	CH2, CH3	↑ 1.75x	↑ 1.5x	
P217S, A378V, S408R	Hinge, CH3	↑ 2x	↑ 4.5x	
P247L, I253N, K334N	CH2	↑ 3x	↑ 2.5x	
P247L	CH2	↑ 0.5x	↑ 4x	0.91; 0.84
F372Y	CH3	↑ 0.75x	↑ 5.5x	0.88; 0.59
K326E	CH2	↑ 2x	↑ 3.5x	1.63; 2
K246I, K334N	CH2	↑ 0.5x	↑ 4x	0.66; 0.6
K320E, K326E	CH2	↑ 1x	↑ 1x	
H224L	Hinge	↑ 0.5x	↑ 5x	0.55; 0.53
S375C, P396L	CH3	↑ 1.5x	↑ 4.5x	
Increased FcγRIIA binding, decreased or no change to FcγRIIB binding				

Mutations	Domain	Binding to FcγRIIA (ELISA)	Binding to FcγRIIB (ELISA)	4-4-20 ADCC
D312E, K327N, J378S	CH2, CH3	↑0.5x	N/C	
K288N, K326N	CH2	↑1x	N/C	
P275Y	CH2	↑3x	N/C	0.64
P247L, N421K	CH2, CH3	↑3x	N/C	2.0
S298N, W381R	CH2, CH3	↑2x	N/C	
D280E, S354F, A431D, L441I	CH2, CH3	↑3x	N/C	0.62
R255Q, K326B	CH2	↑2x	N/C	0.79
K218R, G281D, G385R	H, CH2, CH3	↑3.5x	N/C	0.67
L398V	CH3	↑1.5x	N/C	
P247L, A330T, S440G	CH2, CH3	↑0.75x	↓0.25x	
V284A, F372L	CH2, CH3	1x	N/C	
T335N, P387S, H435Q	CH2, CH3	1.25x	N/C	
P247L, A431V, S442F	CH2, CH3	1x	N/C	
Increased Binding to FcγRIIA and FcγRIIB				
P343S, P353L, S375I, S383N	CH3	↑0.5x	↑6x	
T394M, V397M	CH3	↑0.5x	↑3x	
E216D, E345K, S375I	H, CH2, CH3	↑0.5x	↑4x	
K334N,	CH2	↑0.5x	↑2x	
K288N, A330S, P396L	CH2, CH3	↑0.5x	↑9x	
P247L, E389G	CH2, CH3	↑1.5x	↑9x	
K222N, T335N, K370E, A378V, T394M	H, CH2, CH3	↑1x	↑7x	
G316D, A378V, D399E	CH2, CH3	↑1.5x	↑14x	2.24
N315I, V379M, T394M	CH2, CH3	↑1x	↑9x	1.37
K290T, G371D,	CH2, CH3	↑0.25x	↑6x	
P247L, L398Q	CH2, CH3	↑1.25x	↑10x	

Mutations	Domain	Binding to FcγRIIA (ELISA)	Binding to FcγRIIB (ELISA)	4-4-20 ADCC
K326Q,K334E,T359N,T366S	CH2,CH3	↑ 1.5x	↑ 5x	
S400P	CH3	↑ 1x	↑ 6x	
P247L,I377F	CH2,CH3	↑ 1x	↑ 5x	
A378V,N390L,V422I	CH3	↑ 0.5x	↑ 5x	
K326E,G385E	CH2,CH3	↑ 0.5x	↑ 15x	
V282E,V369L,L406F	CH2,CH3	↑ 0.5x	↑ 7x	
V397M,T411A,S415N	CH3	↑ 0.25x	↑ 5x	
T223I,T256S,L406F	H, CH2,CH3	↑ 0.25x	↑ 6x	
S298N,S407R	CH2,CH3	↑ 0.5x	↑ 7x	
K246R,S298N,I377F	CH2,CH3	↑ 1x	↑ 5x	
S407I	CH3	↑ 0.5x	↑ 4x	
F372Y	CH3	↑ 0.5x	↑ 4x	
L235P,V382M,S304G,V305I,V323I	CH2,CH3	↑ 2x	↑ 2x	
P247L,W313R,E388G	CH2,CH3	↑ 1.5x	↑ 1x	
D221Y,M252L,A330G,A339T,T359N,V422I,H433L	H, CH2,CH3	↑ 2.5x	↑ 6x	
E258D,N384K	CH2,CH3	↑ 1.25x	↑ 4x	
F241L,E258G	CH2	↑ 2x	↑ 2.5x	-0.08
K370N,S440N	CH3	↑ 1x	↑ 3.5x	
K317N,F423-deleted	CH2,CH3	↑ 2.5x	↑ 7x	0.18
F243I,V379L,G420V	CH2,CH3	↑ 2.5x	↑ 3.5x	1.35
P227S,K290E	H, CH2	↑ 1x	↑ 0.5x	
A231V,Q386H,V412M	CH2,CH3	↑ 1.5x	↑ 6x	
T215P,K274N,A287G,K334N,I365V,P396L	H, CH2,CH3	↑ 2x	↑ 4x	
Increased Binding to FcγRIIB but not FcγRIIA				

Mutations	Domain	Binding to FcγRIIA (ELISA)	Binding to FcγRIIB (ELISA)	4-4-20 ADCC
K334E,E380D	CH2,CH3	N/C	↑4.5x	
T366N	CH3	N/C	↑5x	
P244A,K326I,C367R,S375I,K447T	CH2,CH3	N/C	↑3x	
C229Y,A287T,V379M,P396L,I443V	H, CH2,CH3	↓0.25x	↑10x	
Decreased binding to FcγRIIA and FcγRIIB				
R301H, K340E,D399E	CH2,CH3	↓0.50x	↓0.25x	
K414N	CH3	↓0.25x	N/B	
P291S,P353Q	CH2,CH3	↓0.50x	↓0.25x	
V240L, V281M	CH2	↓0.25x	↓0.25x	
P232S, S304G	CH2	N/B	N/B	
E269K,K290N,Q311R,H433Y	CH2,CH3	N/B	N/B	
M352L	CH3	N/B	N/B	
E216D,K334R,S375I	H, CH2,CH3	N/B	N/B	
P247L,I406F	CH2,CH3	N/B	N/B	
T335N,P387S,H435Q	CH2,CH3	N/B	N/B	
T225S	CH2	↓0.25x	↓0.50x	
D399E,M428L	CH3	↓0.50x	↓0.50x	
K246I,Q362H,K370E	CH2,CH3	N/B	↓0.50x	
K334E,E380D,G446V	CH2,CH3	N/B	N/B	
I377N	CH3	↓0.50x	N/B	
V303I,V369F,M428L	CH2,CH3	N/B	N/B	
L251F,F372L	CH2,CH3	N/B	N/B	
K246E,V284M,V308A	CH2,CH3	N/B	N/B	
D399E,G402D	CH3	N/B	N/B	

Mutations	Domain	Binding to FcγRIIA (ELISA)	Binding to FcγRIIB (ELISA)	4-4-20 ADCC
D399E, M428L	CH3	N/B	N/B	
FcγRIIB depletion/FcγRIIA selection: Naive Fc library.				
E293V, Q295E, A327T	CH2	↑0.4x	↑ or N/B	4.29
Y319F, P352L, P396L	CH2, CH3	↑3.4x	↑2x	1.09
K392T, P396L	CH3	↑4.5x	↑2.5x	3.07
K248M	CH2	↑0.4x	↑ or N/B	4.03
H268N, P396L	CH2, CH3	↑2.2x	↑4.5x	2.24
Solution competition 40X FcγRIIB-G2:P396L Library				
D221E, D270E, V308A, Q311H, P396L, G402D		↑3.6x	↑0.1x	3.17
Equilibrium Screen: 0.8 μM FcγRIIA monomer: P396L library				
K290T, N390I, P396L	CH2, CH3	↑2.8x	↑6.1x	1.93
K326L, P396L	CH2, CH3	↑2.9x	↑5.9x	1.16
H268D, P396L	CH2, CH3	↑3.8x	↑13.7x	2.15
K210M, P396L	CH1, CH3	↑1.9x	↑4.6x	2.02
L358P, P396L	CH3	↑1.9x	↑4.2x	1.58
K288R, T307A, K344E, P396L	CH2, CH3	↑4.1x	↑2.3x	3.3
V273I, K326E, L328I, P396L	CH2, CH3	↑1.3x	↑10.8x	0.78
K326I, S408N, P396L	CH2, CH3	↑4x	↑9.3x	1.65
K334N, P396L	CH2, CH3	↑3.1x	↑3x	2.43
V379M, P396L	CH3	↑1.9x	↑5.6x	2.01
P227S, P396L	CH2, CH3	↑1.5x	↑4x	2.01
P217S, P396L	H, CH3	↑1.6x	↑4.5x	2.04
K261N, K210M, P396L	CH2, CH3	↑2x	↑4.2x	2.06
Kinetic Screen: 0.8 μM, 1' with cold 8 μM FcγRIIA: P396L Library				

Mutations	Domain	Binding to FcγRIIA (ELISA)	Binding to FcγRIIB (ELISA)	4-4-20 ADCC
term is M, P396L	CH3	↑1.9x	↑7.2x	3.09
Q419H, P396L	CH3	↑2x	↑6.9x	2.24
K370E, P396L	CH3	↑2x	↑6.6x	2.47
L242F, P396L	CH2, CH3	↑2.5x	↑4.1x	2.4
F243L, V305I, A378D, F404S, P396L	CH2, CH3	↑1.6x	↑5.4x	3.59
R255L, P396L	CH2, CH3	↑1.8x	↑6x	2.79
V240A, P396L	CH2, CH3	↑1.3x	↑4.2x	2.35
T250S, P396L	CH2, CH3	↑1.5x	↑6.8x	1.60
P247S, P396L	CH2, CH3	↑1.2x	↑4.2x	2.10
K290E, V369A, T393A, P396L	CH2, CH3	↑1.3x	↑6.7x	1.55
K210N, K222I, K320M, P396L	H, CH2, CH3	↑2.7x	↑8.7x	1.88
L410H, P396L	CH3	↑1.7x	↑4.5x	2.00
Q419L, P396L	CH3	↑2.2x	↑6.1x	1.70
V427A, P396L	CH3	↑1.9x	↑4.7x	1.67
P217S, V305I, E09L, N390H, P396L	H, CH2, CH3	↑2x	↑7x	1.54
E258D, P396L	CH2, CH3	↑1.9x	↑4.9x	1.54
N384K, P396L	CH3	↑2.2x	↑5.2x	1.49
V323I, P396L	CH2, CH3	↑1.1x	↑8.2x	1.29
K246N, Q419R, P396L	CH2, CH3	↑1.1x	↑4.8x	1.10
P217A, T359A, P396L	H, CH2, CH3	↑1.5x	↑4.8x	1.17
P244H, P396L	CH2, CH3	↑2.5x	↑4x	1.40
V215I, K290V, P396L	H, CH2, CH3	↑2.2x	↑4.6x	1.74
F275L, Q362H, N384K, P396L	CH2, CH3	↑2.2x	↑3.7x	1.51
V305L, P396L	CH2, CH3	↑1.3x	↑5.5x	1.50
S400F, P396L	CH3	↑1.5x	↑4.7x	1.19
V303I, P396L	CH3	↑1.1x	↑4x	1.01
FcγRIIB depletion FcγRIIA 158V solid phase selection: Naïve Library				

Mutations	Domain	Binding to FcγRIIA (ELISA)	Binding to FcγRIIB (ELISA)	4-4-20 ADCC
A330V, H433Q, V427M	CH2,CH3	NT	NT	NT
V263Q, E272D, Q419H	CH2,CH3	NT	NT	NT
N276Y, T393N, W417R	CH2,CH3	NT	NT	NT
V282L, A330V, H433Y, T436R	CH2,CH3	NT	NT	NT
A330V, Q419H	CH2,CH3	NT	NT	NT
V284M, S298N, K334E, R355W	CH2,CH3	NT	NT	NT
A330V, G427M, K438R	CH2,CH3	NT	NT	NT
S219T, T225K, D270E, K360R	CH2,CH3	NT	NT	NT
K222E, V263Q, S298N	CH2	NT	NT	NT
V263Q, E272D	CH2	NT	NT	NT
R292G	CH2	NT	NT	NT
S298N	CH2	NT	NT	NT
E233G, P247S, L306P	CH2	NT	NT	NT
D270E	CH2	NT	NT	NT
S219T, T225K, D270E	CH2	NT	NT	NT
K326E, A330T	CH2	NT	NT	NT
E233G	CH2	NT	NT	NT
S254T, A330V, N361D, P243L	CH2,CH3	NT	NT	NT
FcγRIIB depletion FcγRIIA 158F solid phase selection: Naïve Library				
158F by FACS top 0.2%				
V284M, S298N, K334E, R355W R416T	CH2,CH3	NT	NT	NT

[00138] In preferred embodiments, the invention provides modified immunoglobulin molecules (*e.g.*, antibodies) with variant Fc regions, having one or more amino acid modifications, which one or more amino acid modifications increase the affinity of the molecule for Fc γ RIIIA and/or Fc γ RIIA. Such immunoglobulins include IgG molecules that naturally contain Fc γ R binding regions (*e.g.*, Fc γ RIIIA and/or Fc γ RIIB binding region), or immunoglobulin derivatives that have been engineered to contain an Fc γ R binding region (*e.g.*, Fc γ RIIIA and/or Fc γ RIIB binding region). The modified immunoglobulins of the invention include any immunoglobulin molecule that binds, preferably, immunospecifically, *i.e.*, competes off non-specific binding as determined by immunoassays well known in the art for assaying specific antigen-antibody binding, an antigen and contains an Fc γ R binding region (*e.g.*, a Fc γ RIIIA and/or Fc γ RIIB binding region). Such antibodies include, but are not limited to, polyclonal, monoclonal, bi-specific, multi-specific, human, humanized, chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, disulfide-linked Fvs, and fragments containing either a VL or VH domain or even a complementary determining region (CDR) that specifically binds an antigen, in certain cases, engineered to contain or fused to an Fc γ R binding region.

[00139] In some embodiments, the molecules of the invention comprise portions of an Fc region. As used herein the term "portion of an Fc region" refers to fragments of the Fc region, preferably a portion with effector activity and/or Fc γ R binding activity (or a comparable region of a mutant lacking such activity). The fragment of an Fc region may range in size from 5 amino acids to the entire Fc region minus one amino acids. The portion of an Fc region may be missing up to 10, up to 20, up to 30 amino acids from the N-terminus or C-terminus.

[00140] The IgG molecules of the invention are preferably IgG1 subclass of IgGs, but may also be any other IgG subclasses of given animals. For example, in humans, the IgG class includes IgG1, IgG2, IgG3, and IgG4; and mouse IgG includes IgG1, IgG2a, IgG2b, IgG2c and IgG3.

[00141] The immunoglobulins (and other polypeptides used herein) may be from any animal origin including birds and mammals. Preferably, the antibodies are human, rodent (*e.g.*, mouse and rat), donkey, sheep, rabbit, goat, guinea pig, camel, horse, or chicken. As used herein, "human" antibodies include antibodies having the amino acid sequence of a human immunoglobulin and include antibodies isolated from human immunoglobulin libraries or from animals transgenic for one or more human immunoglobulin and that do not

express endogenous immunoglobulins, as described *infra* and, for example, in U.S. Patent No. 5,939,598 by Kucherlapati *et al.*

[00142] The antibodies of the present invention may be monospecific, bispecific, trispecific or of greater multispecificity. Multispecific antibodies may be specific for different epitopes of a polypeptide or may be specific for heterologous epitopes, such as a heterologous polypeptide or solid support material. *See, e.g.*, PCT publications WO 93/17715; WO 92/08802; WO 91/00360; WO 92/05793; Tutt, *et al.*, *J. Immunol.*, 147:60-69, 1991; U.S. Patent Nos. 4,474,893; 4,714,681; 4,925,648; 5,573,920; 5,601,819; Kostelny *et al.*, *J. Immunol.*, 148:1547-1553, 1992.

10 [00143] The antibodies of the invention include derivatives that are otherwise modified, *i.e.*, by the covalent attachment of any type of molecule to the antibody such that covalent attachment does not prevent the antibody from binding antigen and/or generating an anti-idiotypic response. For example, but not by way of limitation, the antibody derivatives include antibodies that have been modified, *e.g.*, by glycosylation, acetylation, 15 pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to, specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, etc. Additionally, the derivative may contain one or more non-classical amino acids.

20 [00144] For some uses, including *in vivo* use of antibodies in humans and *in vitro* detection assays, it may be preferable to use chimeric, humanized, or human antibodies. A chimeric antibody is a molecule in which different portions of the antibody are derived from different animal species, such as antibodies having a variable region derived from a murine monoclonal antibody and a constant region derived from a human immunoglobulin. Methods for producing chimeric antibodies are known in the art. *See e.g.*, Morrison, *Science*, 229:1202, 1985; Oi *et al.*, *BioTechniques*, 4:214 1986; Gillies *et al.*, *J. Immunol. Methods*, 125:191-202, 1989; U.S. Patent Nos. 5,807,715; 4,816,567; and 4,816,397, which are incorporated herein by reference in their entireties. Humanized antibodies are antibody 25 molecules from non-human species that bind the desired antigen having one or more complementarity determining regions (CDRs) from the non-human species and framework regions and constant domains from a human immunoglobulin molecule. Often, framework residues in the human framework regions will be substituted with the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These 30 framework substitutions are identified by methods well known in the art, *e.g.*, by modeling

of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. *See, e.g., Queen et al., U.S. Patent No. 5,585,089; Riechmann et al., Nature, 332:323, 1988, which are incorporated herein by reference in*

5 their entireties. Antibodies can be humanized using a variety of techniques known in the art including, for example, CDR-grafting (EP 239,400; PCT publication WO 91/09967; U.S. Patent Nos. 5,225,539; 5,530,101 and 5,585,089), veneering or resurfacing (EP 592,106; EP 519,596; Padlan, *Molecular Immunology*, 28(4/5):489-498, 1991; Studnicka et al., *Protein Engineering*, 7(6):805-814, 1994; Roguska et al., *Proc Natl. Acad. Sci. USA*, 91:969-973,

10 1994), and chain shuffling (U.S. Patent No. 5,565,332), all of which are hereby incorporated by reference in their entireties.

[00145] Completely human antibodies are particularly desirable for therapeutic treatment of human patients. Human antibodies can be made by a variety of methods known in the art including phage display methods described above using antibody libraries

15 derived from human immunoglobulin sequences. *See U.S. Patent Nos. 4,444,887 and 4,716,111; and PCT publications WO 98/46645; WO 98/50433; WO 98/24893; WO 98/16654; WO 96/34096; WO 96/33735; and WO 91/10741, each of which is incorporated herein by reference in its entirety.*

[00146] Human antibodies can also be produced using transgenic mice which are

20 incapable of expressing functional endogenous immunoglobulins, but which can express human immunoglobulin genes. For an overview of this technology for producing human antibodies, *see Lonberg and Huszar, Int. Rev. Immunol., 13:65-93, 1995. For a detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, see, e.g., PCT publications WO*

25 *98/24893; WO 92/01047; WO 96/34096; WO 96/33735; European Patent No. 0 598 877; U.S. Patent Nos. 5,413,923; 5,625,126; 5,633,425; 5,569,825; 5,661,016; 5,545,806; 5,814,318; 5,885,793; 5,916,771; and 5,939,598, which are incorporated by reference herein in their entireties. In addition, companies such as Abgenix, Inc. (Freemont, CA), Medarex (NJ) and Genpharm (San Jose, CA) can be engaged to provide human antibodies directed*

30 *against a selected antigen using technology similar to that described above.*

[00147] Completely human antibodies which recognize a selected epitope can be generated using a technique referred to as "guided selection." In this approach a selected non-human monoclonal antibody, *e.g., a mouse antibody, is used to guide the selection of a completely human antibody recognizing the same epitope (Jespers et al., Bio/technology,*

35 *12:899-903, 1988).*

[00148] The invention encompasses engineering human or humanized therapeutic antibodies (*e.g.*, tumor specific monoclonal antibodies) in the Fc region, by modification (*e.g.*, substitution, insertion, deletion) of at least one amino acid residue, which modification increases the affinity of the Fc region for Fc γ RIIA and/or Fc γ RIIA. In another
5 embodiment, the invention relates to engineering human or humanized therapeutic antibodies (*e.g.*, tumor specific monoclonal antibodies) in the Fc region, by modification of at least one amino acid residue, which modification increases the affinity of the Fc region for Fc γ RIIA and/or Fc γ RIIA and further decreases the affinity of the Fc region for Fc γ RIIB. The engineered therapeutic antibodies may further have an enhanced effector
10 function, *e.g.*, enhanced ADCC activity, phagocytosis activity, *etc.*, as determined by standard assays known to those skilled in the art.

[00149] In a specific embodiment, the invention encompasses engineering a humanized monoclonal antibody specific for Her2/neu protooncogene (*e.g.*, Ab4D5 humanized antibody as disclosed in Carter *et al.*, 1992, *Proc. Natl. Acad. Sci. USA* 89:4285-
15 9) by modification (*e.g.*, substitution, insertion, deletion) of at least one amino acid residue which modification increases the affinity of the Fc region for Fc γ RIIA and/or Fc γ RIIA. In another specific embodiment, modification of the humanized Her2/neu monoclonal antibody may also further decrease the affinity of the Fc region for Fc γ RIIB. In yet another specific embodiment, the engineered humanized monoclonal antibodies specific for
20 Her2/neu may further have an enhanced effector function as determined by standard assays known in the art and disclosed and exemplified herein.

[00150] In another specific embodiment, the invention encompasses engineering a mouse human chimeric anti-CD20 monoclonal antibody, 2H7 by modification (*e.g.*, substitution, insertion, deletion) of at least one amino acid residue which modification
25 increases the affinity of the Fc region for Fc γ RIIA and/or Fc γ RIIA. In another specific embodiment, modification of the anti-CD20 monoclonal antibody, 2H7 may also further decrease the affinity of the Fc region for Fc γ RIIB. In yet another specific embodiment, the engineered anti-CD20 monoclonal antibody, 2H7 may further have an enhanced effector function as determined by standard assays known in the art and disclosed and exemplified
30 herein.

[00151] In another specific embodiment, the invention encompasses engineering an anti-Fc γ RIIB antibody including but not limited to any of the antibodies disclosed in U.S. Provisional Application No. 60/403,266 filed on August 12, 2002 and U.S. Application No. 10/643,857 filed on August 14, 2003, having Attorney Docket No. 011183-010-999, by
35 modification (*e.g.*, substitution, insertion, deletion) of at least one amino acid residue which

modification increases the affinity of the Fc region for Fc γ RIIA and/or Fc γ RII. Examples of anti-Fc γ RIIB antibodies that may be engineered in accordance with the methods of the invention are 2B6 monoclonal antibody having ATCC accession number PTA-4591 and 3H7 having ATCC accession number PTA-4592 (deposited at 10801 University Boulevard, Manassas, VA 02209-2011, which are incorporated herein by reference. In another specific embodiment, modification of the anti-Fc γ RIIB antibody may also further decrease the affinity of the Fc region for Fc γ RIIB. In yet another specific embodiment, the engineered anti-Fc γ RIIB antibody may further have an enhanced effector function as determined by standard assays known in the art and disclosed and exemplified herein. In a specific embodiment, the 2B6 monoclonal antibody comprises a modification at position 334 with glutamic acid, at position 359 with asparagine, and at position 366 with serine (MgFc13); or a substitution at position 316 with aspartic acid, at position 378 with valine, and at position 399 with glutamic acid (MgFc27); or a substitution at position 243 with isoleucine, at position 379 with leucine, and at position 420 with valine (MgFc29); or a substitution at position 392 with threonine and at position 396 with leucine (MgFc38); or a substitution at position 221 with glutamic acid, at position 270 with glutamic acid, at position 308 with alanine, at position 311 with histidine, at position 396 with leucine, and at position 402 with aspartic (MgFc42); or a substitution at position 410 with histidine, and at position 396 with leucine (MgFc53); or a substitution at position 243 with leucine, at position 305 with isoleucine, at position 378 with aspartic acid, at position 404 with serine, and at position 396 with leucine (MgFc54); or a substitution at position 255 with isoleucine, and at position 396 with leucine (MgFc55); or a substitution at position 370 with glutamic acid, and at position 396 with leucine (MgFc59) (See Table 5).

5.1.1 POLYPEPTIDE AND ANTIBODY CONJUGATES

[00152] Molecules of the invention (*i.e.*, polypeptides, antibodies) comprising variant Fc regions may be recombinantly fused or chemically conjugated (including both covalently and non-covalently conjugations) to heterologous polypeptides (*i.e.*, an unrelated polypeptide; or portion thereof, preferably at least 10, at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 or at least 100 amino acids of the polypeptide) to generate fusion proteins. The fusion does not necessarily need to be direct, but may occur through linker sequences.

[00153] Further, molecules of the invention (*i.e.*, polypeptides, antibodies) comprising variant Fc regions may be conjugated to a therapeutic agent or a drug moiety that modifies a given biological response. Therapeutic agents or drug moieties are not to be

construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin (*i.e.*, PE-40), or diphtheria toxin, ricin, gelonin, and pokeweed antiviral protein, a protein such as tumor necrosis factor, interferons including, but not limited to, α -interferon (IFN- α), β -interferon (IFN- β), nerve growth factor (NGF), platelet derived growth factor (PDGF), tissue plasminogen activator (TPA), an apoptotic agent (*e.g.*, TNF- α , TNF- β , AIM I as disclosed in PCT Publication No. WO 97/33899), AIM II (*see*, PCT Publication No. WO 97/34911), Fas Ligand (Takahashi *et al.*, *J. Immunol.*, 6:1567-1574, 1994), and VEGI (PCT Publication No. WO 99/23105), a thrombotic agent or an anti-angiogenic agent (*e.g.*, angiostatin or endostatin), or a biological response modifier such as, for example, a lymphokine (*e.g.*, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), and granulocyte colony stimulating factor ("G-CSF"), macrophage colony stimulating factor, ("M-CSF"), or a growth factor (*e.g.*, growth hormone ("GH"); proteases, or ribonucleases.

[00154] Molecules of the invention (*i.e.*, polypeptides, antibodies) can be fused to marker sequences, such as a peptide to facilitate purification. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz *et al.*, 1989, *Proc. Natl. Acad. Sci. USA*, 86:821-824, for instance, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags useful for purification include, but are not limited to, the hemagglutinin "HA" tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson *et al.*, *Cell*, 37:767 1984) and the "flag" tag (Knappik *et al.*, *Biotechniques*, 17(4):754-761, 1994).

[00155] Additional fusion proteins may be generated through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling"). DNA shuffling may be employed to alter the activities of molecules of the invention (*e.g.*, antibodies with higher affinities and lower dissociation rates). *See*, generally, U.S. Patent Nos. 5,605,793; 5,811,238; 5,830,721; 5,834,252; and 5,837,458, and Patten *et al.*, 1997, *Curr. Opinion Biotechnol.* 8:724-33; Harayama, 1998, *Trends Biotechnol.* 16:76; Hansson, *et al.*, 1999, *J. Mol. Biol.* 287:265; and Lorenzo and Blasco, 1998, *BioTechniques* 24:308 (each of these patents and publications are hereby incorporated by reference in its entirety). Molecules of the invention comprising variant Fc regions, or the nucleic acids encoding the molecules of the invention, may be further altered by being

subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. One or more portions of a polynucleotide encoding a molecule of the invention, may be recombined with one or more components, motifs, sections, parts, domains, fragments, *etc.* of one or more heterologous molecules.

- 5 [00156] The present invention also encompasses molecules of the invention comprising variant Fc regions (*i.e.*, antibodies, polypeptides) conjugated to a diagnostic or therapeutic agent or any other molecule for which serum half-life is desired to be increased and/or targeted to a particular subset of cells. The molecules of the invention can be used
- 10 diagnostically to, for example, monitor the development or progression of a disease, disorder or infection as part of a clinical testing procedure to, *e.g.*, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling the molecules of the invention to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, radioactive materials, positron emitting metals, and nonradioactive paramagnetic
- 15 metal ions. The detectable substance may be coupled or conjugated either directly to the molecules of the invention or indirectly, through an intermediate (such as, for example, a linker known in the art) using techniques known in the art. *See*, for example, U.S. Patent No. 4,741,900 for metal ions which can be conjugated to antibodies for use as diagnostics according to the present invention. Such diagnosis and detection can be accomplished by
- 20 coupling the molecules of the invention to detectable substances including, but not limited to, various enzymes, enzymes including, but not limited to, horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; prosthetic group complexes such as, but not limited to, streptavidin/biotin and avidin/biotin; fluorescent materials such as, but not limited to, umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine,
- 25 dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; luminescent material such as, but not limited to, luminol; bioluminescent materials such as, but not limited to, luciferase, luciferin, and aequorin; radioactive material such as, but not limited to, bismuth (^{213}Bi), carbon (^{14}C), chromium (^{51}Cr), cobalt (^{57}Co), fluorine (^{18}F), gadolinium (^{153}Gd , ^{159}Gd), gallium (^{68}Ga , ^{67}Ga), germanium (^{68}Ge), holmium (^{166}Ho), indium (^{115}In , ^{113}In , ^{112}In , ^{111}In), iodine (^{131}I , ^{125}I , ^{123}I , ^{121}I), lanthanum (^{140}La), lutetium (^{177}Lu), manganese (^{54}Mn),
- 30 molybdenum (^{99}Mo), palladium (^{103}Pd), phosphorous (^{32}P), praseodymium (^{142}Pr), promethium (^{149}Pm), rhenium (^{186}Re , ^{188}Re), rhodium (^{105}Rh), ruthenium (^{97}Ru), samarium (^{153}Sm), scandium (^{47}Sc), selenium (^{75}Se), strontium (^{85}Sr), sulfur (^{35}S), technetium (^{99}Tc), thallium (^{201}Tl), tin (^{113}Sn , ^{117}Sn), tritium (^3H), xenon (^{133}Xe), ytterbium (^{169}Yb , ^{175}Yb),

yttrium (^{90}Y), zinc (^{65}Zn); positron emitting metals using various positron emission tomographies, and nonradioactive paramagnetic metal ions.

[00157] Molecules of the invention (*i.e.*, antibodies, polypeptides) comprising a variant Fc region may be conjugated to a therapeutic moiety such as a cytotoxin (*e.g.*, a
5 cytostatic or cytocidal agent), a therapeutic agent or a radioactive element (*e.g.*, alpha-emitters, gamma-emitters, *etc.*). Cytotoxins or cytotoxic agents include any agent that is detrimental to cells. Examples include paclitaxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin,
10 actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (*e.g.*, methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (*e.g.*, mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide,
15 busulfan, dibromomannitol, streptozotocin, mitomycin C, and cisdichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (*e.g.*, daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (*e.g.*, dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC), and anti-mitotic agents (*e.g.*, vincristine and vinblastine).

20 [00158] Moreover, a molecule of the invention can be conjugated to therapeutic moieties such as a radioactive materials or macrocyclic chelators useful for conjugating radiometal ions (*see* above for examples of radioactive materials). In certain embodiments, the macrocyclic chelator is 1,4,7,10-tetraazacyclododecane- $\text{N},\text{N}',\text{N}'',\text{N}'''$ -tetraacetic acid (DOTA) which can be attached to the antibody via a linker molecule. Such linker
25 molecules are commonly known in the art and described in Denardo *et al.*, 1998, *Clin Cancer Res.* 4:2483-90; Peterson *et al.*, 1999, *Bioconjug. Chem.* 10:553; and Zimmerman *et al.*, 1999, *Nucl. Med. Biol.* 26:943-50 each of which is incorporated herein by reference in their entireties.

[00159] Techniques for conjugating such therapeutic moieties to antibodies are well
30 known; *see, e.g.*, Armon *et al.*, "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld *et al.* (eds.), 1985, pp. 243-56, Alan R. Liss, Inc.; Hellstrom *et al.*, "Antibodies For Drug Delivery", in *Controlled Drug Delivery (2nd Ed.)*, Robinson *et al.* (eds.), 1987, pp. 623-53, Marcel Dekker, Inc.; Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A
35 Review", in *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera *et*

al. (eds.), 1985, pp. 475-506); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in *Monoclonal Antibodies For Cancer Detection And Therapy*, Baldwin *et al.* (eds.), 1985, pp. 303-16, Academic Press; and Thorpe *et al.*, *Immunol. Rev.*, 62:119-58, 1982.

- 5 [00160] In one embodiment, where the molecule of the invention is an antibody comprising a variant Fc region, it can be administered with or without a therapeutic moiety conjugated to it, administered alone, or in combination with cytotoxic factor(s) and/or cytokine(s) for use as a therapeutic treatment. Alternatively, an antibody of the invention can be conjugated to a second antibody to form an antibody heteroconjugate as described by
- 10 Segal in U.S. Patent No. 4,676,980, which is incorporated herein by reference in its entirety. Antibodies of the invention may also be attached to solid supports, which are particularly useful for immunoassays or purification of the target antigen. Such solid supports include, but are not limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene.

15 5.2 **SCREENING OF MOLECULES WITH VARIANT Fc REGIONS FOR ENHANCED Fc γ RIII BINDING AND CHARACTERIZATION OF SAME**

- [00161] In preferred embodiments, screening and identifying molecules comprising variant Fc regions with altered Fc γ R affinities (*e.g.*, enhanced Fc γ RIIIA affinity) are done
- 20 using the yeast display technology as described herein in combination with one or more biochemical based assays, preferably in a high throughput manner. The one or more biochemical assays can be any assay known in the art for identifying Fc-Fc γ R interaction, *i.e.*, specific binding of an Fc region to an Fc γ R, including, but not limited to, an ELISA assay, surface plasmon resonance assays, immunoprecipitation assay, affinity
- 25 chromatography, and equilibrium dialysis. In some embodiments, screening and identifying molecules comprising variant Fc regions with altered Fc γ R affinities (*e.g.*, enhanced Fc γ RIIIA affinity) are done using the yeast display technology as described herein in combination with one or more functional based assays, preferably in a high throughput manner. The functional based assays can be any assay known in the art for characterizing
- 30 one or more Fc γ R mediated effector cell functions such as those described herein in Section 5.2.6. Non-limiting examples of effector cell functions that can be used in accordance with the methods of the invention, include but are not limited to, antibody-dependent cell mediated cytotoxicity (ADCC), antibody-dependent phagocytosis, phagocytosis, opsonization, opsonophagocytosis, cell binding, rosetting, C1q binding, and complement

dependent cell mediated cytotoxicity. In some embodiments, screening and identifying molecules comprising variant Fc regions with altered Fc γ R affinities (e.g., enhanced Fc γ RIIIA affinity) are done using the yeast display technology as described herein in combination with one or more biochemical based assays in combination or in parallel with one or more functional based assays, preferably in a high throughput manner.

[00162] The term "specific binding" of an Fc region to an Fc γ R refers to an interaction of the Fc region and a particular Fc γ R which has an affinity constant of at least about 150 nM, in the case of monomeric Fc γ RIIIA and at least about 60 nM in the case of dimeric Fc γ RIIB as determined using, for example, an ELISA or surface plasmon resonance assay (e.g., a BIAcoreTM). The affinity constant of an Fc region for monomeric Fc γ RIIIA may be 150 nM, 200 nM or 300nM. The affinity constant of an Fc region for dimeric Fc γ RIIB may be 60 nM, 80 nM, 90 nM, or 100 nM. Dimeric Fc γ RIIB for use in the methods of the invention may be generated using methods known to one skilled in the art. Typically, the extracellular region of Fc γ RIIB is covalently linked to a heterologous polypeptide which is capable of dimerization, so that the resulting fusion protein is a dimer, e.g., see, U.S. Application No. 60/439,709 filed on January 13, 2003 (Attorney Docket No. 11183-005-888), which is incorporated herein by reference in its entirety. A specific interaction generally is stable under physiological conditions, including, for example, conditions that occur in a living individual such as a human or other vertebrate or invertebrate, as well as conditions that occur in a cell culture such conditions as used for maintaining and culturing mammalian cells or cells from another vertebrate organism or an invertebrate organism.

[00163] In a specific embodiment, screening for and identifying molecules comprising variant Fc regions and altered Fc γ R affinities comprise: displaying the molecule comprising a variant Fc region on the yeast surface; and characterizing the binding of the molecule comprising the variant Fc region to a Fc γ R (one or more), using a biochemical assay for determining Fc-Fc γ R interaction, preferably, an ELISA based assay. Once the molecule comprising a variant Fc region has been characterized for its interaction with one or more Fc γ Rs and determined to have an altered affinity for one or more Fc γ Rs, by at least one biochemical based assay, e.g., an ELISA assay, the molecule maybe engineered into a complete immunoglobulin, using standard recombinant DNA technology methods known in the art, and the immunoglobulin comprising the variant Fc region expressed in mammalian cells for further biochemical characterization. The immunoglobulin into which a variant Fc region of the invention is introduced (e.g., replacing the Fc region of the immunoglobulin)

- can be any immunoglobulin including, but not limited to, polyclonal antibodies, monoclonal antibodies, bispecific antibodies, multi-specific antibodies, humanized antibodies, and chimeric antibodies. In preferred embodiments, a variant Fc region is introduced into an immunoglobulin specific for a cell surface receptor, a tumor antigen, or a cancer antigen.
- 5 The immunoglobulin into which a variant Fc region of the invention is introduced may specifically bind a cancer or tumor antigen for example, including, but not limited to, KS 1/4 pan-carcinoma antigen (Perez and Walker, 1990, *J. Immunol.* 142: 3662-3667; Bumal, 1988, *Hybridoma* 7(4): 407-415), ovarian carcinoma antigen (CA125) (Yu *et al.*, 1991, *Cancer Res.* 51(2): 468-475), prostatic acid phosphate (Tailor *et al.*, 1990, *Nucl. Acids Res.* 18(16): 4928), prostate specific antigen (Henttu and Vihko, 1989, *Biochem. Biophys. Res. Comm.* 160(2): 903-910; Israeli *et al.*, 1993, *Cancer Res.* 53: 227-230), melanoma-associated antigen p97 (Estin *et al.*, 1989, *J. Natl. Cancer Instit.* 81(6): 445-446), melanoma antigen gp75 (Vijayasardahl *et al.*, 1990, *J. Exp. Med.* 171(4): 1375-1380), high molecular weight melanoma antigen (HMW-MAA) (Natali *et al.*, 1987, *Cancer* 59: 55-63; Mittelman
- 10 *et al.*, 1990, *J. Clin. Invest.* 86: 2136-2144), prostate specific membrane antigen, carcinoembryonic antigen (CEA) (Foon *et al.*, 1994, *Proc. Am. Soc. Clin. Oncol.* 13: 294), polymorphic epithelial mucin antigen, human milk fat globule antigen, colorectal tumor-associated antigens such as: CEA, TAG-72 (Yokata *et al.*, 1992, *Cancer Res.* 52: 3402-3408), CO17-1A (Ragnhammar *et al.*, 1993, *Int. J. Cancer* 53: 751-758); GICA 19-9
- 20 (Herlyn *et al.*, 1982, *J. Clin. Immunol.* 2: 135), CTA-1 and LEA, Burkitt's lymphoma antigen-38.13, CD19 (Ghetie *et al.*, 1994, *Blood* 83: 1329-1336), human B-lymphoma antigen-CD20 (Reff *et al.*, 1994, *Blood* 83:435-445), CD33 (Sgouros *et al.*, 1993, *J. Nucl. Med.* 34:422-430), melanoma specific antigens such as ganglioside GD2 (Saleh *et al.*, 1993, *J. Immunol.*, 151, 3390-3398), ganglioside GD3 (Shitara *et al.*, 1993, *Cancer Immunol.*
- 25 *Immunother.* 36:373-380), ganglioside GM2 (Livingston *et al.*, 1994, *J. Clin. Oncol.* 12: 1036-1044), ganglioside GM3 (Hoon *et al.*, 1993, *Cancer Res.* 53: 5244-5250), tumor-specific transplantation type of cell-surface antigen (TSTA) such as virally-induced tumor antigens including T-antigen DNA tumor viruses and Envelope antigens of RNA tumor viruses, oncofetal antigen-alpha-fetoprotein such as CEA of colon, bladder tumor oncofetal
- 30 antigen (Hellstrom *et al.*, 1985, *Cancer. Res.* 45:2210-2188), differentiation antigen such as human lung carcinoma antigen L6, L20 (Hellstrom *et al.*, 1986, *Cancer Res.* 46: 3917-3923), antigens of fibrosarcoma, human leukemia T cell antigen-Gp37 (Bhattacharya-Chatterjee *et al.*, 1988, *J. of Immun.* 141:1398-1403), neoglycoprotein, sphingolipids, breast cancer antigen such as EGFR (Epidermal growth factor receptor), HER2 antigen
- 35 (p185^{HER2}), polymorphic epithelial mucin (PEM) (Hilkens *et al.*, 1992, *Trends in Bio.*

Chem. Sci. 17:359), malignant human lymphocyte antigen-APO-1 (Bernhard *et al.*, 1989, *Science* 245: 301-304), differentiation antigen (Feizi, 1985, *Nature* 314: 53-57) such as I antigen found in fetal erythrocytes, primary endoderm I antigen found in adult erythrocytes, preimplantation embryos, I(Ma) found in gastric adenocarcinomas, M18, M39 found in breast epithelium, SSEA-1 found in myeloid cells, VEP8, VEP9, Myl, VIM-D5, D₁₅₆₋₂₂ found in colorectal cancer, TRA-1-85 (blood group H), C14 found in colonic adenocarcinoma, F3 found in lung adenocarcinoma, AH6 found in gastric cancer, Y hapten, Le^y found in embryonal carcinoma cells, TL5 (blood group A), EGF receptor found in A431 cells, E₁ series (blood group B) found in pancreatic cancer, FC10.2 found in embryonal carcinoma cells, gastric adenocarcinoma antigen, CO-514 (blood group Le^a) found in Adenocarcinoma, NS-10 found in adenocarcinomas, CO-43 (blood group Le^b), G49 found in EGF receptor of A431 cells, MH2 (blood group ALe^b/Le^y) found in colonic adenocarcinoma, 19.9 found in colon cancer, gastric cancer mucins, T_{5A7} found in myeloid cells, R₂₄ found in melanoma, 4.2, G_{D3}, D1.1, OFA-1, G_{M2}, OFA-2, G_{D2}, and M1:22:25:8 found in embryonal carcinoma cells, and SSEA-3 and SSEA-4 found in 4 to 8-cell stage embryos. In one embodiment, the antigen is a T cell receptor derived peptide from a Cutaneous Tcell Lymphoma (*see*, Edelson, 1998, *The Cancer Journal* 4:62).

[00164] In some embodiments, a variant Fc region of the invention is introduced into an anti-fluoresceine monoclonal antibody, 4-4-20 (Kranz *et al.*, 1982 *J. Biol. Chem.* 257(12): 6987-6995; which is incorporated herein by reference in its entirety). In other embodiments, a variant Fc region of the invention is introduced into a mouse-human chimeric anti-CD20 monoclonal antibody 2H7, which recognizes the CD20 cell surface phosphoprotein on B cells (Liu *et al.*, 1987, *Journal of Immunology*, 139: 3521-6; which is incorporated herein by reference in its entirety). In yet other embodiments, a variant Fc region of the invention is introduced into a humanized antibody (Ab4D5) against the human epidermal growth factor receptor 2 (p185 HER2) as described by Carter *et al.* (1992, *Proc. Natl. Acad. Sci. USA* 89: 4285-9; which is incorporated herein by reference in its entirety). In yet other embodiments, a variant Fc region of the invention is introduced into a humanized anti-TAG72 antibody (CC49) (Sha *et al.*, 1994 *Cancer Biother.* 9(4): 341-9). In other embodiments, a variant Fc region of the invention is introduced into Rituxan which is used for treating lymphomas.

[00165] In another specific embodiment, the invention encompasses engineering an anti-FcγRIIB antibody including but not limited to any of the antibodies disclosed in U.S. Provisional Application No. 60/403,266 filed on August 12, 2002 and U.S. Application No. ___ filed on August 14, 2003, having Attorney Docket No. 011183-010-999, by

modification (*e.g.*, substitution, insertion, deletion) of at least one amino acid residue which modification increases the affinity of the Fc region for Fc γ RIIA and/or Fc γ RIIA. Examples of anti-Fc γ RIIB antibodies that may be engineered in accordance with the methods of the invention are 2B6 monoclonal antibody having ATCC accession number PTA-4591 and

5 3H7 having ATCC accession number PTA-4592. In another specific embodiment, modification of the anti-Fc γ RIIB antibody may also further decrease the affinity of the Fc region for Fc γ RIIB. In yet another specific embodiment, the engineered anti-Fc γ RIIB antibody may further have an enhanced effector function as determined by standard assays known in the art and disclosed and exemplified herein. In some embodiments, a variant Fc

10 region of the invention is introduced into a therapeutic monoclonal antibody specific for a cancer antigen or cell surface receptor including but not limited to, ErbituxTM (also known as IMC-C225) (ImClone Systems Inc.), a chimerized monoclonal antibody against EGFR; HERCEPTIN[®] (Trastuzumab) (Genentech, CA) which is a humanized anti-HER2 monoclonal antibody for the treatment of patients with metastatic breast cancer; REOPRO[®]

15 (abciximab) (Centocor) which is an anti-glycoprotein IIb/IIIa receptor on the platelets for the prevention of clot formation; ZENAPAX[®] (daclizumab) (Roche Pharmaceuticals, Switzerland) which is an immunosuppressive, humanized anti-CD25 monoclonal antibody for the prevention of acute renal allograft rejection. Other examples are a humanized anti-CD18 F(ab')₂ (Genentech); CDP860 which is a humanized anti-CD18 F(ab')₂ (Celltech,

20 UK); PRO542 which is an anti-HIV gp120 antibody fused with CD4 (Progenics/Genzyme Transgenics); C14 which is an anti-CD14 antibody (ICOS Pharm); a humanized anti-VEGF IgG1 antibody (Genentech); OVAREXTM which is a murine anti-CA 125 antibody (Altarex); PANOREXTM which is a murine anti-17-IA cell surface antigen IgG2a antibody (Glaxo Wellcome/Centocor); IMC-C225 which is a chimeric anti-EGFR IgG antibody

25 (ImClone System); VITAXINTM which is a humanized anti- α V β 3 integrin antibody (Applied Molecular Evolution/MedImmune); Campath 1H/LDP-03 which is a humanized anti CD52 IgG1 antibody (Leukosite); Smart M195 which is a humanized anti-CD33 IgG antibody (Protein Design Lab/Kanebo); RITUXANTM which is a chimeric anti-CD20 IgG1 antibody (IDEC Pharm/Genentech, Roche/Zettyaku); LYMPHOCIDETM which is a

30 humanized anti-CD22 IgG antibody (Immunomedics); Smart ID10 which is a humanized anti-HLA antibody (Protein Design Lab); ONCOLYMTM (Lym-1) is a radiolabelled murine anti-HLA DR antibody (Techniclone); anti-CD11a is a humanized IgG1 antibody (Genetech/Xoma); ICM3 is a humanized anti-ICAM3 antibody (ICOS Pharm); IDEC-114 is a primatized anti-CD80 antibody (IDEC Pharm/Mitsubishi); ZEVALINTM is a radiolabelled

murine anti-CD20 antibody (IDEC/Schering AG); IDEC-131 is a humanized anti-CD40L antibody (IDEC/Eisai); IDEC-151 is a primatized anti-CD4 antibody (IDEC); IDEC-152 is a primatized anti-CD23 antibody (IDEC/Seikagaku); SMART anti-CD3 is a humanized anti-CD3 IgG (Protein Design Lab); 5G1.1 is a humanized anti-complement factor 5 (C5) antibody (Alexion Pharm); IDEC-151 is a primatized anti-CD4 IgG1 antibody (IDEC Pharm/SmithKline Beecham); MDX-CD4 is a human anti-CD4 IgG antibody (Medarex/Eisai/Genmab); CDP571 is a humanized anti-TNF- α IgG4 antibody (Celltech); LDP-02 is a humanized anti- $\alpha 4\beta 7$ antibody (LeukoSite/Genentech); OrthoClone OKT4A is a humanized anti-CD4 IgG antibody (Ortho Biotech); ANTOVA™ is a humanized anti-CD40L IgG antibody (Biogen); ANTEGREN™ is a humanized anti-VLA-4 IgG antibody (Elan); MDX-33 is a human anti-CD64 (Fc γ R) antibody (Medarex/Centcon); rhuMab-E25 is a humanized anti-IgE IgG1 antibody (Genentech/Norartis/Tanox Biosystems); IDEC-152 is a primatized anti-CD23 antibody (IDEC Pharm); ABX-CBL is a murine anti CD-147 IgM antibody (Abgenix); BTI-322 is a rat anti-CD2 IgG antibody (Medimmune/Bio Transplant); Orthoclone/OKT3 is a murine anti-CD3 IgG2a antibody (ortho Biotech); SIMULECT™ is a chimeric anti-CD25 IgG1 antibody (Novartis Pharm); LDP-01 is a humanized anti- β_2 -integrin IgG antibody (LeukoSite); Anti-LFA-1 is a murine anti CD18 F(ab')₂ (Pasteur-Merieux/Immunotech); CAT-152 is a human anti-TGF- β_2 antibody (Cambridge Ab Tech); and Corsevin M is a chimeric anti-Factor VII antibody (Centacor).

[00166] The variant Fc regions of the invention, preferably in the context of an immunoglobulin, can be further characterized using one or more biochemical assays and/or one or more functional assays, preferably in a high throughput manner. In some alternate embodiments, the variant Fc regions of the inventions are not introduced into an immunoglobulin and are further characterized using one or more biochemical based assays and/or one or more functional assays, preferably in a high throughput manner. The one or more biochemical assays can be any assay known in the art for identifying Fc-Fc γ R interactions, including, but not limited to, an ELISA assay, and surface plasmon resonance-based assay for determining the kinetic parameters of Fc-Fc γ R interaction, *e.g.*, BIAcore assay. The one or more functional assays can be any assay known in the art for characterizing one or more Fc γ R mediated effector cell function as known to one skilled in the art or described herein. In specific embodiments, the immunoglobulins comprising the variant Fc regions are assayed in an ELISA assay for binding to one or more Fc γ Rs, *e.g.*, Fc γ RIIA, Fc γ RIIA, Fc γ RIIA; followed by one or more ADCC assays. In some embodiments, the immunoglobulins comprising the variant Fc regions are assayed further

using a surface plasmon resonance-based assay, *e.g.*, BIAcore. Surface plasmon resonance-based assays are well known in the art, and are further discussed in Section 5.2.7, and exemplified herein in Example 6.8.

[00167] An exemplary high throughput assay for characterizing immunoglobulins comprising variant Fc regions may comprise: introducing a variant Fc region of the invention, *e.g.*, by standard recombinant DNA technology methods, in a 4-4-20 antibody; characterizing the specific binding of the 4-4-20 antibody comprising the variant Fc region to an Fc γ R (*e.g.*, Fc γ RIIA, Fc γ RIIB) in an ELISA assay; characterizing the 4-4-20 antibody comprising the variant Fc region in an ADCC assay (using methods disclosed herein) wherein the target cells are opsonized with the 4-4-20 antibody comprising the variant Fc region; the variant Fc region may then be cloned into a second immunoglobulin, *e.g.*, 4D5, 2H7, and that second immunoglobulin characterized in an ADCC assay, wherein the target cells are opsonized with the second antibody comprising the variant Fc region. The second antibody comprising the variant Fc region is then further analyzed using an ELISA-based assay to confirm the specific binding to an Fc γ R.

[00168] Preferably, a variant Fc region of the invention binds Fc γ RIIA and/or Fc γ RIIA with a higher affinity than a wild type Fc region as determined in an ELISA assay. Most preferably, a variant Fc region of the invention binds Fc γ RIIA and/or Fc γ RIIA with a higher affinity and binds Fc γ RIIB with a lower affinity than a wild type Fc region as determined in an ELISA assay. In some embodiments, the variant Fc region binds Fc γ RIIA and/or Fc γ RIIA with at least 2-fold higher, at least 4-fold higher, more preferably at least 6-fold higher, most preferably at least 8 to 10-fold higher affinity than a wild type Fc region binds Fc γ RIIA and/or Fc γ RIIA and binds Fc γ RIIB with at least 2-fold lower, at least 4-fold lower, more preferably at least 6-fold lower, most preferably at least 8 to 10-fold lower affinity than a wild type Fc region binds Fc γ RIIB as determined in an ELISA assay.

[00169] The immunoglobulin comprising the variant Fc regions may be analyzed at any point using a surface plasmon based resonance based assay, *e.g.*, BIAcore, for defining the kinetic parameters of the Fc-Fc γ R interaction, using methods disclosed herein and known to those of skill in the art. Preferably, the K_d of a variant Fc region of the invention for binding to a monomeric Fc γ RIIA and/or Fc γ RIIA as determined by BIAcore analysis is about 100 nM, preferably about 70 nM, most preferably about 40 nM.; and the K_d of the variant Fc region of the invention for binding a dimeric Fc γ RIIB is about 80 nM, about 100 nM, more preferably about 200 nM.

[00170] In most preferred embodiments, the immunoglobulin comprising the variant Fc regions is further characterized in an animal model for interaction with an FcγR.

Preferred animal models for use in the methods of the invention are, for example, transgenic mice expressing human FcγRs, *e.g.*, any mouse model described in U.S. Patent No.

- 5 5,877,397, which is incorporated herein by reference in its entirety. Transgenic mice for use in the methods of the invention include, but are not limited to, nude knockout FcγRIIIA mice carrying human FcγRIIIA; nude knockout FcγRIIIA mice carrying human FcγRIIA; nude knockout FcγRIIIA mice carrying human FcγRIIB and human FcγRIIIA; nude knockout FcγRIIIA mice carrying human FcγRIIB and human FcγRIIA.

10 5.2.1 FcγR-Fc BINDING ASSAY

- [00171] An FcγR-Fc binding assay was developed for determining the binding of the molecules of the invention comprising variant Fc regions to FcγR, which allowed detection and quantitation of the interaction, despite the inherently weak affinity of the receptor for its ligand, *e.g.*, in the micromolar range for FcγRIIB and FcγRIIIA. The method involves the
- 15 formation of an FcγR complex that has an improved avidity for an Fc region, relative to an uncomplexed FcγR. According to the invention, the preferred molecular complex is a tetrameric immune complex, comprising: (a) the soluble region of FcγR (*e.g.*, the soluble region of FcγRIIIA, FcγRIIA or FcγRIIB); (b) a biotinylated 15 amino acid AVITAG sequence (AVITAG) operably linked to the C-terminus of the soluble region of FcγR (*e.g.*,
- 20 the soluble region of FcγRIIIA, FcγRIIA or FcγRIIB); and (c) streptavidin-phycoerythrin (SA-PE); in a molar ratio to form a tetrameric FcγR complex (preferably in a 5:1 molar ratio). According to a preferred embodiment of the invention, the fusion protein is biotinylated enzymatically, using for example, the *E.coli* Bir A enzyme, a biotin ligase which specifically biotinylates a lysine residue in the 15 amino acid AVITAG sequence. In
- 25 a specific embodiment of the invention, 85% of the fusion protein is biotinylated, as determined by standard methods known to those skilled in the art, including but not limited to streptavidin shift assay. According to preferred embodiments of the invention, the biotinylated soluble FcγR proteins are mixed with SA-PE in a 1X SA-PE:5X biotinylated soluble FcγR molar ratio to form a tetrameric FcγR complex.

- 30 [00172] In a preferred embodiment of the invention, polypeptides comprising Fc regions bind the tetrameric FcγR complexes, formed according to the methods of the invention, with at least an 8-fold higher affinity than the monomeric uncomplexed FcγR. The binding of polypeptides comprising Fc regions to the tetrameric FcγR complexes may be determined using standard techniques known to those skilled in the art, such as for

example, fluorescence activated cell sorting (FACS), radioimmunoassays, ELISA assays, etc.

[00173] The invention encompasses the use of the immune complexes formed according to the methods described above, for determining the functionality of molecules comprising an Fc region in cell-based or cell-free assays.

[00174] As a matter of convenience, the reagents may be provided in an assay kit, i.e., a packaged combination of reagents for assaying the ability of molecules comprising variant Fc regions to bind Fc γ R tetrameric complexes. Other forms of molecular complexes for use in determining Fc-Fc γ R interactions are also contemplated for use in the methods of the invention, e.g., fusion proteins formed as described in U.S. Provisional Application 60/439,709, filed on January 13, 2003 (Attorney Docket No. 11183-005-888); which is incorporated herein by reference in its entirety.

5.2.2 MUTAGENESIS AND CONSTRUCTION OF YEAST DISPLAY LIBRARIES

[00175] An initial library of molecules comprising variant Fc regions is produced using any random based mutagenesis techniques known in the art. It will be appreciated by one of skill in the art that amino acid sequence variants of Fc regions may be obtained by any mutagenesis technique known to those skilled in the art. Some of these techniques are briefly described herein, however, it will be recognized that alternative procedures may produce an equivalent result. In a preferred embodiment molecules of the invention comprising variant Fc regions are prepared by error-prone PCR as exemplified in Example 6, *infra* (See Leung *et al.*, 1989, *Technique*, 1:11). It is especially preferred to have error rates of 2-3 bp/Kb for use in the methods of the invention. In one embodiment, using error prone PCR a mutation frequency of 2-3 mutations/kb is obtained.

[00176] Mutagenesis may be performed in accordance with any of the techniques known in the art including, but not limited to, synthesizing an oligonucleotide having one or more modifications within the sequence of the Fc region of an antibody or a polypeptide comprising an Fc region (e.g., the CH2 or CH3 domain) to be modified. Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences which encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Typically, a primer of about 30 to about 45 nucleotides or more in length is preferred, with about 10 to about 25 or more residues on both sides of the junction of the

sequence being altered. A number of such primers introducing a variety of different mutations at one or more positions may be used to generate a library of mutants.

[00177] The technique of site-specific mutagenesis is well known in the art, as exemplified by various publications (*see, e.g., Kunkel et al., Methods Enzymol.*, 154:367-82, 1987, which is hereby incorporated by reference in its entirety). In general, site-directed mutagenesis is performed by first obtaining a single-stranded vector or melting apart of two strands of a double stranded vector which includes within its sequence a DNA sequence which encodes the desired peptide. An oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically. This primer is then annealed with the single-stranded vector, and subjected to DNA polymerizing enzymes such as T7 DNA polymerase, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform or transfect appropriate cells, such as *E. coli* cells, and clones are selected which include recombinant vectors bearing the mutated sequence arrangement. As will be appreciated, the technique typically employs a phage vector which exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage. These phage are readily commercially available and their use is generally well known to those skilled in the art. Double stranded plasmids are also routinely employed in site directed mutagenesis which eliminates the step of transferring the gene of interest from a plasmid to a phage.

[00178] Alternatively, the use of PCRTM with commercially available thermostable enzymes such as *Taq* DNA polymerase may be used to incorporate a mutagenic oligonucleotide primer into an amplified DNA fragment that can then be cloned into an appropriate cloning or expression vector. *See, e.g., Tomic et al., Nucleic Acids Res.*, 18(6):1656, 1987, and Upender *et al., Biotechniques*, 18(1):29-30, 32, 1995, for PCRTM - mediated mutagenesis procedures, which are hereby incorporated in their entireties. PCRTM employing a thermostable ligase in addition to a thermostable polymerase may also be used to incorporate a phosphorylated mutagenic oligonucleotide into an amplified DNA fragment that may then be cloned into an appropriate cloning or expression vector (*see e.g., Michael, Biotechniques*, 16(3):410-2, 1994, which is hereby incorporated by reference in its entirety).

[00179] Another method for preparing variants for use in the invention, is cassette mutagenesis based on the technique described by Wells *et al.* (1985, *Gene*, 34: 315). The starting material is the plasmid comprising the desired DNA encoding the protein to be mutated (*e.g., the DNA encoding a polypeptide comprising an Fc region*). The codon(s) in

- the DNA sequence to be mutated are identified; there must be a unique restriction endonuclease site on each side of the identified mutations site(s). If no such restriction site exists, it may be generated by oligonucleotide directed mutagenesis. After the restriction sites have been introduced into the plasmid, the plasmid is cut at these sites and linearized.
- 5 A double-stranded oligonucleotide encoding the sequence of the DNA between the restriction sites but containing the mutation is synthesized using standard procedures known to those skilled in the art. The double stranded oligonucleotide is referred to as the cassette. This cassette is designed to have 3' and 5' ends that are compatible with the ends of the linearized plasmid, such that it can be directly ligated to the plasmid.
- 10 [00180] Other methods known to those of skill in the art for producing sequence variants of the Fc region of an antibody or polypeptides comprising an Fc region can be used. For example, recombinant vectors encoding the amino acid sequence of the constant domain of an antibody or a fragment thereof may be treated with mutagenic agents, such as hydroxylamine, to obtain sequence variants.
- 15 [00181] Once a mutant library is produced according to the methods described, the mutagenized library is transformed into a yeast strain, preferably EBY100 (Invitrogen), *MATa ura3-52 trp1 leu2Δ1 his3Δ200 pep4::HIS3 prb1Δ1.6R can1 GAL::GAL-AGA1* using a standard lithium acetate transformation protocol known to those skilled in the art (ref).
- [00182] It will be appreciated by one of skill in the art, that once molecules of the
- 20 invention with desired binding properties (e.g., molecules with variant Fc regions with at least one amino acid modification, which modification enhances the affinity of the variant Fc region for FcγRIIIA relative to a comparable molecule, comprising a wild-type Fc region) have been identified (See Section 5.1 and Table 2) according to the methods of the invention, other molecules (i.e., therapeutic antibodies) may be engineered using standard
- 25 recombinant DNA techniques and any known mutagenesis techniques, as described in this section to produce engineered molecules carrying the identified mutation sites.

5.2.3 YEAST SURFACE DISPLAY

- [00183] The preferred method for screening and identifying molecules comprising variant Fc regions with altered FcγR affinities (i.e., enhanced FcγRIIIA affinity and/or
- 30 FcγRIIA) is yeast surface display technology (for review see Boder and Wittrup, 2000, *Methods in Enzymology*, 328: 430-444, which is incorporated herein by reference in its entirety) which addresses the deficiency in the prior art for screening binding interactions of extracellular post-translationally modified proteins. Specifically, the yeast surface display is a genetic method whereby polypeptides comprising Fc mutants are expressed on the yeast

cell wall in a form accessible for interacting with Fc γ R. Yeast surface display of the mutant Fc containing polypeptides of the invention may be performed in accordance with any of the techniques known to those skilled in the art. See U.S. Patent No.'s 6,423,538; 6,114,147; and 6,300,065, all of which are incorporated herein by reference in their entirety.

- 5 See Boder *et al.*, 1997 *Nat. Biotechnol.*, 15:553-7; Boder *et al.*, 1998 *Biotechnol. Prog.*, 14:55-62; Boder *et al.*, 2000 *Methods Enzymol.*, 328:430-44; Boder *et al.*, 2000 *Proc. Natl. Acad. Sci. U.S.A.*, 2000, 97:10701-5; Shusta *et al.*, 1998 *Nat. Biotechnol.*, 1998, 16:773-7; Shusta *et al.*, 1999 *J. Mol. Biol.*, 292:949-56; Shusta *et al.*, 1999 *Curr. Opin. Biotechnol.*, 10:117-22; Shusta *et al.*, 2000 *Nat. Biotechnol.*, 18:754-9; Wittrup *et al.*, 1994 *Ann. N.Y. Acad. Sci.*, 745:321-30; Wittrup *et al.*, 1994 *Cytometry*, 16:206-13; Wittrup, 1995 *Curr. Opin. Biotechnol.*, 6:203-8; Wittrup, 1999 *Trends Biotechnol.*, 17:423-4; Wittrup, 2000 *Nat. Biotechnol.*, 18:1039-40; Wittrup, 2001 *Curr. Opin. Biotechnol.*, 12:395-9.

- [00184] The invention provides methods for constructing an Fc mutant library in yeast for displaying molecules comprising Fc regions, which have been mutated as
- 15 described in Section 5.2.2. Preferably, the Fc mutant libraries for use in the methods of the invention contain at least 10^7 cells, up to 10^9 cells. One exemplary method for constructing a Fc library for use in the methods of the invention comprises the following: nucleic acids encoding molecules comprising Fc regions are cloned into the multiple cloning site of a vector derived from a yeast replicating vector, *e.g.*, pCT302; such that the Fc encoding
- 20 nucleic acids are expressed under the control of the GAL1 galactose-inducible promoter and in-frame with a nucleotide sequence encoding Aga2p, the mating agglutinin cell wall protein. In a preferred embodiment, nucleic acids encoding molecules comprising Fc regions are cloned C-terminal to the Aga2p coding region, such that a Fc-region Aga2p fusion protein is encoded. A fusion protein comprising the Aga2p protein and polypeptides
- 25 comprising Fc regions will be secreted extracellularly and displayed on the cell wall via disulfide linkage to the Aga1p protein, an integral cell wall protein, using the preferred construct of the invention. In an alternative embodiment, the constructs may further comprise nucleotide sequences encoding epitope tags. Any epitope tag nucleotide coding sequence known to those skilled in the art can be used in accordance with the invention,
- 30 including, but not limited to nucleotide sequences encoding hemagglutinin (HA), c-myc Xpress TAG, His - TAG, or V5TAG. The presence of the fusion protein on the yeast cell surface may be detected using FACS analysis, confocal fluorescence microscopy or standard immunostaining methods, all of which are known to those skilled in the art. In one embodiment, the presence of the Fc fusion proteins of the invention on the yeast cell surface
- 35 are detected using Fc-specific monoclonal antibodies (CH3 specific), including but not

limited to IgG1 Fc-specific monoclonal antibody, HP6017 (Sigma), JL512 (Immunotech), and any antibody disclosed in Partridge *et al.*, 1986, *Molecular Immunology*, 23 (12): 1365-72, which is incorporated herein by reference in its entirety. In another embodiment, the presence of the Fc fusion proteins of the invention are detected by immunofluorescent labeling of epitope tags using techniques known to those skilled in the art. It is particularly useful in the methods of the invention, to use nucleotide sequences encoding epitope tags to flank the nucleic acids encoding the Fc fusion proteins, as an internal control, to detect if the fusion proteins are displayed on the cell wall in a partially proteolyzed form.

5.2.4 SCREENING OF YEAST DISPLAY LIBRARIES

10 [00185] The invention encompasses screening the yeast display libraries using immunological based assays including but not limited to cell based assays, solution based assays, and solid phase based assays.

[00186] In some embodiments, the invention encompasses identification of Fc mutants with altered Fc γ R affinities using affinity maturation methods which are known to those skilled in the art and encompassed herein. Briefly, affinity maturation creates novel alleles by randomly recombining individual mutations present in a mutant library, *see, e.g.*, Hawkins *et al.*, 1992, *J. Mol. Biol.* 226: 889-896; Stemmer *et al.*, 1994 *Nature*, 370: 389-91; both of which are incorporated herein by reference in their entireties. It has been used successfully to increase the affinity of antibodies, T cell receptors and other proteins. The invention encompasses using mutations that show increased Fc γ R binding as a baseline to construct new mutant libraries with enhanced phenotypes. Using the methods of the invention, a population of IgG1 Fc mutants enriched by yeast surface display for increased binding to an Fc γ R, *e.g.*, Fc γ RIIIA, may be selected. Following DNA preparation, Fc regions can be amplified by PCR using flanking primers that selectively amplify the mutated region of the Fc, which is about ~700 bp using methods known to one skilled in the art and exemplified or disclosed herein. Novel mutants can thus be constructed by reshuffling of mutations in the Fc region for example via DNaseI treatment of the amplified DNA and isolation of fragments using methods such as those disclosed by Stemmer *et al.*, 1994 *Proc. Natl. Acad. Sci. USA* 91: 10747-51, which is incorporated herein by reference in its entirety. Fragments can then be religated, PCR amplified with nested primers and cloned into the yeast display vector, *e.g.*, pYD1 using methods known to one skilled in the art. The recombined library can then be reselected in the yeast Fc display screen. As the K_D decreases, below 10 nM, conditions can be established to allow for further increases in affinity based on the reduction of the off rate of the Fc γ RIIIA ligand from the Fc receptor

using methods known in the art such as those disclosed in Boder *et al.*, 1998, *Biotechnol. Prog.* 14: 55-62, which is incorporated herein by reference in its entirety. The invention encompasses a kinetic screen of the yeast library. A kinetic screen may be established by labeling of the Fc displaying cells to saturation with a labeled ligand, e.g., a fluorescent ligand followed by incubation with an excess of non-labeled ligand for a predetermined period. After termination of the reaction by the addition of excess buffer (e.g., 1X PBS, 0.5 mg/ml BSA) cells will be analyzed by FACS and sort gates set for selection. After each round of enrichment individual mutants can be tested for fold increases in affinity and sequenced for diversity. The *in vitro* recombination process can be repeated. In some embodiments, the *in vitro* is repeated at least 3 times.

[00187] Selection of the Fc variants of the invention may be done using any FcγR including but not limited to polymorphic variants of FcγR. In some embodiments, selection of the Fc variants is done using a polymorphic variant of FcγRIIIA which contains a phenylalanine at position 158. In other embodiments, selection of the Fc variants is done using a polymorphic variant of FcγRIIIA which contains a valine at position 158. FcγRIIIA 158V displays a higher affinity for IgG1 than 158F and an increased ADCC activity (*see, e.g., Koene et al.*, 1997, *Blood*, 90:1109-14; *Wu et al.*, 1997, *J. Clin. Invest.* 100: 1059-70, both of which are incorporated herein by reference in their entireties); this residue in fact directly interacts with the lower hinge region of IgG1 as recently shown by IgG1-FcγRIIIA co-crystallization studies, *see, e.g., Sonderman et al.*, 2000, *Nature*, 100: 1059-70, which is incorporated herein by reference in its entirety. Studies have shown that in some cases therapeutic antibodies have improved efficacy in FcγRIIIA-158V homozygous patients. For example, humanized anti-CD20 monoclonal antibody Rituximab was therapeutically more effective in FcγRIIIA158V homozygous patients compared to FcγRIIIA 158F homozygous patients (*See, e.g., Cartron et al.*, 2002 *Blood*, 99(3): 754-8). Although not intending to be bound by a particular mechanism of action, selection of Fc variants of the invention with FcγRIIIA 158F allotype may provide for variants that once engineered into therapeutic antibodies will be clinically more efficacious for FcγRIIIA 158F homozygous patients.

[00188] The invention encompasses screening yeast libraries based on FcγRIIB depletion and FcγRIIIA selection so that Fc mutants are selected that not only have an enhanced affinity for FcγRIIIA but also have a reduced affinity for FcγRIIB. Yeast libraries may be enriched for clones that have a reduced affinity for FcγRIIB by sequential depletion methods, for example, by incubating the yeast library with magnetic beads coated with FcγRIIB. FcγRIIB depletion is preferably carried out sequentially so that the library is

enriched in clones that have a reduced affinity for Fc γ RIIB. In some embodiments, the Fc γ RIIB depletion step results in a population of cells so that only 30% , preferably only 10%, more preferably only 5%, most preferably less than 1% bind Fc γ RIIB. In some
5 6 cycles. The Fc γ RIIB depletion step is preferably combined with an Fc γ RIIIA selection step, for example using FACS sorting so that Fc variants with an enhanced affinity for Fc γ RIIIA are selected.

5.2.4.1 FACS ASSAYS; SOLID PHASED ASSAYS AND IMMUNOLOGICAL BASED ASSAYS

10 [00189] The invention encompasses characterization of the mutant Fc fusion proteins that are displayed on the yeast surface cell wall, according to the methods described in Section 5.2.3. One aspect of the invention provides a method for selecting mutant Fc fusion proteins with a desirable binding property, specifically, the ability of the mutant Fc fusion
15 polypeptide comprising a wild-type Fc region binds Fc γ RIIIA and/or Fc γ RIIA. In another embodiment, the invention provides a method for selecting mutant Fc fusion proteins with a desirable binding property, specifically, the ability of the mutant Fc fusion protein to bind Fc γ RIIIA and/or Fc γ RIIA with a greater affinity than a comparable polypeptide comprising
20 Fc fusion protein to bind Fc γ RIIB with a lower affinity than a comparable polypeptide comprising a wild-type Fc region binds Fc γ RIIB. It will be appreciated by one skilled in the art, that the methods of the invention can be used for identifying and screening any mutations in the Fc regions of molecules, with any desired binding characteristic.

[00190] Yeast cells displaying the mutant Fc fusion proteins can be screened and
25 characterized by any biochemical or immunological based assays known to those skilled in the art for assessing binding interactions.

[00191] Preferably, fluorescence activated cell sorting (FACS), using any of the techniques known to those skilled in the art, is used for screening the mutant Fc fusion proteins displayed on the yeast cell surface for binding Fc γ RIIIA, preferably the Fc γ RIIIA
30 tetrameric complex, or optionally Fc γ RIIB. Flow sorters are capable of rapidly examining a large number of individual cells that contain library inserts (e.g., 10-100 million cells per hour) (Shapiro *et al.*, *Practical Flow Cytometry*, 1995). Additionally, specific parameters used for optimization including, but not limited to, ligand concentration (i.e., Fc γ RIIIA tetrameric complex), kinetic competition time, or FACS stringency may be varied in order

to select for the cells which display Fc fusion proteins with specific binding properties, *e.g.*, higher affinity for FcγRIIIA compared to a comparable polypeptide comprising a wild-type Fc region. Flow cytometers for sorting and examining biological cells are well known in the art. Known flow cytometers are described, for example, in U.S. Patent Nos. 4,347,935; 5,464,581; 5,483,469; 5,602,039; 5,643,796; and 6,211,477; the entire contents of which are
5 incorporated by reference herein. Other known flow cytometers are the FACS Vantage™ system manufactured by Becton Dickinson and Company, and the COPAS™ system manufactured by Union Biometrica.

[00192] According to a preferred embodiment of the invention, yeast cells are
10 analyzed by fluorescence activated cell sorting (FACS). In most preferred embodiments, the FACS analysis of the yeast cells is done in an iterative manner, at least twice, at least three times, or at least 5 times. Between each round of selection cells are regrown and induced so the Fc regions are displayed on the maximum number of yeast cell surfaces. Although not intending to be bound by a particular mode of action, this iterative process
15 helps enrich the population of the cells with a particular phenotype, *e.g.*, high binding to FcγRIIIA.

[00193] In preferred embodiments, screening for Fc variants of the invention comprises a selection process that has multiple rounds of screening, *e.g.*, at least two rounds of screening. In one embodiment, screening for Fc variants that have an enhanced affinity
20 for FcγRIIIA may comprise the following steps: in the first round of screening, a library of yeast cells, *e.g.*, a naïve library of 10^7 cells is enriched by FACS, preferably in an iterative manner, using for example labeled tetrameric FcγRIIIA to select for Fc variants that have an enhanced affinity for FcγRIIIA; the variant Fc region that is selected with the desired phenotype, *e.g.*, enhanced binding to FcγRIIIA, is then introduced into an antibody, *e.g.*, a
25 4-4-20 antibody, and the engineered antibody is assayed using a secondary screen, *e.g.*, ELISA for binding to an FcγR. In the second round of screening, a single mutation library may be generated based on the first screen so that the Fc region harbors the variant displaying the enhanced affinity for FcγRIIIA; and enriched by FACS using for example labeled monomeric FcγRIIIA in both the presence and absence of unlabeled receptor; and
30 the variant Fc region is then introduced into an antibody, *e.g.*, a 4-4-20 antibody, and the engineered antibody is assayed using a secondary screen, *e.g.*, ELISA for binding to an FcγR. In some embodiments, the secondary screen may further comprise characterizing the antibodies comprising Fc variants in an ADCC or BIAcore based assay using methods disclosed herein

[00194] The invention encompasses FACS screening of the mutant yeast library under equilibrium or kinetic conditions. When the screening is performed under equilibrium conditions, an excess of the yeast library carrying Fc mutants is incubated with Fc γ RIIIA, preferably labeled Fc γ RIIIA at a concentration 5-10 fold below the K_d, for at least one hour to allow binding of Fc mutants to Fc γ RIIIA under equilibrium conditions. When the screening is performed under kinetic conditions, the mutant yeast library is incubated with labeled Fc γ RIIIA; the cells are then incubated with equimolar unlabeled Fc γ RIIIA for a pre-selected time, bound Fc γ RIIIA is then monitored.

[00195] One exemplary method of analyzing the yeast cells expressing mutant Fc fusion proteins with FACS is containing the cells with Fc γ RIIIA-tetrameric complex which has been labeled with a fluorescent label such as, PE and an anti-Fc antibody, such as F(ab)₂ anti-Fc which has been fluorescently labeled. Fluorescence measurements of a yeast library produced according to the methods of the invention preferably involves comparisons with controls; for example, yeast cells that lack the insert encoding molecules comprising an Fc region (negative control). The flow sorter has the ability not only to measure fluorescence signals in cells at a rapid rate, but also to collect cells that have specified fluorescent properties. This feature may be employed in a preferred embodiment of the invention to enrich the initial library population for cells expressing Fc fusion proteins with specific binding characteristics, *e.g.*, higher affinity for Fc γ RIIIA compared to a comparable polypeptide comprising a wild-type Fc region. In a preferred embodiment of the invention, yeast cells are analyzed by FACS and sort gates established to select for cells that show the highest affinity for Fc γ RIIIA relative to the amount of Fc expression on the yeast cell surface. According to a preferred embodiment, four consecutive sorts are established, wherein the gates for each successive sort is 5.5%, 1%, 0.2%, and 0.1%. It is preferred that the yeast display library formed according to the methods of the invention be over-sampled by at least 10-fold to improve the probability of isolating rare clones (*e.g.*, analyze $\sim 10^8$ cells from a library of 10^7 clones). Alternatively, 2-5 sorts are established to select for cells of the desired phenotype. Sort gates can be established empirically by one skilled in the art.

[00196] In other preferred embodiments, mutant Fc fusion proteins displayed on the yeast cell surface are screened using solid phase based assays, for example assays using magnetic beads, *e.g.*, supplied by Dynal, preferably in a high through put manner for binding to an Fc γ R, *e.g.*, Fc γ RIIIA. In one embodiment, magnetic bead assays may be used to identify mutants with enhanced affinity for Fc γ RIIIA and/or reduced affinity for Fc γ RIIB. An exemplary assay to identify mutants with enhanced affinity for Fc γ RIIIA and reduced

- affinity for Fc γ RIIB may comprise selecting mutants by a sequential solid phase depletion using magnetic beads coated with Fc γ RIIB followed by selection with magnetic beads coated with Fc γ RIIA. For example one assay may comprise the following steps: incubating the library of yeast cells generated in accordance with the methods of the
- 5 invention with magnetic beads coated with Fc γ RIIB; separating yeast cells bound to beads from the non bound fraction by placing the mixture in a magnetic field, removing the non-bound yeast cells and placing them in a fresh media; binding the yeast cells to beads coated with Fc γ RIIA, separating yeast cells bound to beads from the non bound fraction by placing the mixture in a magnetic field, removing the non-bound yeast cells; removing the
- 10 bound cells by rigorous vortexing; growing the recovered cells in glucose containing media; re-inducing in selective media containing galactose. The selection process is repeated at least once. Inserts containing the Fc domain are then amplified using common methodologies known in the art, e.g., PCR, and introduced into an antibody by methods already described for further characterization.
- 15 [00197] In an alternative embodiment, a non-yeast based system is used to characterize the binding properties of the molecules of the invention. One exemplary system for characterizing the molecules of the invention comprises a mammalian expression vector containing the heavy chain of the anti-fluorescein monoclonal antibody 4-4-20, into which the nucleic acids encoding the molecules of the invention with variant Fc regions are
- 20 cloned. The resulting recombinant clone is expressed in a mammalian host cell line (i.e., human kidney cell line 293H), and the resulting recombinant immunoglobulin is analyzed for binding to Fc γ R using any standard assay known to those in the art, including but not limited to ELISA and FACS.
- [00198] Molecules of the present invention (e.g., antibodies, fusion proteins,
- 25 conjugated molecules) may be characterized in a variety of ways. In particular, molecules of the invention comprising modified Fc regions may be assayed for the ability to immunospecifically bind to a ligand, e.g., Fc γ RIIA tetrameric complex. Such an assay may be performed in solution (e.g., Houghten, *Bio/Techniques*, 13:412-421, 1992), on beads (Lam, *Nature*, 354:82-84, 1991, on chips (Fodor, *Nature*, 364:555-556, 1993), on bacteria
- 30 (U.S. Patent No. 5,223,409), on spores (U.S. Patent Nos. 5,571,698; 5,403,484; and 5,223,409), on plasmids (Cull *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:1865-1869, 1992) or on phage (Scott and Smith, *Science*, 249:386-390, 1990; Devlin, *Science*, 249:404-406, 1990; Cwirla *et al.*, *Proc. Natl. Acad. Sci. USA*, 87:6378-6382, 1990; and Felici, *J. Mol. Biol.*, 222:301-310, 1991) (each of these references is incorporated by reference herein in its

entirety). Molecules that have been identified to immunospecifically bind to an ligand, *e.g.*, Fc γ RIIA can then be assayed for their specificity and affinity for the ligand.

[00199] Molecules of the invention that have been engineered to comprise modified Fc regions (*e.g.*, therapeutic antibodies) or have been identified in the yeast display system to have the desired phenotype (*see* Section 5.1) may be assayed for immunospecific binding to an antigen (*e.g.*, cancer antigen and cross-reactivity with other antigens (*e.g.*, Fc γ R) by any method known in the art. Immunoassays which can be used to analyze immunospecific binding and cross-reactivity include, but are not limited to, competitive and non-competitive assay systems using techniques such as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, to name but a few. Such assays are routine and well known in the art (*see, e.g.*, Ausubel *et al.*, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York, which is incorporated by reference herein in its entirety).

[00200] The binding affinity of the molecules of the present invention comprising modified Fc regions to a ligand, *e.g.*, Fc γ R tetrameric complex and the off-rate of the interaction can be determined by competitive binding assays. One example of a competitive binding assay is a radioimmunoassay comprising the incubation of labeled ligand, such as tetrameric Fc γ R (*e.g.*, ^3H or ^{125}I) with a molecule of interest (*e.g.*, molecules of the present invention comprising modified Fc regions) in the presence of increasing amounts of unlabeled ligand, such as tetrameric Fc γ R, and the detection of the molecule bound to the labeled ligand. The affinity of the molecule of the present invention for the ligand and the binding off-rates can be determined from the saturation data by scatchard analysis.

[00201] In a preferred embodiment, BIAcore kinetic analysis is used to determine the binding on and off rates of molecules of the present invention to a ligand such as Fc γ R. BIAcore kinetic analysis comprises analyzing the binding and dissociation of a ligand from chips with immobilized molecules (*e.g.*, molecules comprising modified Fc regions) on their surface.

5.2.5 SEQUENCING OF MUTANTS

[00202] Any of a variety of sequencing reactions known in the art can be used to directly sequence the molecules of the invention comprising variant Fc regions. Examples of sequencing reactions include those based on techniques developed by Maxim and Gilbert

(*Proc. Natl. Acad. Sci. USA*, 74:560, 1977) or Sanger (*Proc. Natl. Acad. Sci. USA*, 74:5463, 1977). It is also contemplated that any of a variety of automated sequencing procedures can be utilized (*Bio/Techniques*, 19:448, 1995), including sequencing by mass spectrometry (see, e.g., PCT Publication No. WO 94/16101, Cohen *et al.*, *Adv. Chromatogr.*, 36:127-162, 1996, and Griffin *et al.*, *Appl. Biochem. Biotechnol.*, 38:147-159, 1993).

5.2.6 FUNCTIONAL ASSAYS OF MOLECULES WITH VARIANT Fc REGIONS

[00203] The invention encompasses characterization of the molecules of the invention (e.g., an antibody comprising a variant Fc region identified by the yeast display technology described *supra*; or therapeutic monoclonal antibodies engineered according to the methods of the invention) using assays known to those skilled in the art for identifying the effector cell function of the molecules. In particular, the invention encompasses characterizing the molecules of the invention for Fc γ R-mediated effector cell function. Examples of effector cell functions that can be assayed in accordance with the invention, include but are not limited to, antibody-dependent cell mediated cytotoxicity, phagocytosis, opsonization, opsonophagocytosis, C1q binding, and complement dependent cell mediated cytotoxicity. Any cell-based or cell free assay known to those skilled in the art for determining effector cell function activity can be used (For effector cell assays, see Perussia *et al.*, 2000, *Methods Mol. Biol.* 121: 179-92; Baggiolini *et al.*, 1998 *Experientia*, 44(10): 841-8; Lehmann *et al.*, 2000 *J. Immunol. Methods*, 243(1-2): 229-42; Brown EJ. 1994, *Methods Cell Biol.*, 45: 147-64; Munn *et al.*, 1990 *J. Exp. Med.*, 172: 231-237, Abdul-Majid *et al.*, 2002 *Scand. J. Immunol.* 55: 70-81; Ding *et al.*, 1998, *Immunity* 8:403-411, each of which is incorporated by reference herein in its entirety).

[00204] In one embodiment, the molecules of the invention can be assayed for Fc γ R-mediated phagocytosis in human monocytes. Alternatively, the Fc γ R-mediated phagocytosis of the molecules of the invention may be assayed in other phagocytes, e.g., neutrophils (polymorphonuclear leukocytes; PMN); human peripheral blood monocytes, monocyte-derived macrophages, which can be obtained using standard procedures known to those skilled in the art (e.g., see Brown EJ. 1994, *Methods Cell Biol.*, 45: 147-164). In one embodiment, the function of the molecules of the invention is characterized by measuring the ability of THP-1 cells to phagocytose fluoresceinated IgG-opsonized sheep red blood cells (SRBC) by methods previously described (Tridandapani *et al.*, 2000, *J. Biol. Chem.* 275: 20480-7). For example, an exemplary assay for measuring phagocytosis of the molecules of the invention comprising variant Fc regions with enhanced affinities for

Fc γ RIIIA, comprises of: treating THP-1 cells with a molecule of the invention or with a control antibody that does not bind to Fc γ RIIIA, comparing the activity levels of said cells, wherein a difference in the activities of the cells (*e.g.*, rosetting activity (the number of THP-1 cells binding IgG-coated SRBC), adherence activity (the total number of SRBC bound to THP-1 cells), and phagocytic rate) would indicate the functionality of the molecule of the invention. It can be appreciated by one skilled in the art that this exemplary assay can be used to assay any of the molecules identified by the methods of the invention.

[00205] Another exemplary assay for determining the phagocytosis of the molecules of the invention is an antibody-dependent opsonophagocytosis assay (ADCP) which can comprise the following: coating a target bioparticle such as *Escherichia coli*-labeled FITC (Molecular Probes) or *Staphylococcus aureus*-FITC with (i) wild-type 4-4-20 antibody, an antibody to fluorescein (*See* Bedzyk *et al.*, 1989, *J. Biol. Chem.*, 264(3): 1565-1569, which is incorporated herein by reference in its entirety), as the control antibody for Fc γ R-dependent ADCP; or (ii) 4-4-20 antibody harboring the D265A mutation that knocks out binding to Fc γ RIII, as a background control for Fc γ R-dependent ADCP (iii) 4-4-20 antibody carrying variant Fc regions identified by the methods of the invention and produced as exemplified in Example 6.6; and forming the opsonized particle; adding any of the opsonized particles described (i-iii) to THP-1 effector cells (a monocytic cell line available from ATCC) in a 60:1 ratio to allow Fc γ R-mediated phagocytosis to occur; preferably incubating the cells and *E. coli*-FITC/antibody at 37°C for 1.5 hour; adding trypan blue after incubation (preferably at room temperature for 2-3 min.) to the cells to quench the fluorescence of the bacteria that are adhered to the outside of the cell surface without being internalized; transferring cells into a FACS buffer (*e.g.*, 0.1%, BSA in PBS, 0.1%, sodium azide), analyzing the fluorescence of the THP1 cells using FACS (*e.g.*, BD FACS Calibur). Preferably, the THP-1 cells used in the assay are analyzed by FACS for expression of Fc γ R on the cell surface. THP-1 cells express both CD32A and CD64. CD64 is a high affinity Fc γ R that is blocked in conducting the ADCP assay in accordance with the methods of the invention. The THP-1 cells are preferably blocked with 100 μ g/mL soluble IgG1 or 10% human serum. To analyze the extent of ADCP, the gate is preferably set on THP-1 cells and median fluorescence intensity is measured. The ADCP activity for individual mutants is calculated and reported as a normalized value to the wild type chMab 4-4-20 obtained. The opsonized particles are added to THP-1 cells such that the ratio of the opsonized particles to THP-1 cells is 30:1 or 60:1. In most preferred embodiments, the ADCP assay is conducted with controls, such as *E. coli*-FITC in medium, *E. coli*-FITC and THP-1 cells (to serve as Fc γ R-independent ADCP activity), *E. coli*-FITC, THP-1 cells and

wild-type 4-4-20 antibody (to serve as FcγR-dependent ADCP activity), *E coli*-FITC, THP-1 cells, 4-4-20 D265A (to serve as the background control for FcγR-dependent ADCP activity).

[00206] In another embodiment, the molecules of the invention can be assayed for FcγR-mediated ADCC activity in effector cells, *e.g.*, natural killer cells, using any of the standard methods known to those skilled in the art (*See e.g.*, Perussia *et al.*, 2000, *Methods Mol. Biol.* 121: 179-92). An exemplary assay for determining ADCC activity of the molecules of the invention is based on a ⁵¹Cr release assay comprising of: labeling target cells with [⁵¹Cr]Na₂CrO₄ (this cell-membrane permeable molecule is commonly used for labeling since it binds cytoplasmic proteins and although spontaneously released from the cells with slow kinetics, it is released massively following target cell necrosis); opsonizing the target cells with the molecules of the invention comprising variant Fc regions; combining the opsonized radiolabeled target cells with effector cells in a microtitre plate at an appropriate ratio of target cells to effector cells; incubating the mixture of cells for 16-18 hours at 37°C; collecting supernatants; and analyzing radioactivity. The cytotoxicity of the molecules of the invention can then be determined, for example using the following formula: % lysis = (experimental cpm - target leak cpm)/(detergent lysis cpm - target leak cpm) x 100%. Alternatively, % lysis =(ADCC-AICC)/(maximum release-spontaneous release). Specific lysis can be calculated using the formula: specific lysis = % lysis with the molecules of the invention - % lysis in the absence of the molecules of the invention. A graph can be generated by varying either the target: effector cell ratio or antibody concentration.

[00207] In yet another embodiment, the molecules of the invention are characterized for antibody dependent cellular cytotoxicity (ADCC) *see, e.g.*, Ding *et al.*, *Immunity*, 1998, 8:403-11; which is incorporated herein by reference in its entirety.

[00208] Preferably, the effector cells used in the ADCC assays of the invention are peripheral blood mononuclear cells (PBMC) that are preferably purified from normal human blood, using standard methods known to one skilled in the art, *e.g.*, using Ficoll-Paque density gradient centrifugation. Preferred effector cells for use in the methods of the invention express different FcγR activating receptors. The invention encompasses, effector cells, THP-1, expressing FcγRI, FcγRIIA and FcγRIIB, and monocyte derived primary macrophages derived from whole human blood expressing both FcγRIIA and FcγRIIB, to determine if Fc antibody mutants show increased ADCC activity and phagocytosis relative to wild type IgG1 antibodies.

[00209] The human monocyte cell line, THP-1, activates phagocytosis through expression of the high affinity receptor FcγRI and the low affinity receptor FcγRIIA (Fleit et al., 1991, *J. Leuk. Biol.* 49: 556). THP-1 cells do not constitutively express FcγRIIA or FcγRIIB. Stimulation of these cells with cytokines effects the FcR expression pattern
5 (Pricop et al., 2000 *J. Immunol.* 166: 531-7). Growth of THP-1 cells in the presence of the cytokine IL4 induces FcγRIIB expression and causes a reduction in FcγRIIA and FcγRI expression. FcγRIIB expression can also be enhanced by increased cell density (Tridandapani et al., 2002, *J. Biol Chem.* 277: 5082-9). In contrast, it has been reported that IFNγ can lead to expression of FcγRIIA (Pearse et al., 1993 *PNAS USA* 90: 4314-8). The
10 presence or absence of receptors on the cell surface can be determined by FACS using common methods known to one skilled in the art. Cytokine induced expression of FcγR on the cell surface provides a system to test both activation and inhibition in the presence of FcγRIIB. If THP-1 cells are unable to express the FcγRIIB the invention also encompasses another human monocyte cell line, U937. These cells have been shown to terminally
15 differentiate into macrophages in the presence of IFNγ and TNF (Koren et al., 1979, *Nature* 279: 328-331).

[00210] FcγR dependent tumor cell killing is mediated by macrophage and NK cells in mouse tumor models (Clynes et al., 1998, *PNAS USA* 95: 652-656). The invention encompasses the use of elutriated monocytes from donors as effector cells to analyze the
20 efficiency Fc mutants to trigger cell cytotoxicity of target cells in both phagocytosis and ADCC assays. Expression patterns of FcγRI, FcγRIIA, and FcγRIIB are affected by different growth conditions. FcγR expression from frozen elutriated monocytes, fresh elutriated monocytes, monocytes maintained in 10% FBS, and monocytes cultured in FBS + GM-CSF and or in human serum may be determined using common methods known to
25 those skilled in the art. For example, cells can be stained with FcγR specific antibodies and analyzed by FACS to determine FcR profiles. Conditions that best mimic macrophage *in vivo* FcγR expression is then used for the methods of the invention.

[00211] In some embodiments, the invention encompasses the use of mouse cells especially when human cells with the right FcγR profiles are unable to be obtained. In some
30 embodiments, the invention encompasses the mouse macrophage cell line RAW264.7(ATCC) which can be transfected with human FcγRIIA and stable transfectants isolated using methods known in the art, *see, e.g.,* Ralph et al., *J. Immunol.* 119: 950-4). Transfectants can be quantitated for FcγRIIA expression by FACS analysis using routine experimentation and high expressors can be used in the ADCC assays of the invention. In

other embodiments, the invention encompasses isolation of spleen peritoneal macrophage expressing human FcγR from knockout transgenic mice such as those disclosed herein.

[00212] Lymphocytes may be harvested from peripheral blood of donors (PBM) using a Ficoll-Paque gradient (Pharmacia). Within the isolated mononuclear population of cells the majority of the ADCC activity occurs via the natural killer cells (NK) containing FcγRIIIA but not FcγRIIB on their surface. Results with these cells indicate the efficacy of the mutants on triggering NK cell ADCC and establish the reagents to test with elutriated monocytes.

[00213] Target cells used in the ADCC assays of the invention include, but are not limited to, breast cancer cell lines, *e.g.*, SK-BR-3 with ATCC accession number HTB-30 (*see, e.g.*, Tremp *et al.*, 1976, *Cancer Res.* 33-41); B-lymphocytes; cells derived from Burkitts lymphoma, *e.g.*, Raji cells with ATCC accession number CCL-86 (*see, e.g.*, Epstein *et al.*, 1965, *J. Natl. Cancer Inst.* 34: 231-240), and Daudi cells with ATCC accession number CCL-213 (*see, e.g.*, Klein *et al.*, 1968, *Cancer Res.* 28: 1300-10). The target cells must be recognized by the antigen binding site of the immunoglobulin to be assayed.

[00214] The ADCC assay is based on the ability of NK cells to mediate cell death via an apoptotic pathway. NK cells mediate cell death in part by FcγRIIIA's recognition of IgG bound to an antigen on a cell surface. The ADCC assays used in accordance with the methods of the invention may be radioactive based assays or fluorescence based assays. The ADCC assay used to characterize the molecules of the invention comprising variant Fc regions comprises labeling target cells, *e.g.*, SK-BR-3, MCF-7, OVCAR3, Raji, Daudi cells, opsonizing target cells with an antibody that recognizes a cell surface receptor on the target cell via its antigen binding site; combining the labeled opsonized target cells and the effector cells at an appropriate ratio, which can be determined by routine experimentation; harvesting the cells; detecting the label in the supernatant of the lysed target cells, using an appropriate detection scheme based on the label used. The target cells may be labeled either with a radioactive label or a fluorescent label, using standard methods known in the art. For example the labels include, but are not limited to, [⁵¹Cr]Na₂CrO₄; and the acetoxymethyl ester of the fluorescence enhancing ligand, 2,2':6',2''-terpyridine-6-6''-dicarboxylate (TDA).

[00215] In a specific preferred embodiment, a time resolved fluorimetric assay is used for measuring ADCC activity against target cells that have been labeled with the acetoxymethyl ester of the fluorescence enhancing ligand, 2,2':6',2''-terpyridine-6-6''-dicarboxylate (TDA). Such fluorimetric assays are known in the art, *e.g.*, *see*, Blomberg

et al., 1996, *Journal of Immunological Methods*, 193: 199-206; which is incorporated herein by reference in its entirety. Briefly, target cells are labeled with the membrane permeable acetoxymethyl diester of TDA (bis(acetoxymethyl) 2,2':6',2''-terpyridine-6-6''-dicarboxylate, (BATDA), which rapidly diffuses across the cell membrane of viable
5 cells. Intracellular esterases split off the ester groups and the regenerated membrane impermeable TDA molecule is trapped inside the cell. After incubation of effector and target cells, *e.g.*, for at least two hours, up to 3.5 hours, at 37°C, under 5% CO₂, the TDA released from the lysed target cells is chelated with Eu³⁺ and the fluorescence of the Europium-TDA chelates formed is quantitated in a time-resolved fluorometer (*e.g.*, Victor
10 1420, Perkin Elmer/Wallac).

[00216] In another specific embodiment, the ADCC assay used to characterize the molecules of the invention comprising variant Fc regions comprises the following steps: Preferably 4-5x10⁶ target cells (*e.g.*, SK-BR-3, MCF-7, OVCAR3, Raji cells) are labeled with bis(acetoxymethyl) 2,2':6',2''-terpyridine-t-6''-dicarboxylate (DELFIATDA
15 Reagent, Perkin Elmer/Wallac). For optimal labeling efficiency, the number of target cells used in the ADCC assay should preferably not exceed 5x10⁶. BATDA reagent is added to the cells and the mixture is incubated at 37°C preferably under 5% CO₂, for at least 30 minutes. The cells are then washed with a physiological buffer, *e.g.*, PBS with 0.125 mM sulfinpyrazole, and media containing 0.125 mM sulfinpyrazole. The labeled target cells are
20 then opsonized (coated) with a molecule of the invention comprising a variant Fc region, *i.e.*, an immunoglobulin comprising a variant Fc region of the invention, including, but not limited to, a polyclonal antibody, a monoclonal antibody, a bispecific antibody, a multi-specific antibody, a humanized antibody, or a chimeric antibody. In preferred embodiments, the immunoglobulin comprising a variant Fc region used in the ADCC assay
25 is specific for a cell surface receptor, a tumor antigen, or a cancer antigen. The immunoglobulin into which a variant Fc region of the invention is introduced may specifically bind any cancer or tumor antigen, such as those listed in section 5.4. Additionally, the immunoglobulin into which a variant Fc region of the invention is introduced may be any therapeutic antibody specific for a cancer antigen, such as those
30 listed in section 5.4. In some embodiments, the immunoglobulin comprising a variant Fc region used in the ADCC assay is an anti-fluorescein monoclonal antibody, 4-4-20 (Kranz *et al.*, 1982 *J. Biol. Chem.* 257(12): 6987-6995) a mouse-human chimeric anti-CD20 monoclonal antibody 2H7 (Liu *et al.*, 1987, *Journal of Immunology*, 139: 3521-6); or a humanized antibody (Ab4D5) against the human epidermal growth factor receptor 2 (p185
35 HER2) (Carter *et al.* (1992, *Proc. Natl. Acad. Sci. USA* 89: 4285-9). The target cells in the

ADCC assay are chosen according to the immunoglobulin into which a variant Fc region of the invention has been introduced so that the immunoglobulin binds a cell surface receptor of the target cell specifically. Preferably, the ADCC assays of the invention are performed using more than one engineered antibody, *e.g.*, anti Her2/neu, 4-4-20, 2B6, Rituxan, and 2H7, harboring the Fc variants of the invention. In a most preferred embodiment, the Fc variants of the invention are introduced into at least 3 antibodies and their ADCC activities is tested. Although not intending to be bound by a particular mechanism of action, examining at least 3 antibodies in these functional assays will diminish the chance of eliminating a viable Fc mutation erroneously.

10 [00217] Opsonized target cells are added to effector cells, *e.g.*, PBMC, to produce effector:target ratios of approximately 50:1, 75:1, or 100:1. In a specific embodiment, when the immunoglobulin comprising a variant Fc region has the variable domain of 4-4-20, the effector:target is 75:1. The effector and target cells are incubated for at least two hours, up to 3.5 hours, at 37°C, under 5% CO₂. Cell supernatants are harvested and added to an

15 acidic europium solution (*e.g.*, DELFIA Europium Solution, Perkin Elmer/Wallac). The fluorescence of the Europium-TDA chelates formed is quantitated in a time-resolved fluorometer (*e.g.*, Victor 1420, Perkin Elmer/Wallac). Maximal release (MR) and spontaneous release (SR) are determined by incubation of target cells with 1% TX-100 and media alone, respectively. Antibody independent cellular cytotoxicity (AICC) is measured

20 by incubation of target and effector cells in the absence of antibody. Each assay is preferably performed in triplicate. The mean percentage specific lysis is calculated as: Experimental release (ADCC) - AICC / (MR - SR) x 100.

5.2.7 OTHER ASSAYS

[00218] The molecules of the invention comprising variant Fc regions may also be

25 assayed using any surface plasmon resonance based assays known in the art for characterizing the kinetic parameters of Fc-FcγR interaction binding. Any SPR instrument commercially available including, but not limited to, BIAcore Instruments, available from BIAcore AB (Uppsala, Sweden); IAsys instruments available from Affinity Sensors (Franklin, MA.); IBIS system available from Windsor Scientific Limited (Berks, UK), SPR-

30 CELLIA systems available from Nippon Laser and Electronics Lab (Hokkaido, Japan), and SPR Detector Spreeta available from Texas Instruments (Dallas, TX) can be used in the instant invention. For a review of SPR-based technology see Mullet *et al.*, 2000, *Methods* 22: 77-91; Dong *et al.*, 2002, *Review in Mol. Biotech.*, 82: 303-23; Fivash *et al.*, 1998, *Current Opinion in Biotechnology* 9: 97-101; Rich *et al.*, 2000, *Current Opinion in*

Biotechnology 11: 54-61; all of which are incorporated herein by reference in their entirety. Additionally, any of the SPR instruments and SPR based methods for measuring protein-protein interactions described in U.S. Patent No.'s 6,373,577; 6,289,286; 5,322,798; 5,341,215; 6,268,125 are contemplated in the methods of the invention, all of which are
5 incorporated herein by reference in their entirety.

[00219] Briefly, SPR based assays involve immobilizing a member of a binding pair on a surface, and monitoring its interaction with the other member of the binding pair in solution in real time. SPR is based on measuring the change in refractive index of the solvent near the surface that occurs upon complex formation or dissociation. The surface
10 onto which the immobilization occur is the sensor chip, which is at the heart of the SPR technology; it consists of a glass surface coated with a thin layer of gold and forms the basis for a range of specialized surfaces designed to optimize the binding of a molecule to the surface. A variety of sensor chips are commercially available especially from the companies listed *supra*, all of which may be used in the methods of the invention.

15 Examples of sensor chips include those available from BIAcore AB, Inc., *e.g.*, Sensor Chip CM5, SA, NTA, and HPA. A molecule of the invention may be immobilized onto the surface of a sensor chip using any of the immobilization methods and chemistries known in the art, including but not limited to, direct covalent coupling via amine groups, direct covalent coupling via sulfhydryl groups, biotin attachment to avidin coated surface,
20 aldehyde coupling to carbohydrate groups, and attachment through the histidine tag with NTA chips.

[00220] In some embodiments, the kinetic parameters of the binding of molecules of the invention comprising variant Fc regions, *e.g.*, immunoglobulins comprising variant Fc region, to an FcγR may be determined using a BIAcore instrument (*e.g.*, BIAcore
25 instrument 1000, BIAcore Inc., Piscataway, NJ). Any FcγR can be used to assess the interaction with the molecules of the invention comprising variant Fc regions. In a specific embodiment the FcγR is FcγRIIIA, preferably a soluble monomeric FcγRIIIA. For example, in one embodiment, the soluble monomeric FcγRIIIA is the extracellular region of FcγRIIIA joined to the linker-AVITAG sequence (*see*, U.S. Provisional Application No.
30 60/439,498, filed on January 9, 2003 (Attorney Docket No. 11183-004-888) and U.S. Provisional Application No. 60/456,041 filed on March 19, 2003, which are incorporated herein by reference in their entireties). In another specific embodiment, the FcγR is FcγRIIB, preferably a soluble dimeric FcγRIIB. For example in one embodiment, the soluble dimeric FcγRIIB protein is prepared in accordance with the methodology described

in U.S. Provisional application No. 60/439,709 filed on January 13, 2003, which is incorporated herein by reference in its entirety.

[00221] An exemplary assay for determining the kinetic parameters of a molecule comprising a variant Fc region, wherein the molecule is the 4-4-20 antibody, to an FcγR using a BIAcore instrument comprises the following: BSA-FITC is immobilized on one of the four flow cells of a sensor chip surface, preferably through amine coupling chemistry such that about 5000 response units (RU) of BSA-FITC is immobilized on the surface. Once a suitable surface is prepared, 4-4-20 antibodies carrying the Fc mutations are passed over the surface, preferably by one minute injections of a 20 μg/mL solution at a 5 μL/mL flow rate. The level of 4-4-20 antibodies bound to the surface ranges between 400 and 700 RU. Next, dilution series of the receptor (FcγRIIA and FcγRIIB-Fc fusion protein) in HBS-P buffer (20mM HEPES, 150 mM NaCl, 3mM EDTA, pH 7.5) are injected onto the surface at 100 μL/min. Antibody regeneration between different receptor dilutions is carried out preferably by single 5 second injections of 100mM NaHCO₃ pH 9.4; 3M NaCl. Any regeneration technique known in the art is contemplated in the method of the invention.

[00222] Once an entire data set is collected, the resulting binding curves are globally fitted using computer algorithms supplied by the SPR instrument manufacturer, e.g., BIAcore, Inc. (Piscataway, NJ). These algorithms calculate both the K_{on} and K_{off} , from which the apparent equilibrium binding constant, K_d is deduced as the ratio of the two rate constants (i.e., K_{off}/K_{on}). More detailed treatments of how the individual rate constants are derived can be found in the BIAevaluation Software Handbook (BIAcore, Inc., Piscataway, NJ). The analysis of the generated data may be done using any method known in the art. For a review of the various methods of interpretation of the kinetic data generated see Myszkka, 1997, *Current Opinion in Biotechnology* 8: 50-7; Fisher *et al.*, 1994, *Current Opinion in Biotechnology* 5: 389-95; O'Shannessy, 1994, *Current Opinion in Biotechnology*, 5:65-71; Chaiken *et al.*, 1992, *Analytical Biochemistry*, 201: 197-210; Morton *et al.*, 1995, *Analytical Biochemistry* 227: 176-85; O'Shannessy *et al.*, 1996, *Analytical Biochemistry* 236: 275-83; all of which are incorporated herein by reference in their entirety.

[00223] In preferred embodiments, the kinetic parameters determined using an SPR analysis, e.g., BIAcore, may be used as a predictive measure of how a molecule of the invention will function in a functional assay, e.g., ADCC. An exemplary method for predicting the efficacy of a molecule of the invention based on kinetic parameters obtained from an SPR analysis may comprise the following: determining the K_{off} values for binding

of a molecule of the invention to FcγRIIIA and FcγRIIB; plotting (1) $K_{\text{off}}(\text{wt})/K_{\text{off}}(\text{mut})$ for FcγRIIIA; (2) $K_{\text{off}}(\text{mut})/K_{\text{off}}(\text{wt})$ for FcγRIIB against the ADCC data. Numbers higher than one show a decreased dissociation rate for FcγRIIIA and an increased dissociation rate for FcγRIIB relative to wild tyoe; and possess and enhanced ADCC function.

5

5.3 METHODS OF RECOMBINANTLY PRODUCING MOLECULES OF THE INVENTION

5.3.1 POLYNUCLEOTIDES ENCODING MOLECULES OF THE INVENTION

- 10 [00224] The present invention also includes polynucleotides that encode the molecules, including the polypeptides and antibodies, of the invention identified by the methods of the invention. The polynucleotides encoding the molecules of the invention may be obtained, and the nucleotide sequence of the polynucleotides determined, by any method known in the art.
- 15 [00225] Once the nucleotide sequence of the molecules (*e.g.*, antibodies) that are identified by the methods of the invention is determined, the nucleotide sequence may be manipulated using methods well known in the art, *e.g.*, recombinant DNA techniques, site directed mutagenesis, PCR, *etc.* (*see*, for example, the techniques described in Sambrook *et al.*, 2001, Molecular Cloning, A Laboratory Manual, 3rd Ed., Cold Spring Harbor
- 20 Laboratory, Cold Spring Harbor, NY; and Ausubel *et al.*, eds., 1998, Current Protocols in Molecular Biology, John Wiley & Sons, NY, which are both incorporated by reference herein in their entirety), to generate, for example, antibodies having a different amino acid sequence, for example by generating amino acid substitutions, deletions, and/or insertions.
- [00226] In a specific embodiment, when the nucleic acids encode antibodies, one or
- 25 more of the CDRs are inserted within framework regions using routine recombinant DNA techniques. The framework regions may be naturally occurring or consensus framework regions, and preferably human framework regions (*see, e.g.*, Chothia *et al.*, 1998, *J. Mol. Biol.* 278: 457-479 for a listing of human framework regions).
- [00227] In another embodiment, human libraries or any other libraries available in
- 30 the art, can be screened by standard techniques known in the art, to clone the nucleic acids encoding the molecules of the invention.

5.3.2 RECOMBINANT EXPRESSION OF MOLECULES OF THE INVENTION

- [00228] Once a nucleic acid sequence encoding molecules of the invention (*i.e.*, antibodies) has been obtained, the vector for the production of the molecules may be produced by recombinant DNA technology using techniques well known in the art. Methods which are well known to those skilled in the art can be used to construct expression vectors containing the coding sequences for the molecules of the invention and appropriate transcriptional and translational control signals. These methods include, for example, *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. (*See*, for example, the techniques described in Sambrook *et al.*, 1990, Molecular Cloning, A Laboratory Manual, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY and Ausubel *et al.* eds., 1998, Current Protocols in Molecular Biology, John Wiley & Sons, NY).
- [00229] An expression vector comprising the nucleotide sequence of a molecule identified by the methods of the invention (*i.e.*, an antibody) can be transferred to a host cell by conventional techniques (*e.g.*, electroporation, liposomal transfection, and calcium phosphate precipitation) and the transfected cells are then cultured by conventional techniques to produce the molecules of the invention. In specific embodiments, the expression of the molecules of the invention is regulated by a constitutive, an inducible or a tissue, specific promoter.
- [00230] The host cells used to express the molecules identified by the methods of the invention may be either bacterial cells such as *Escherichia coli*, or, preferably, eukaryotic cells, especially for the expression of whole recombinant immunoglobulin molecule. In particular, mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus is an effective expression system for immunoglobulins (Foecking *et al.*, 1998, *Gene* 45:101; Cockett *et al.*, 1990, *Bio/Technology* 8:2).
- [00231] A variety of host-expression vector systems may be utilized to express the molecules identified by the methods of the invention. Such host-expression systems represent vehicles by which the coding sequences of the molecules of the invention may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express the molecules of the invention *in situ*. These include, but are not limited to, microorganisms such as bacteria (*e.g.*, *E. coli* and *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing coding sequences for the molecules

- identified by the methods of the invention; yeast (*e.g.*, *Saccharomyces Pichia*) transformed with recombinant yeast expression vectors containing sequences encoding the molecules identified by the methods of the invention; insect cell systems infected with recombinant virus expression vectors (*e.g.*, baculovirus) containing the sequences encoding the
- 5 molecules identified by the methods of the invention; plant cell systems infected with recombinant virus expression vectors (*e.g.*, cauliflower mosaic virus (CaMV) and tobacco mosaic virus (TMV) or transformed with recombinant plasmid expression vectors (*e.g.*, Ti plasmid) containing sequences encoding the molecules identified by the methods of the invention; or mammalian cell systems (*e.g.*, COS, CHO, BHK, 293, 293T, 3T3 cells,
- 10 lymphotic cells (*see* U.S. 5,807,715), Per C.6 cells (human retinal cells developed by Crucell) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (*e.g.*, metallothionein promoter) or from mammalian viruses (*e.g.*, the adenovirus late promoter; the vaccinia virus 7.5K promoter).
- [00232] In bacterial systems, a number of expression vectors may be advantageously
- 15 selected depending upon the use intended for the molecule being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of an antibody, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther *et al.*, 1983,
- 20 *EMBO J.* 2:1791), in which the antibody coding sequence may be ligated individually into the vector in frame with the *lac Z* coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, 1985, *Nucleic Acids Res.* 13:3101-3109; Van Heeke & Schuster, 1989, *J. Biol. Chem.* 24:5503-5509); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In
- 25 general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption and binding to a matrix glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.
- 30 [00233] In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The antibody coding sequence may be cloned individually into non-essential regions (*e.g.*, the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (*e.g.*, the polyhedrin promoter).

[00234] In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the antibody coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, *e.g.*, the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (*e.g.*, region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the immunoglobulin molecule in infected hosts (*e.g.*, see Logan & Shenk, 1984, *Proc. Natl. Acad. Sci. USA* 81:355-359). Specific initiation signals may also be required for efficient translation of inserted antibody coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, *etc.* (see Bittner *et al.*, 1987, *Methods in Enzymol.* 153:51-544).

[00235] In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (*e.g.*, glycosylation) and processing (*e.g.*, cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERY, BHK, Hela, COS, MDCK, 293, 293T, 3T3, WI38, BT483, Hs578T, HTB2, BT20 and T47D, CRL7030 and Hs578Bst.

[00236] For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express an antibody of the invention may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (*e.g.*, promoter, enhancer, sequences, transcription terminators, polyadenylation sites, *etc.*), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media,

and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which
5 express the antibodies of the invention. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that interact directly or indirectly with the antibodies of the invention.

[00237] A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler *et al.*, 1977, *Cell* 11: 223), hypoxanthine-
10 guanine phosphoribosyltransferase (Szybalska & Szybalski, 1992, *Proc. Natl. Acad. Sci. USA* 48: 202), and adenine phosphoribosyltransferase (Lowy *et al.*, 1980, *Cell* 22: 817) genes can be employed in tk-, hgp^{rt}- or ap^{rt}- cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler *et al.*, 1980, *Proc. Natl. Acad. Sci. USA* 77:357; O'Hare
15 *et al.*, 1981, *Proc. Natl. Acad. Sci. USA* 78: 1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, *Proc. Natl. Acad. Sci. USA* 78: 2072); neo, which confers resistance to the aminoglycoside G-418 *Clinical Pharmacy* 12: 488-505; Wu and Wu, 1991, 3:87-95; Tolstoshev, 1993, *Ann. Rev. Pharmacol. Toxicol.* 32:573-596; Mulligan, 1993, *Science* 260:926-932; and Morgan and Anderson, 1993, *Ann. Rev.*
20 *Biochem.* 62:191-217; May, 1993, *TIB TECH* 11(5):155-215). Methods commonly known in the art of recombinant DNA technology which can be used are described in Ausubel *et al.* (eds.), 1993, *Current Protocols in Molecular Biology*, John Wiley & Sons, NY; Kriegler, 1990, *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY; and in Chapters 12 and 13, Dracopoli *et al.* (eds), 1994, *Current Protocols in Human Genetics*,
25 John Wiley & Sons, NY.; Colberre-Garapin *et al.*, 1981, *J. Mol. Biol.* 150:1; and hyg^{ro}, which confers resistance to hygromycin (Santerre *et al.*, 1984, *Gene* 30:147).

[00238] The expression levels of an antibody of the invention can be increased by vector amplification (for a review, see Bebbington and Hentschel, The use of vectors based on gene amplification for the expression of cloned genes in mammalian cells in DNA
30 cloning, Vol. 3 (Academic Press, New York, 1987). When a marker in the vector system expressing an antibody is amplifiable, increase in the level of inhibitor present in culture of host cell will increase the number of copies of the marker gene. Since the amplified region is associated with the nucleotide sequence of the antibody, production of the antibody will also increase (Crouse *et al.*, 1983, *Mol. Cell. Biol.* 3:257).

- [00239] The host cell may be co-transfected with two expression vectors of the invention, the first vector encoding a heavy chain derived polypeptide and the second vector encoding a light chain derived polypeptide. The two vectors may contain identical selectable markers which enable equal expression of heavy and light chain polypeptides.
- 5 Alternatively, a single vector may be used which encodes both heavy and light chain polypeptides. In such situations, the light chain should be placed before the heavy chain to avoid an excess of toxic free heavy chain (Proudfoot, 1986, *Nature* 322:52; Kohler, 1980, *Proc. Natl. Acad. Sci. USA* 77:2197). The coding sequences for the heavy and light chains may comprise cDNA or genomic DNA.
- 10 [00240] Once a molecule of the invention (*i.e.*, antibodies) has been recombinantly expressed, it may be purified by any method known in the art for purification of polypeptides or antibodies, for example, by chromatography (*e.g.*, ion exchange, affinity, particularly by affinity for the specific antigen after Protein A, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique
- 15 for the purification of polypeptides or antibodies.

5.4 PROPHYLACTIC AND THERAPEUTIC METHODS

- [00241] The present invention encompasses administering one or more of the molecules of the invention (*e.g.*, antibodies) to an animal, preferably a mammal, and most preferably a human, for preventing, treating, or ameliorating one or more symptoms
- 20 associated with a disease, disorder, or infection. The molecules of the invention are particularly useful for the treatment or prevention of a disease or disorder where an enhanced efficacy of effector cell function (*e.g.*, ADCC) mediated by Fc γ R is desired. The methods and compositions of the invention are particularly useful for the treatment or prevention of primary or metastatic neoplastic disease (*i.e.*, cancer), and infectious diseases.
- 25 Molecules of the invention may be provided in pharmaceutically acceptable compositions as known in the art or as described herein. As detailed below, the molecules of the invention can be used in methods of treating or preventing cancer (particularly in passive immunotherapy), autoimmune disease, inflammatory disorders or infectious diseases.
- [00242] The molecules of the invention may also be advantageously utilized in
- 30 combination with other therapeutic agents known in the art for the treatment or prevention of a cancer, autoimmune disease, inflammatory disorders or infectious diseases. In a specific embodiment, molecules of the invention may be used in combination with monoclonal or chimeric antibodies, lymphokines, or hematopoietic growth factors (such as, *e.g.*, IL-2, IL-3 and IL-7), which, for example, serve to increase the number or activity of

effector cells which interact with the molecules and, increase immune response. The molecules of the invention may also be advantageously utilized in combination with one or more drugs used to treat a disease, disorder, or infection such as, for example anti-cancer agents, anti-inflammatory agents or anti-viral agents, *e.g.*, as detailed in sections 5.4.1.2 and 5.4.2.1 below.

5.4.1 CANCERS

- [00243] The invention encompasses methods and composition for treatment or prevention of cancer or metastasis in a subject comprising administering to the subject a therapeutically effective amount of one or more molecules comprising a variant Fc region.
- 10 [00244] Molecules of the invention (*i.e.*, polypeptides, antibodies) comprising variant Fc regions can be used to prevent, inhibit or reduce the growth of primary tumors or metastasis of cancerous cells. In one embodiment, the molecule of the invention comprises a variant Fc that binds Fc γ RIIA and/or Fc γ RIIA with a greater affinity than a comparable polypeptide comprising a wild type Fc region binds Fc γ RIIA and/or Fc γ RIIA, and/or said
- 15 variant Fc region has an enhanced effector function, *e.g.*, ADCC, CDC, phagocytosis, opsonization, *etc.* Such molecules can be used alone to treat or prevent cancer. In another embodiment, the molecule of the invention comprises a variant Fc region that binds Fc γ RIIA and/or Fc γ RIIA with a greater affinity than a comparable polypeptide comprising a wild type Fc region binds Fc γ RIIA and/or Fc γ RIIA, and further binds Fc γ RIIB with a
- 20 lower affinity than a comparable polypeptide comprising a wild-type Fc region binds Fc γ RIIB, and/or said variant Fc region has an enhanced effector function, *e.g.*, ADCC, CDC, phagocytosis, opsonization, *etc.* Such molecules can also be used alone to treat or prevent cancer.
- [00245] In some embodiments, the invention encompasses methods and compositions
- 25 for the treatment or prevention of cancer in a subject with Fc γ R polymorphisms such as those homozygous for the Fc γ RIIA-158V or Fc γ RIIA-158F alleles. In some embodiments, the invention encompasses engineering therapeutic antibodies, *e.g.*, tumor specific monoclonal antibodies in accordance with the methods of the invention such that the engineered antibodies have enhanced efficacy in patients homozygous for the low affinity
- 30 allele of Fc γ RIIA (158F). In other embodiments, the invention encompasses engineering therapeutic antibodies, *e.g.*, tumor specific monoclonal antibodies in accordance with the methods of the invention such that the engineered antibodies have enhanced efficacy in patients homozygous for the high affinity allele of Fc γ RIIA (158V).

[00246] In some embodiments, the engineered antibodies of the invention are particularly effective in treating and/or preventing non-Hodgkin's lymphoma (NHL). The engineered antibodies of the invention are therapeutically more effective than current therapeutic regimens for NHL, including but not limited to chemotherapy, and
5 immunotherapy using anti-CD20 mAb, Rituximab. The efficacy of anti-CD20 monoclonal antibodies however depends on the FcγR polymorphism of the subject (Carton *et al.*, 2002 *Blood*, 99: 754-8; Weng *et al.*, 2003 *J Clin Oncol*.21(21):3940-7 both of which are incorporated herein by reference in their entireties). These receptors are expressed on the surface of the effector cells and mediate ADCC. High affinity alleles, of the low affinity
10 activating receptors, improve the effector cells' ability to mediate ADCC. The methods of the invention allow engineering anti-CD20 antibodies harboring Fc mutations to enhance their affinity to FcγR on effector cells via their altered Fc domains. The engineered antibodies of the invention provide better immunotherapy reagents for patients regardless of their FcγR polymorphism.

15 [00247] An exemplary method for determining the efficacy of the engineered anti-CD20 antibodies in a subject may include the following: Plasmids harboring chimeric anti-HER2/neu heavy chain genes with Fc mutations that show substantially increased killing in ADCC can be used as a backbone to transfer in the variable domain from the Rituximab heavy chain gene. The variable region from the anti-HER2/neu Fc variant is replaced with
20 the variable region from Rituximab. Plasmids containing wild type Fc domains or a D265A mutation to abrogate FcR binding, or the anti-CD20 Fc variants are transiently cotransfected with the Rituximab light chain gene into 293H cells, conditioned media and the antibody is purified over a protein G column using routine methods.

[00248] Anti-CD20 mAbs harboring the Fc variants are tested by ADCC using a
25 cultured B cell line to determine the ability of the Fc mutations to enhance ADCC. Standard ADCC is performed using methods disclosed herein. Lymphocytes are harvested from peripheral blood using a Ficoll-Paque gradient (Pharmacia). Target Daudi cells, a B-cell line expressing CD20, are loaded with Europium (PerkinElmer) and incubated with effectors for 4 hrs at 37°C. Released Europium is detected using a fluorescent plate reader
30 (Wallac). The resulting ADCC data indicates the efficacy of the Fc variants to trigger NK cell mediated cytotoxicity and establish which anti-CD20 Fc variants can be tested with both patient samples and elutriated monocytes. Fc variants showing the greatest potential for enhancing the efficacy of the anti-CD20 antibody are then tested in an ADCC assay using PBMCs from patients. PBMC from healthy donors are used as effector cells. *In vitro*
35 ADCC assays using anti-CD20 variants and Rituximab are performed in primary

lymphoma cells from patients with follicular lymphoma. The specific FcγR polymorphism of the donors is determined and cataloged using methods known in the art. ADCC assay is performed by effector cells from patients with different FcγRIIIA and FcγRIIA genotypes.

[00249] According to an aspect of the invention, molecules (*e.g.*, antibodies) of the invention comprising variant Fc regions enhance the efficacy of cancer immunotherapy by increasing the potency of the antibody effector function relative to a molecule containing the wild-type Fc region, *e.g.*, ADCC, CDC, phagocytosis, opsonization, *etc.* In a specific embodiment, antibody dependent cellular toxicity and/or phagocytosis of tumor cells is enhanced using the molecules of the invention with variant Fc regions. Molecules of the invention may enhance the efficacy of immunotherapy cancer treatment by enhancing at least one antibody-mediated effector function. In one particular embodiment, a molecule of the invention comprising a variant Fc region enhances the efficacy of immunotherapy treatment by enhancing the complement dependent cascade. In another embodiment of the invention, the molecule of the invention comprising a variant Fc region enhances the efficacy of immunotherapy treatment by enhancing the phagocytosis and/or opsonization of the targeted tumor cells. In another embodiment of the invention, the molecule of the invention comprising a variant Fc region enhances the efficacy of treatment by enhancing antibody-dependent cell-mediated cytotoxicity ("ADCC") in destruction of the targeted tumor cells.

[00250] The invention further contemplates engineering therapeutic antibodies (*e.g.*, tumor specific monoclonal antibodies) for enhancing the therapeutic efficacy of the therapeutic antibody, for example, by enhancing the effector function of the therapeutic antibody (*e.g.*, ADCC). Preferably the therapeutic antibody is a cytotoxic and/or opsonizing antibody. It will be appreciated by one of skill in the art, that once molecules of the invention with desired binding properties (*e.g.*, molecules with variant Fc regions with at least one amino acid modification, which modification enhances the affinity of the variant Fc region for FcγRIIIA and/or FcγRIIA relative to a comparable molecule, comprising a wild-type Fc region) have been identified (*See* Section 5.2 and Table 5) according to the methods of the invention, therapeutic antibodies may be engineered using standard recombinant DNA techniques and any known mutagenesis techniques, as described in Section 5.2.2 to produce engineered therapeutic carrying the identified mutation sites with the desired binding properties. Any of the therapeutic antibodies listed in Table 6 that have demonstrated therapeutic utility in cancer treatment, may be engineered according to the methods of the invention, for example, by modifying the Fc region to have an enhanced affinity for FcγRIIIA and/or FcγRIIA compared to a therapeutic antibody having a wild-

type Fc region, and used for the treatment and or prevention of a cancer characterized by a cancer antigen.

- [00251] The invention also encompasses engineering any other polypeptide comprising an Fc region which has therapeutic utility, including but not limited to
- 5 ENBREL, according to the methods of the invention, in order to enhance the therapeutic efficacy of such polypeptides, for example, by enhancing the effector function of the polypeptide comprising an Fc region.

TABLE 6. THERAPEUTIC ANTIBODIES THAT CAN BE ENGINEERED ACCORDING TO THE METHODS OF THE INVENTION

Company	Product	Disease	Target
Abgenix	ABX-EGF	Cancer	EGF receptor
AltaRex	OvaRex	ovarian cancer	tumor antigen CA125
	BravaRex	metastatic cancers	tumor antigen MUC1
Antisoma	Theragyn (pentumomabytrrium-90)	ovarian cancer	PEM antigen
	Therex	breast cancer	PEM antigen
Boehringer Ingelheim	Blvatuzumab	head & neck cancer	CD44
Centocor/J&J	Panorex	Colorectal cancer	17-1A
	ReoPro	PTCA	gp IIIb/IIIa
	ReoPro	Acute MI	gp IIIb/IIIa
	ReoPro	Ischemic stroke	gp IIIb/IIIa
Corixa	Bexocar	NHL	CD20
CRC Technology	MAb, idiotypic 105AD7	colorectal cancer vaccine	gp72
Crucell	Anti-EpCAM	cancer	Ep-CAM
Cytoclonal	MAB, lung cancer	non-small cell lung cancer	NA
Genentech	Herceptin	metastatic breast cancer	HER-2
	Herceptin	early stage breast cancer	HER-2
	Rituxan	Relapsed/refractory low-grade or follicular NHL	CD20
	Rituxan	intermediate &	CD20

Company	Product	Disease	Target
		high-grade NHL	
	MAB-VEGF	NSCLC, metastatic	VEGF
	MAB-VEGF	Colorectal cancer, metastatic	VEGF
	AMD Fab	age-related macular degeneration	CD18
	E-26 (2 nd gen. IgE)	allergic asthma & rhinitis	IgE
IDEC	Zevalin (Rituxan + yttrium-90)	low grade of follicular, relapsed or refractory, CD20-positive, B-cell NHL and Rituximab- refractory NHL	CD20
ImClone	Cetuximab + innotecan	refractory colorectal carcinoma	EGF receptor
	Cetuximab + cisplatin & radiation	newly diagnosed or recurrent head & neck cancer	EGF receptor
	Cetuximab + gemcitabine	newly diagnosed metastatic pancreatic carcinoma	EGF receptor
	Cetuximab + cisplatin + 5FU or Taxol	recurrent or metastatic head & neck cancer	EGF receptor
	Cetuximab + carboplatin + paclitaxel	newly diagnosed non-small cell lung carcinoma	EGF receptor
	Cetuximab + cisplatin	head & neck cancer (extensive incurable local- regional disease & distant metastases)	EGF receptor
	Cetuximab + radiation	locally advanced head & neck carcinoma	EGF receptor

Company	Product	Disease	Target
	BEC2 + Bacillus Calmette Guerin	small cell lung carcinoma	mimics ganglioside GD3
	BEC2 + Bacillus Calmette Guerin	melanoma	mimics ganglioside GD3
	IMC-1C11	colorectal cancer with liver metastases	VEGF-receptor
ImmonoGen	nuC242-DM1	Colorectal, gastric, and pancreatic cancer	nuC242
ImmunoMedics	LymphoCide	Non-Hodgkins lymphoma	CD22
	LymphoCide Y-90	Non-Hodgkins lymphoma	CD22
	CEA-Cide	metastatic solid tumors	CEA
	CEA-Cide Y-90	metastatic solid tumors	CEA
	CEA-Scan (Tc-99m-labeled arcitumomab)	colorectal cancer (radioimaging)	CEA
	CEA-Scan (Tc-99m-labeled arcitumomab)	Breast cancer (radioimaging)	CEA
	CEA-Scan (Tc-99m-labeled arcitumomab)	lung cancer (radioimaging)	CEA
	CEA-Scan (Tc-99m-labeled arcitumomab)	intraoperative tumors (radio imaging)	CEA
	LeukoScan (Tc-99m-labeled sulesomab)	soft tissue infection (radioimaging)	CEA
	LymphoScan (Tc-99m-labeled)	lymphomas (radioimaging)	CD22
	AFP-Scan (Tc-99m-labeled)	liver 7 gem-cell cancers (radioimaging)	AFP
Intracel	HumaRAD-HN (+ yttrium-90)	head & neck cancer	NA
	HumaSPECT	colorectal imaging	NA
Medarex	MDX-101 (CTLA-4)	Prostate and other cancers	CTLA-4
	MDX-210 (her-2)	Prostate cancer	HER-2

Company	Product	Disease	Target
MedImmune Merck KGaA	overexpression)		
	MDX-210/MAK	Cancer	HER-2
	Vitaxin	Cancer	$\alpha v \beta_3$
	MAB 425	Various cancers	EGF receptor
Millennium	IS-IL-2	Various cancers	Ep-CAM
	Campath (alemtuzumab)	chronic lymphocytic leukemia	CD52
NeoRx	CD20-streptavidin (+ biotin-yttrium 90)	Non-Hodgkins lymphoma	CD20
	Avidicin (albumin + NRLU13)	metastatic cancer	NA
Peregrine	Oncolym (+ iodine-131)	Non-Hodgkins lymphoma	HLA-DR 10 beta
	Cotara (+ iodine-131)	unresectable malignant glioma	DNA-associated proteins
Pharmacia Corporation	C215 (+ staphylococcal enterotoxin)	pancreatic cancer	NA
	MAB, lung/kidney cancer	lung & kidney cancer	NA
	nacolomab tafenatox (C242 + staphylococcal enterotoxin)	colon & pancreatic cancer	NA
Protein Design Labs	Nuvion	T cell malignancies	CD3
	SMART M195	AML	CD33
Titan	SMART 1D10	NHL	HLA-DR antigen
	CEAVac	colorectal cancer, advanced	CEA
	TriGem	metastatic melanoma & small cell lung cancer	GD2-ganglioside
	TriAb	metastatic breast cancer	MUC-1
Trilex	CEAVac	colorectal cancer, advanced	CEA
	TriGem	metastatic melanoma &	GD2-ganglioside

Company	Product	Disease	Target
	TriAb	small cell lung cancer metastatic breast cancer	MUC-1
Viventia Biotech	NovoMAb-G2 radiolabeled	Non-Hodgkins lymphoma	NA
	Monopharm C	colorectal & pancreatic carcinoma	SK-1 antigen
	GlioMAb-H (+ gelonin toxin)	glioma, melanoma & neuroblastoma	NA
Xoma	Rituxan	Relapsed/refractory low-grade or follicular NHL	CD20
	Rituxan	intermediate & high-grade NHL	CD20
	ING-1	adenomcarcinoma	Ep-CAM

[00252] Accordingly, the invention provides methods of preventing or treating cancer characterized by a cancer antigen, using a therapeutic antibody that binds a cancer antigen and is cytotoxic and has been modified at one or more sites in the Fc region, according to the invention, to bind Fc γ RIIA and/or Fc γ RIIA with a higher affinity than the parent therapeutic antibody, and/or mediates effector function (*e.g.*, ADCC, phagocytosis) more effectively. In another embodiment, the invention provides methods of preventing or treating cancer characterized by a cancer antigen, using a therapeutic antibody that binds a cancer antigen and is cytotoxic, and has been engineered according to the invention to bind Fc γ RIIA and/or Fc γ RIIA with a higher affinity and bind Fc γ RIIB with a lower affinity than the parent therapeutic antibody, and/or mediates effector function (*e.g.*, ADCC, phagocytosis) more effectively. The therapeutic antibodies that have been engineered according to the invention are useful for prevention or treatment of cancer, since they have an enhanced cytotoxic activity (*e.g.*, enhanced tumor cell killing and/or enhanced for example, ADCC activity or CDC activity).

[00253] Cancers associated with a cancer antigen may be treated or prevented by administration of a therapeutic antibody that binds a cancer antigen and is cytotoxic, and has been engineered according to the methods of the invention to have, for example, an enhanced effector function. In one particular embodiment, the therapeutic antibodies

engineered according to the methods of the invention enhance the antibody-mediated cytotoxic effect of the antibody directed at the particular cancer antigen. For example, but not by way of limitation, cancers associated with the following cancer antigens may be treated or prevented by the methods and compositions of the invention: KS 1/4 pan-

5 carcinoma antigen (Perez and Walker, 1990, *J. Immunol.* 142:32-37; Bumal, 1988, *Hybridoma* 7(4):407-415), ovarian carcinoma antigen (CA125) (Yu *et al.*, 1991, *Cancer Res.* 51(2):48-475), prostatic acid phosphate (Tailor *et al.*, 1990, *Nucl. Acids Res.* 18(1):4928), prostate specific antigen (Henttu and Vihko, 1989, *Biochem. Biophys. Res. Comm.* 10(2):903-910; Israeli *et al.*, 1993, *Cancer Res.* 53:227-230), melanoma-associated

10 antigen p97 (Estin *et al.*, 1989, *J. Natl. Cancer Instit.* 81(6):445-44), melanoma antigen gp75 (Vijayasardahl *et al.*, 1990, *J. Exp. Med.* 171(4):1375-1380), high molecular weight melanoma antigen (HMW-MAA) (Natali *et al.*, 1987, *Cancer* 59:55-3; Mittelman *et al.*, 1990, *J. Clin. Invest.* 86:2136-2144), prostate specific membrane antigen, carcinoembryonic antigen (CEA) (Foon *et al.*, 1994, *Proc. Am. Soc. Clin. Oncol.* 13:294),

15 polymorphic epithelial mucin antigen, human milk fat globule antigen, Colorectal tumor-associated antigens such as: CEA, TAG-72 (Yokata *et al.*, 1992, *Cancer Res.* 52:3402-3408), CO17-1A (Ragnhammar *et al.*, 1993, *Int. J. Cancer* 53:751-758); GICA 19-9 (Herlyn *et al.*, 1982, *J. Clin. Immunol.* 2:135), CTA-1 and LEA, Burkitt's lymphoma antigen-38.13, CD19 (Ghetie *et al.*, 1994, *Blood* 83:1329-1336), human B-lymphoma

20 antigen-CD20 (Reff *et al.*, 1994, *Blood* 83:435-445), CD33 (Sgouros *et al.*, 1993, *J. Nucl. Med.* 34:422-430), melanoma specific antigens such as ganglioside GD2 (Saleh *et al.*, 1993, *J. Immunol.*, 151, 3390-3398), ganglioside GD3 (Shitara *et al.*, 1993, *Cancer Immunol. Immunother.* 36:373-380), ganglioside GM2 (Livingston *et al.*, 1994, *J. Clin. Oncol.* 12:1036-1044), ganglioside GM3 (Hoon *et al.*, 1993, *Cancer Res.* 53:5244-5250), tumor-

25 specific transplantation type of cell-surface antigen (TSTA) such as virally-induced tumor antigens including T-antigen DNA tumor viruses and envelope antigens of RNA tumor viruses, oncofetal antigen-alpha-fetoprotein such as CEA of colon, bladder tumor oncofetal antigen (Hellstrom *et al.*, 1985, *Cancer Res.* 45:2210-2188), differentiation antigen such as human lung carcinoma antigen L6, L20 (Hellstrom *et al.*, 1986, *Cancer Res.* 46:3917-3923),

30 antigens of fibrosarcoma, human leukemia T cell antigen-Gp37 (Bhattacharya-Chatterjee *et al.*, 1988, *J. of Immun.* 141:1398-1403), neoglycoprotein, sphingolipids, breast cancer antigen such as EGFR (Epidermal growth factor receptor), HER2 antigen (p185^{HER2}), polymorphic epithelial mucin (PEM) (Hilkens *et al.*, 1992, *Trends in Bio. Chem. Sci.* 17:359), malignant human lymphocyte antigen-APO-1 (Bernhard *et al.*, 1989, *Science*

35 245:301-304), differentiation antigen (Feizi, 1985, *Nature* 314:53-57) such as I antigen

found in fetal erythrocytes and primary endoderm, I(Ma) found in gastric adenocarcinomas, M18 and M39 found in breast epithelium, SSEA-1 found in myeloid cells, VEP8, VEP9, Myl, VIM-D5, and D156-22 found in colorectal cancer, TRA-1-85 (blood group H), C14 found in colonic adenocarcinoma, F3 found in lung adenocarcinoma, AH6 found in gastric cancer, Y hapten, Le^y found in embryonal carcinoma cells, TL5 (blood group A), EGF receptor found in A431 cells, E₁ series (blood group B) found in pancreatic cancer, FC10.2 found in embryonal carcinoma cells, gastric adenocarcinoma, CO-514 (blood group Le^a) found in adenocarcinoma, NS-10 found in adenocarcinomas, CO-43 (blood group Le^b), G49, EGF receptor, (blood group ALe^b/Le^y) found in colonic adenocarcinoma, 19.9 found in colon cancer, gastric cancer mucins, T₅A₇ found in myeloid cells, R₂₄ found in melanoma, 4.2, G_{D3}, D1:1, OFA-1, G_{M2}, OFA-2, G_{D2}, M1:22:25:8 found in embryonal carcinoma cells and SSEA-3, SSEA-4 found in 4-8-cell stage embryos. In another embodiment, the antigen is a T cell receptor derived peptide from a cutaneous T cell lymphoma (see Edelson, 1998, *The Cancer Journal* 4:62).

15 [00254] Cancers and related disorders that can be treated or prevented by methods and compositions of the present invention include, but are not limited to, the following:

Leukemias including, but not limited to, acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemias such as myeloblastic, promyelocytic, myelomonocytic, monocytic, erythroleukemia leukemias and myelodysplastic syndrome, chronic leukemias such as but not limited to, chronic myelocytic (granulocytic) leukemia, chronic lymphocytic leukemia, hairy cell leukemia; polycythemia vera; lymphomas such as but not limited to Hodgkin's disease, non-Hodgkin's disease; multiple myelomas such as but not limited to smoldering multiple myeloma, nonsecretory myeloma, osteosclerotic myeloma, plasma cell leukemia, solitary plasmacytoma and extramedullary plasmacytoma; Waldenström's

25 macroglobulinemia; monoclonal gammopathy of undetermined significance; benign monoclonal gammopathy; heavy chain disease; bone and connective tissue sarcomas such as but not limited to bone sarcoma, osteosarcoma, chondrosarcoma, Ewing's sarcoma, malignant giant cell tumor, fibrosarcoma of bone, chordoma, periosteal sarcoma, soft-tissue sarcomas, angiosarcoma (hemangiosarcoma), fibrosarcoma, Kaposi's sarcoma,

30 leiomyosarcoma, liposarcoma, lymphangiosarcoma, neurilemmoma, rhabdomyosarcoma, synovial sarcoma; brain tumors including but not limited to, glioma, astrocytoma, brain stem glioma, ependymoma, oligodendroglioma, nonglial tumor, acoustic neurinoma, craniopharyngioma, medulloblastoma, meningioma, pineocytoma, pineoblastoma, primary brain lymphoma; breast cancer including, but not limited to, adenocarcinoma, lobular (small

35 cell) carcinoma, intraductal carcinoma, medullary breast cancer, mucinous breast cancer,

tubular breast cancer, papillary breast cancer, Paget's disease, and inflammatory breast cancer; adrenal cancer, including but not limited to, pheochromocytom and adrenocortical carcinoma; thyroid cancer such as but not limited to papillary or follicular thyroid cancer, medullary thyroid cancer and anaplastic thyroid cancer; pancreatic cancer, including but not limited to, insulinoma, gastrinoma, glucagonoma, vipoma, somatostatin-secreting tumor, and carcinoid or islet cell tumor; pituitary cancers including but not limited to, Cushing's disease, prolactin-secreting tumor, acromegaly, and diabetes insipidus; eye cancers including but not limited to, ocular melanoma such as iris melanoma, choroidal melanoma, and ciliary body melanoma, and retinoblastoma; vaginal cancers, including but not limited to, squamous cell carcinoma, adenocarcinoma, and melanoma; vulvar cancer, including but not limited to, squamous cell carcinoma, melanoma, adenocarcinoma, basal cell carcinoma, sarcoma, and Paget's disease; cervical cancers including but not limited to, squamous cell carcinoma, and adenocarcinoma; uterine cancers including but not limited to, endometrial carcinoma and uterine sarcoma; ovarian cancers including but not limited to, ovarian epithelial carcinoma, borderline tumor, germ cell tumor, and stromal tumor; esophageal cancers including but not limited to, squamous cancer, adenocarcinoma, adenoid cystic carcinoma, mucoepidermoid carcinoma, adenosquamous carcinoma, sarcoma, melanoma, plasmacytoma, verrucous carcinoma, and oat cell (small cell) carcinoma; stomach cancers including but not limited to, adenocarcinoma, fungating (polypoid), ulcerating, superficial spreading, diffusely spreading, malignant lymphoma, liposarcoma, fibrosarcoma, and carcinosarcoma; colon cancers; rectal cancers; liver cancers including but not limited to hepatocellular carcinoma and hepatoblastoma, gallbladder cancers including but not limited to, adenocarcinoma; cholangiocarcinomas including but not limited to, papillary, nodular, and diffuse; lung cancers including but not limited to, non-small cell lung cancer, squamous cell carcinoma (epidermoid carcinoma), adenocarcinoma, large-cell carcinoma and small-cell lung cancer; testicular cancers including but not limited to, germinal tumor, seminoma, anaplastic, classic (typical), spermatocytic, nonseminoma, embryonal carcinoma, teratoma carcinoma, choriocarcinoma (yolk-sac tumor), prostate cancers including but not limited to, adenocarcinoma, leiomyosarcoma, and rhabdomyosarcoma; penile cancers; oral cancers including but not limited to, squamous cell carcinoma; basal cancers; salivary gland cancers including but not limited to, adenocarcinoma, mucoepidermoid carcinoma, and adenoidcystic carcinoma; pharynx cancers including but not limited to, squamous cell cancer, and verrucous; skin cancers including but not limited to, basal cell carcinoma, squamous cell carcinoma and melanoma, superficial spreading melanoma, nodular melanoma, lentigo malignant melanoma, acral lentiginous melanoma; kidney cancers

including but not limited to, renal cell cancer, adenocarcinoma, hypernephroma, fibrosarcoma, transitional cell cancer (renal pelvis and/ or uterer); Wilms' tumor; bladder cancers including but not limited to, transitional cell carcinoma, squamous cell cancer, adenocarcinoma, carcinosarcoma. In addition, cancers include myxosarcoma, osteogenic sarcoma, endotheliosarcoma, lymphangioendotheliosarcoma, mesothelioma, synovioma, hemangioblastoma, epithelial carcinoma, cystadenocarcinoma, bronchogenic carcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma and papillary adenocarcinomas (for a review of such disorders, see Fishman *et al.*, 1985, *Medicine*, 2d Ed., J.B. Lippincott Co., Philadelphia and Murphy *et al.*, 1997, *Informed Decisions: The Complete Book of Cancer Diagnosis, Treatment, and Recovery*, Viking Penguin, Penguin Books U.S.A., Inc., United States of America).

[00255] Accordingly, the methods and compositions of the invention are also useful in the treatment or prevention of a variety of cancers or other abnormal proliferative diseases, including (but not limited to) the following: carcinoma, including that of the bladder, breast, colon, kidney, liver, lung, ovary, pancreas, stomach, prostate, cervix, thyroid and skin; including squamous cell carcinoma; hematopoietic tumors of lymphoid lineage, including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, Burkett's lymphoma; hematopoietic tumors of myeloid lineage, including acute and chronic myelogenous leukemias and promyelocytic leukemia; tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma; other tumors, including melanoma, seminoma, teratocarcinoma, neuroblastoma and glioma; tumors of the central and peripheral nervous system, including astrocytoma, neuroblastoma, glioma, and schwannomas; tumors of mesenchymal origin, including fibrosarcoma, rhabdomyosarcoma, and osteosarcoma; and other tumors, including melanoma, xeroderma pigmentosum, keratoacanthoma, seminoma, thyroid follicular cancer and teratocarcinoma. It is also contemplated that cancers caused by aberrations in apoptosis would also be treated by the methods and compositions of the invention. Such cancers may include but not be limited to follicular lymphomas, carcinomas with p53 mutations, hormone dependent tumors of the breast, prostate and ovary, and precancerous lesions such as familial adenomatous polyposis, and myelodysplastic syndromes. In specific embodiments, malignancy or dysproliferative changes (such as metaplasias and dysplasias), or hyperproliferative disorders, are treated or prevented by the methods and compositions of the invention in the ovary, bladder, breast, colon, lung, skin, pancreas, or uterus. In other specific embodiments, sarcoma, melanoma, or leukemia is treated or prevented by the methods and compositions of the invention.

[00256] In a specific embodiment, a molecule of the invention (*e.g.*, an antibody comprising a variant Fc region, or a therapeutic monoclonal antibody engineered according to the methods of the invention) inhibits or reduces the growth of primary tumor or metastasis of cancerous cells by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to the growth of primary tumor or metastasis in the absence of said molecule of the invention.

[00257]

10

5.4.1.1 COMBINATION THERAPY

[00258] The invention further encompasses administering the molecules of the invention in combination with other therapies known to those skilled in the art for the treatment or prevention of cancer, including but not limited to, current standard and experimental chemotherapies, hormonal therapies, biological therapies, immunotherapies, radiation therapies, or surgery. In some embodiments, the molecules of the invention may be administered in combination with a therapeutically or prophylactically effective amount of one or more anti-cancer agents, therapeutic antibodies (*e.g.*, antibodies listed in Table 3), or other agents known to those skilled in the art for the treatment and/or prevention of cancer (*See* Section 5.4.1.2).

20 [00259] In certain embodiments, one or more molecule of the invention is administered to a mammal, preferably a human, concurrently with one or more other therapeutic agents useful for the treatment of cancer. The term "concurrently" is not limited to the administration of prophylactic or therapeutic agents at exactly the same time, but rather it is meant that a molecule of the invention and the other agent are administered to a mammal in a sequence and within a time interval such that the molecule of the invention can act together with the other agent to provide an increased benefit than if they were administered otherwise. For example, each prophylactic or therapeutic agent (*e.g.*, chemotherapy, radiation therapy, hormonal therapy or biological therapy) may be administered at the same time or sequentially in any order at different points in time; however, if not administered at the same time, they should be administered sufficiently close in time so as to provide the desired therapeutic or prophylactic effect. Each therapeutic agent can be administered separately, in any appropriate form and by any suitable route. In various embodiments, the prophylactic or therapeutic agents are administered less than 1 hour apart, at about 1 hour apart, at about 1 hour to about 2 hours

apart, at about 2 hours to about 3 hours apart, at about 3 hours to about 4 hours apart, at about 4 hours to about 5 hours apart, at about 5 hours to about 6 hours apart, at about 6 hours to about 7 hours apart, at about 7 hours to about 8 hours apart, at about 8 hours to about 9 hours apart, at about 9 hours to about 10 hours apart, at about 10 hours to about 11 hours apart, at about 11 hours to about 12 hours apart, no more than 24 hours apart or no more than 48 hours apart. In preferred embodiments, two or more components are administered within the same patient visit.

[00260] In other embodiments, the prophylactic or therapeutic agents are administered at about 2 to 4 days apart, at about 4 to 6 days apart, at about 1 week apart, at about 1 to 2 weeks apart, or more than 2 weeks apart. In preferred embodiments, the prophylactic or therapeutic agents are administered in a time frame where both agents are still active. One skilled in the art would be able to determine such a time frame by determining the half life of the administered agents.

[00261] In certain embodiments, the prophylactic or therapeutic agents of the invention are cyclically administered to a subject. Cycling therapy involves the administration of a first agent for a period of time, followed by the administration of a second agent and/or third agent for a period of time and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or improves the efficacy of the treatment.

[00262] In certain embodiments, prophylactic or therapeutic agents are administered in a cycle of less than about 3 weeks, about once every two weeks, about once every 10 days or about once every week. One cycle can comprise the administration of a therapeutic or prophylactic agent by infusion over about 90 minutes every cycle, about 1 hour every cycle, about 45 minutes every cycle. Each cycle can comprise at least 1 week of rest, at least 2 weeks of rest, at least 3 weeks of rest. The number of cycles administered is from about 1 to about 12 cycles, more typically from about 2 to about 10 cycles, and more typically from about 2 to about 8 cycles.

[00263] In yet other embodiments, the therapeutic and prophylactic agents of the invention are administered in metronomic dosing regimens, either by continuous infusion or frequent administration without extended rest periods. Such metronomic administration can involve dosing at constant intervals without rest periods. Typically the therapeutic agents, in particular cytotoxic agents, are used at lower doses. Such dosing regimens encompass the chronic daily administration of relatively low doses for extended periods of time. In preferred embodiments, the use of lower doses can minimize toxic side effects and

eliminate rest periods. In certain embodiments, the therapeutic and prophylactic agents are delivered by chronic low-dose or continuous infusion ranging from about 24 hours to about 2 days, to about 1 week, to about 2 weeks, to about 3 weeks to about 1 month to about 2 months, to about 3 months, to about 4 months, to about 5 months, to about 6 months. The
5 scheduling of such dose regimens can be optimized by the skilled oncologist.

[00264] In other embodiments, courses of treatment are administered concurrently to a mammal, *i.e.*, individual doses of the therapeutics are administered separately yet within a time interval such that molecules of the invention can work together with the other agent or agents. For example, one component may be administered one time per week in
10 combination with the other components that may be administered one time every two weeks or one time every three weeks. In other words, the dosing regimens for the therapeutics are carried out concurrently even if the therapeutics are not administered simultaneously or within the same patient visit.

[00265] When used in combination with other prophylactic and/or therapeutic agents,
15 the molecules of the invention and the prophylactic and/or therapeutic agent can act additively or, more preferably, synergistically. In one embodiment, a molecule of the invention is administered concurrently with one or more therapeutic agents in the same pharmaceutical composition. In another embodiment, a molecule of the invention is administered concurrently with one or more other therapeutic agents in separate
20 pharmaceutical compositions. In still another embodiment, a molecule of the invention is administered prior to or subsequent to administration of another prophylactic or therapeutic agent. The invention contemplates administration of a molecule of the invention in combination with other prophylactic or therapeutic agents by the same or different routes of administration, *e.g.*, oral and parenteral. In certain embodiments, when a molecule of the
25 invention is administered concurrently with another prophylactic or therapeutic agent that potentially produces adverse side effects including, but not limited to, toxicity, the prophylactic or therapeutic agent can advantageously be administered at a dose that falls below the threshold that the adverse side effect is elicited.

[00266] The dosage amounts and frequencies of administration provided herein are
30 encompassed by the terms therapeutically effective and prophylactically effective. The dosage and frequency further will typically vary according to factors specific for each patient depending on the specific therapeutic or prophylactic agents administered, the severity and type of cancer, the route of administration, as well as age, body weight, response, and the past medical history of the patient. Suitable regimens can be selected by
35 one skilled in the art by considering such factors and by following, for example, dosages

reported in the literature and recommended in the *Physician's Desk Reference* (56th ed., 2002).

5.4.1.2 OTHER THERAPEUTIC/PROPHYLACTIC AGENTS

[00267] In a specific embodiment, the methods of the invention encompass the administration of one or more molecules of the invention with one or more therapeutic agents used for the treatment and/or prevention of cancer. In one embodiment, angiogenesis inhibitors may be administered in combination with the molecules of the invention. Angiogenesis inhibitors that can be used in the methods and compositions of the invention include but are not limited to: Angiostatin (plasminogen fragment); antiangiogenic antithrombin III; Angiozyme; ABT-627; Bay 12-9566; Benefin; Bevacizumab; BMS-275291; cartilage-derived inhibitor (CDI); CAI; CD59 complement fragment; CEP-7055; Col 3; Combretastatin A-4; Endostatin (collagen XVIII fragment); Fibronectin fragment; Gro-beta; Halofuginone; Heparinases; Heparin hexasaccharide fragment; HMV833; Human chorionic gonadotropin (hCG); IM-862; Interferon alpha/beta/gamma; Interferon inducible protein (IP-10); Interleukin-12; Kringle 5 (plasminogen fragment); Marimastat; Metalloproteinase inhibitors (TIMPs); 2-Methoxyestradiol; MMI 270 (CGS 27023A); MoAb IMC-1C11; Neovastat; NM-3; Panzem; PI-88; Placental ribonuclease inhibitor; Plasminogen activator inhibitor; Platelet factor-4 (PF4); Prinomastat; Prolactin 16kD fragment; Proliferin-related protein (PRP); PTK 787/ZK 222594; Retinoids; Solimastat; Squalamine; SS 3304; SU 5416; SU6668; SU11248; Tetrahydrocortisol-S; tetrathiomolybdate; thalidomide; Thrombospondin-1 (TSP-1); TNP-470; Transforming growth factor-beta (TGF-b); Vascuostatin; Vasostatin (calreticulin fragment); ZD6126; ZD 6474; farnesyl transferase inhibitors (FTI); and bisphosphonates.

[00268] Anti-cancer agents that can be used in combination with the molecules of the invention in the various embodiments of the invention, including pharmaceutical compositions and dosage forms and kits of the invention, include, but are not limited to: acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; aminoglutethimide; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; bropiramine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefingol; chlorambucil; cirolemycin; cisplatin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; dactinomycin; daunorubicin hydrochloride;

- decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziquone; docetaxel; doxorubicin; doxorubicin hydrochloride; droloxifene; droloxifene citrate; dromostanolone propionate; duazomycin; edatrexate; eflornithine hydrochloride; elsamitrucin; enloplatin; enpromate; epipropidine; epirubicin hydrochloride; erbulozole; esorubicin hydrochloride; estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; flurocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; ilmofofine; interleukin II (including recombinant interleukin II, or rIL2), interferon alfa-2a; interferon alfa-2b; interferon alfa-n1 ; interferon alfa-n3; interferon beta-I a; interferon gamma-I b; iproplatin; irinotecan hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride; lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprocol; maytansine; mechlorethamine hydrochloride; megestrol acetate; melengestrol acetate; melphalan; menogaril; mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedpa; mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane; mitoxantrone hydrochloride; mycophenolic acid; nocodazole; nogalamycin; ormaplatin; oxisuran; paclitaxel; pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide; pipobroman; piposulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium; porfiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride; pyrazofurin; riboprine; rogletimide; safingol; safingol hydrochloride; semustine; simtrazene; sparfosate sodium; sparsomycin; spirogermanium hydrochloride; spiromustine; spiroplatin; streptonigrin; streptozocin; sulofenur; talisomycin; tecogalan sodium; tegafur; teloxantrone hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprine; thioguanine; thiotepa; tiazofurin; tirapazamine; toremifene citrate; trestolone acetate; triciribine phosphate; trimetrexate; trimetrexate glucuronate; triptorelin; tubulozole hydrochloride; uracil mustard; uredepa; vapreotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine sulfate; vinepidine sulfate; vinglycinate sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; zorubicin hydrochloride. Other anti-cancer drugs include, but are not limited to: 20-epi-1,25 dihydroxyvitamin D3; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; anti-dorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen;

- antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin III derivatives; balanol; batimastat; BCR/ABL antagonists;
- 5 benzochlorins; benzoylstauroporine; beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; bropirimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives; canarypox IL-2; capecitabine; carboxamide-amino-triazole; carboxyamidotriazole; CaRest M3; CARN 700; cartilage
- 10 derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B; cetorelix; chlorins; chloroquinoxaline sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin analogue; conagenin; crambescidin 816; crisnatol; cryptophycin 8; cryptophycin A derivatives; curacin A; cyclopentantraquinones; cycloplatam; cypemycin;
- 15 cytarabine ocfosfate; cytolytic factor; cytostatin; daclicximab; decitabine; dehydrodidemnin B; deslorelin; dexamethasone; dexifosfamide; dexrazoxane; dexverapamil; diaziquone; didemnin B; didox; diethylnorspermine; dihydro-5-azacytidine; dihydrotaxol, 9-; dioxamycin; diphenyl spiromustine; docetaxel; docosanol; dolasetron; doxifluridine; droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine; edrecolomab;
- 20 eflornithine; elemene; emitefur; epirubicin; episteride; estramustine analogue; estrogen agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane; fadrozole; fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; flezelastine; fluasterone; fludarabine; fluorodaunorubicin hydrochloride; forfenimex; formestane; fostriecin; fotemustine; gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase
- 25 inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; heregulin; hexamethylene bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene; idramantone; ilmofofosine; ilomastat; imidazoacridones; imiquimod; immunostimulant peptides; insulin-like growth factor-1 receptor inhibitor; interferon agonists; interferons; interleukins; iobenguane; iododoxorubicin; ipomeanol, 4-; iroplact; irsogladine; isobengazole; isohomohalicondrin B;
- 30 itasetron; jasplakinolide; kahalalide F; lamellarin-N triacetate; lanreotide; leinamycin; lenograstim; lentinan sulfate; leptolstatin; letrozole; leukemia inhibiting factor; leukocyte alpha interferon; leuprolide+estrogen+progesterone; leuprorelin; levamisole; liarozole; linear polyamine analogue; lipophilic disaccharide peptide; lipophilic platinum compounds; lissoclinamide 7; lobaplatin; lombricine; lometrexol; lonidamine; losoxantrone; lovastatin;
- 35 loxoribine; lurtotecan; lutetium texaphyrin; lysofylline; lytic peptides; maitansine;

- mannostatin A; marimastat; masoprocol; maspin; matrilysin inhibitors; matrix metalloproteinase inhibitors; menogaril; merbarone; meterelin; methioninase; metoclopramide; MIF inhibitor; mifepristone; miltefosine; mirimostim; mismatched double stranded RNA; mitoguazone; mitolactol; mitomycin analogues; mitonafide; mitotoxin
- 5 fibroblast growth factor-saporin; mitoxantrone; mofarotene; molgramostim; monoclonal antibody, human chorionic gonadotrophin; monophosphoryl lipid A+myobacterium cell wall sk; mopidamol; multiple drug resistance gene inhibitor; multiple tumor suppressor 1-based therapy; mustard anticancer agent; mycaperoxide B; mycobacterial cell wall extract; myriaporone; N-acetyldinaline; N-substituted benzamides; nafarelin; nagrestip;
- 10 naloxone+pentazocine; napavin; naphterpin; nartograstim; nedaplatin; nemorubicin; neridronic acid; neutral endopeptidase; nilutamide; nisamycin; nitric oxide modulators; nitroxide antioxidant; nitrullyn; O6-benzylguanine; octreotide; okicenone; oligonucleotides; onapristone; ondansetron; ondansetron; oracin; oral cytokine inducer; ormaplatin; osaterone; oxaliplatin; oxaunomycin; paclitaxel; paclitaxel analogues; paclitaxel derivatives;
- 15 palauamine; palmitoylrhizoxin; pamidronic acid; panaxytriol; panomifene; parabactin; pazelliptine; pegaspargase; peldesine; pentosan polysulfate sodium; pentostatin; pentrozole; perflubron; perfosfamide; perillyl alcohol; phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride; pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum complex; platinum compounds;
- 20 platinum-triamine complex; porfimer sodium; porfiromycin; prednisone; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based immune modulator; protein kinase C inhibitor; protein kinase C inhibitors, microalgal; protein tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists;
- 25 raltitrexed; ramosetron; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII retinamide; roglitimide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safingol; saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; signal
- 30 transduction modulators; single chain antigen binding protein; sizofiran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin; spongistatin 1; squalamine; stem cell inhibitor; stem-cell division inhibitors; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista;
- 35 suramin; swainsonine; synthetic glycosaminoglycans; tallimustine; tamoxifen methiodide;

tauromustine; tazarotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors;
 temoporfin; temozolomide; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine;
 thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor
 agonist; thymotrinan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine;
 5 titanocene bichloride; topsentin; toremifene; totipotent stem cell factor; translation
 inhibitors; tretinoin; triacetyluridine; triciribine; trimetrexate; triptorelin; tropisetron;
 turosteride; tyrosine kinase inhibitors; tyrphostins; UBC inhibitors; ubenimex; urogenital
 sinus-derived growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin
 B; vector system, erythrocyte gene therapy; velaresol; veramine; verdins; verteporfin;
 10 vinorelbine; vinxaltine; vitaxin; vorozole; zanoterone; zeniplatin; zilascorb; and zinostatin
 stimalamer. Preferred additional anti-cancer drugs are 5-fluorouracil and leucovorin.

[00269] Examples of therapeutic antibodies that can be used in methods of the
 invention include but are not limited to ZENAPAX® (daclizumab) (Roche Pharmaceuticals,
 Switzerland) which is an immunosuppressive, humanized anti-CD25 monoclonal antibody
 15 for the prevention of acute renal allograft rejection; PANOREX™ which is a murine anti-
 17-IA cell surface antigen IgG2a antibody (Glaxo Wellcome/Centocor); BEC2 which is a
 murine anti-idiotypic (GD3 epitope) IgG antibody (ImClone System); IMC-C225 which is a
 chimeric anti-EGFR IgG antibody (ImClone System); VITAXIN™ which is a humanized
 anti- α V β 3 integrin antibody (Applied Molecular Evolution/MedImmune); Smart M195
 20 which is a humanized anti-CD33 IgG antibody (Protein Design Lab/Kanebo);
 LYMPHOCIDE™ which is a humanized anti-CD22 IgG antibody (Immunomedics); ICM3
 is a humanized anti-ICAM3 antibody (ICOS Pharm); IDEC-114 is a primatized anti-CD80
 antibody (IDEC Pharm/Mitsubishi); IDEC-131 is a humanized anti-CD40L antibody
 (IDEC/Eisai); IDEC-151 is a primatized anti-CD4 antibody (IDEC); IDEC-152 is a
 25 primatized anti-CD23 antibody (IDEC/Seikagaku); SMART anti-CD3 is a humanized anti-
 CD3 IgG (Protein Design Lab); 5G1.1 is a humanized anti-complement factor 5 (C5)
 antibody (Alexion Pharm); D2E7 is a humanized anti-TNF- α antibody (CAT/BASF);
 CDP870 is a humanized anti-TNF- α Fab fragment (Celltech); IDEC-151 is a primatized
 anti-CD4 IgG1 antibody (IDEC Pharm/SmithKline Beecham); MDX-CD4 is a human anti-
 30 CD4 IgG antibody (Medarex/Eisai/Genmab); CDP571 is a humanized anti-TNF- α IgG4
 antibody (Celltech); LDP-02 is a humanized anti- α 4 β 7 antibody (LeukoSite/Genentech);
 OrthoClone OKT4A is a humanized anti-CD4 IgG antibody (Ortho Biotech); ANTOVA™
 is a humanized anti-CD40L IgG antibody (Biogen); ANTEGREN™ is a humanized anti-
 VLA-4 IgG antibody (Elan); and CAT-152 is a human anti-TGF- β 2 antibody (Cambridge

Ab Tech). Other examples of therapeutic antibodies that can be used in accordance with the invention are presented in Table 6.

5.4.2 AUTOIMMUNE DISEASE AND INFLAMMATORY DISEASES

5 [00270] In some embodiments, molecules of the invention comprise a variant Fc region, having one or more amino acid modifications in one or more regions, which modification increases the affinity of the variant Fc region for Fc γ RIIB but decreases the affinity of the variant Fc region for Fc γ RIIA and/or Fc γ RIIA. Molecules of the invention with such binding characteristics are useful in regulating the immune response, *e.g.*, in
10 inhibiting the immune response in connection with autoimmune diseases or inflammatory diseases. Although not intending to be bound by any mechanism of action, molecules of the invention with an enhanced affinity for Fc γ RIIB and a decreased affinity for Fc γ RIIA and/or Fc γ RIIA may lead to dampening of the activating response to Fc γ R and inhibition of cellular responsiveness.

15 [00271] In some embodiments, a molecule of the invention comprising a variant Fc region is not an immunoglobulin, and comprises at least one amino acid modification which modification increases the affinity of the variant Fc region for Fc γ RIIB relative to a molecule comprising a wild-type Fc region. In other embodiments, said molecule further comprises one or more amino acid modifications, which modifications decreases the affinity
20 of the molecule for an activating Fc γ R. In some embodiments, the molecule is a soluble (*i.e.*, not membrane bound) Fc region. The invention contemplates other amino acid modifications within the soluble Fc region which modulate its affinity for various Fc receptors, including those known to one skilled in the art as described herein. In other embodiments, the molecule (*e.g.*, the Fc region comprising at least one or more amino acid
25 modification) is modified using techniques known to one skilled in the art and as described herein to increase the *in vivo* half life of the Fc region. Such molecules have therapeutic utility in treating and/or preventing an autoimmune disorder. Although not intending to be bound by any mechanism of actions, such molecules with enhanced affinity for Fc γ RIIB will lead to a dampening of the activating receptors and thus a dampening of the immune
30 response and have therapeutic efficacy for treating and/or preventing an autoimmune disorder.

[00272] In certain embodiments, the one or more amino acid modifications, which increase the affinity of the variant Fc region for Fc γ RIIB but decrease the affinity of the variant Fc region for Fc γ RIIA comprise a substitution at position 246 with threonine and at

position 396 with histidine; or a substitution at position 268 with aspartic acid and at position 318 with aspartic acid; or a substitution at position 217 with serine, at position 378 with valine, and at position 408 with arginine; or a substitution at position 375 with cysteine and at position 396 with leucine; or a substitution at position 246 with isoleucine and at position 334 with asparagine. In one embodiment, the one or more amino acid modifications, which increase the affinity of the variant Fc region for Fc γ RIIB but decrease the affinity of the variant Fc region for Fc γ RIIA comprise a substitution at position 247 with leucine. In another embodiment, the one or more amino acid modification, which increases the affinity of the variant Fc region for Fc γ RIIB but decreases the affinity of the variant Fc region for Fc γ RIIA comprise a substitution at position 372 with tyrosine. In yet another embodiment, the one or more amino acid modification, which increases the affinity of the variant Fc region for Fc γ RIIB but decreases the affinity of the variant Fc region for Fc γ RIIA comprise a substitution at position 326 with glutamic acid. In one embodiment, the one or more amino acid modification, which increases the affinity of the variant Fc region for Fc γ RIIB but decreases the affinity of the variant Fc region for Fc γ RIIA comprise a substitution at position 224 with leucine.

[00273] The variant Fc regions that have an enhanced affinity for Fc γ RIIB and a decreased affinity for Fc γ RIIA and/or Fc γ RII relative to a comparable molecule comprising a wild-type Fc region, may be used to treat or prevent autoimmune diseases or inflammatory diseases. The present invention provides methods of preventing, treating, or managing one or more symptoms associated with an autoimmune or inflammatory disorder in a subject, comprising administering to said subject a therapeutically or prophylactically effective amount of one or more molecules of the invention with variant Fc regions that have an enhanced affinity for Fc γ RIIB and a decreased affinity for Fc γ RIIA and or Fc γ RII relative to a comparable molecule comprising a wild type Fc region.

[00274] The invention also provides methods for preventing, treating, or managing one or more symptoms associated with an inflammatory disorder in a subject further comprising, administering to said subject a therapeutically or prophylactically effective amount of one or more anti-inflammatory agents. The invention also provides methods for preventing, treating, or managing one or more symptoms associated with an autoimmune disease further comprising, administering to said subject a therapeutically or prophylactically effective amount of one or more immunomodulatory agents. Section 5.4.2.1 provides non-limiting examples of anti-inflammatory agents and immunomodulatory agents.

[00275] Examples of autoimmune disorders that may be treated by administering the molecules of the present invention include, but are not limited to, alopecia areata, ankylosing spondylitis, antiphospholipid syndrome, autoimmune Addison's disease, autoimmune diseases of the adrenal gland, autoimmune hemolytic anemia, autoimmune hepatitis, autoimmune oophoritis and orchitis, autoimmune thrombocytopenia, Behcet's disease, bullous pemphigoid, cardiomyopathy, celiac sprue-dermatitis, chronic fatigue immune dysfunction syndrome (CFIDS), chronic inflammatory demyelinating polyneuropathy, Churg-Strauss syndrome, cicatricial pemphigoid, CREST syndrome, cold agglutinin disease, Crohn's disease, discoid lupus, essential mixed cryoglobulinemia, fibromyalgia-fibromyositis, glomerulonephritis, Graves' disease, Guillain-Barre, Hashimoto's thyroiditis, idiopathic pulmonary fibrosis, idiopathic thrombocytopenia purpura (ITP), IgA neuropathy, juvenile arthritis, lichen planus, lupus erythematosus, Ménière's disease, mixed connective tissue disease, multiple sclerosis, type 1 or immune-mediated diabetes mellitus, myasthenia gravis, pemphigus vulgaris, pernicious anemia, polyarteritis nodosa, polychondritis, polyglandular syndromes, polymyalgia rheumatica, polymyositis and dermatomyositis, primary agammaglobulinemia, primary biliary cirrhosis, psoriasis, psoriatic arthritis, Raynaud's phenomenon, Reiter's syndrome, Rheumatoid arthritis, sarcoidosis, scleroderma, Sjögren's syndrome, stiff-man syndrome, systemic lupus erythematosus, lupus erythematosus, takayasu arteritis, temporal arteritis/ giant cell arteritis, ulcerative colitis, uveitis, vasculitides such as dermatitis herpetiformis vasculitis, vitiligo, and Wegener's granulomatosis. Examples of inflammatory disorders include, but are not limited to, asthma, encephalitis, inflammatory bowel disease, chronic obstructive pulmonary disease (COPD), allergic disorders, septic shock, pulmonary fibrosis, undifferentiated spondyloarthropathy, undifferentiated arthropathy, arthritis, inflammatory osteolysis, and chronic inflammation resulting from chronic viral or bacteria infections. As described herein in Section 2.2.2, some autoimmune disorders are associated with an inflammatory condition. Thus, there is overlap between what is considered an autoimmune disorder and an inflammatory disorder. Therefore, some autoimmune disorders may also be characterized as inflammatory disorders. Examples of inflammatory disorders which can be prevented, treated or managed in accordance with the methods of the invention include, but are not limited to, asthma, encephalitis, inflammatory bowel disease, chronic obstructive pulmonary disease (COPD), allergic disorders, septic shock, pulmonary fibrosis, undifferentiated spondyloarthropathy, undifferentiated arthropathy, arthritis, inflammatory osteolysis, and chronic inflammation resulting from chronic viral or bacteria infections.

[00276] Molecules of the invention with variant Fc regions that have an enhanced affinity for Fc γ RIIB and a decreased affinity for Fc γ RIIA relative to a comparable molecule comprising a wild-type Fc region can also be used to reduce the inflammation experienced by animals, particularly mammals, with inflammatory disorders. In a specific embodiment, a molecule of the invention reduces the inflammation in an animal by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to the inflammation in an animal, which is not administered the said molecule.

10 [00277] Molecules of the invention with variant Fc regions that have an enhanced affinity for Fc γ RIIB and a decreased affinity for Fc γ RIIA relative to a comparable molecule comprising a wild-type Fc region can also be used to prevent the rejection of transplants.

[00278] The invention further contemplates engineering any of the antibodies known in the art for the treatment and/or prevention of autoimmune disease or inflammatory disease, so that the antibodies comprise a variant Fc region comprising one or more amino acid modifications, which have been identified by the methods of the invention to have an enhanced affinity for Fc γ RIIB and a decreased affinity for Fc γ RIIA relative to a comparable molecule comprising a wild type Fc region. A non-limiting example of the antibodies that are used for the treatment or prevention of inflammatory disorders which can be engineered according to the invention is presented in Table 7A, and a non-limiting example of the antibodies that are used for the treatment or prevention of autoimmune disorder is presented in Table 7B.

25 **TABLE 7A: ANTIBODIES FOR INFLAMMATORY DISEASES AND AUTOIMMUNE DISEASES THAT CAN ENGINEERED IN ACCORDANCE WITH THE INVENTION.**

Antibody Name	Target Antigen	Product Type	Isotype	Sponsors	Indication
5G1.1	Complement (C5)	Humanized	IgG	Alexion Pharm Inc	Rheumatoid Arthritis
5G1.1	Complement (C5)	Humanized	IgG	Alexion Pharm Inc	SLE
5G1.1	Complement (C5)	Humanized	IgG	Alexion Pharm Inc	Nephritis
5G1.1-SC	Complement (C5)	Humanized	ScFv	Alexion Pharm Inc	Cardiopulmonary Bypass

Antibody Name	Target Antigen	Product Type	Isotype	Sponsors	Indication
5G1.1-SC	Complement (C5)	Humanized	ScFv	Alexion Pharm Inc	Myocardial Infarction
5G1.1-SC	Complement (C5)	Humanized	ScFv	Alexion Pharm Inc	Angioplasty
ABX-CBL	CBL	Human		Abgenix Inc	GvHD
ABX-CBL	CD147	Murine	IgG	Abgenix Inc	Allograft rejection
ABX-IL8	IL-8	Human	IgG2	Abgenix Inc	Psoriasis
Antegren	VLA-4	Humanized	IgG	Athena/Elan	Multiple Sclerosis
Anti-CD11a	CD11a	Humanized	IgG1	Genentech Inc/Xoma	Psoriasis
Anti-CD18	CD18	Humanized	Fab'2	Genentech Inc	Myocardial infarction
Anti-LFA1	CD18	Murine	Fab'2	Pasteur-Merieux/Immunotech	Allograft rejection
Antova	CD40L	Humanized	IgG	Biogen	Allograft rejection
Antova	CD40L	Humanized	IgG	Biogen	SLE
BTI-322	CD2	Rat	IgG	Medimmune Inc	GvHD, Psoriasis
CDP571	TNF-alpha	Humanized	IgG4	Celltech	Crohn's
CDP571	TNF-alpha	Humanized	IgG4	Celltech	Rheumatoid Arthritis
CDP850	E-selectin	Humanized		Celltech	Psoriasis
Corsevin M	Fact VII	Chimeric		Centocor	Anticoagulant
D2E7	TNF-alpha	Human		CAT/BASF	Rheumatoid Arthritis
Hu23F2G	CD11/18	Humanized		ICOS Pharm Inc	Multiple Sclerosis
Hu23F2G	CD11/18	Humanized	IgG	ICOS Pharm Inc	Stroke
IC14	CD14			ICOS Pharm Inc	Toxic shock
ICM3	ICAM-3	Humanized		ICOS Pharm Inc	Psoriasis
IDEC-114	CD80	Primates		IDEC Pharm/Mitsubishi	Psoriasis
IDEC-131	CD40L	Humanized		IDEC	SLE

Antibody Name	Target Antigen	Product Type	Isotype	Sponsors	Indication
				Pharm/Eisai	
IDEC-131	CD40L	Humanized		IDEC Pharm/Eisai	Multiple Sclerosis
IDEC-151	CD4	Primatised	IgG1	IDEC Pharm/Glaxo SmithKline	Rheumatoid Arthritis
IDEC-152	CD23	Primatised		IDEC Pharm	Asthma/Allergy
Infliximab	TNF-alpha	Chimeric	IgG1	Centocor	Rheumatoid Arthritis
Infliximab	TNF-alpha	Chimeric	IgG1	Centocor	Crohn's
LDP-01	beta2-integrin	Humanized	IgG	Millennium Inc (LeukoSite Inc.)	Stroke
LDP-01	beta2-integrin	Humanized	IgG	Millennium Inc (LeukoSite Inc.)	Allograft rejection
LDP-02	alpha4beta7	Humanized		Millennium Inc (LeukoSite Inc.)	Ulcerative Colitis
MAK-195F	TNF alpha	Murine	Fab'2	Knoll Pharm, BASF	Toxic shock
MDX-33	CD64 (FcR)	Human		Medarex/Cent eon	Autoimmune haematological disorders
MDX-CD4	CD4	Human	IgG	Medarex/Eisai / Genmab	Rheumatoid Arthritis
MEDI-507	CD2	Humanized		Medimmune Inc	Psoriasis
MEDI-507	CD2	Humanized		Medimmune Inc	GvHD
OKT4A	CD4	Humanized	IgG	Ortho Biotech	Allograft rejection
OrthoClone OKT4A	CD4	Humanized	IgG	Ortho Biotech	Autoimmune disease
Orthoclone/anti-CD3 OKT3	CD3	Murine	mIgG2a	Ortho Biotech	Allograft rejection

Antibody Name	Target Antigen	Product Type	Isotype	Sponsors	Indication
RepPro/Abciximab	gpIIbIIIa	Chimeric	Fab	Centocor/Lilly	Complications of coronary angioplasty
rhuMab-E25	IgE	Humanized	IgG1	Genentech/Novartis/Tanox Biosystems	Asthma/Allergy
SB-240563	IL5	Humanized		GlaxoSmithKline	Asthma/Allergy
SB-240683	IL-4	Humanized		GlaxoSmithKline	Asthma/Allergy
SCH55700	IL-5	Humanized		Celltech/Schering	Asthma/Allergy
Simulect	CD25	Chimeric	IgG1	Novartis Pharm	Allograft rejection
SMART a-CD3	CD3	Humanized		Protein Design Lab	Autoimmune disease
SMART a-CD3	CD3	Humanized		Protein Design Lab	Allograft rejection
SMART a-CD3	CD3	Humanized	IgG	Protein Design Lab	Psoriasis
Zenapax	CD25	Humanized	IgG1	Protein Design Lab/Hoffman-La Roche	Allograft rejection

TABLE 7B: ANTIBODIES FOR AUTOIMMUNE DISORDERS THAT CAN BE ENGINEERED IN ACCORDANCE WITH THE INVENTION

Antibody	Indication	Target Antigen
ABX-RB2		antibody to CBL antigen on T cells, B cells and NK cells fully human antibody from the Xenomouse
5c8 (Anti CD-40 ligand antibody)	Phase II trials were halted in Oct. 99 examine "adverse events"	CD-40
IDEC 131	systemic lupus erythematosus (SLE)	anti CD40 humanized
IDEC 151	rheumatoid arthritis	primatized ; anti-CD4
IDEC 152	Asthma	primatized; anti-CD23
IDEC 114	Psoriasis	primatized anti-CD80
MEDI-507	rheumatoid arthritis; multiple sclerosis Crohn's disease Psoriasis	anti-CD2
LDP-02 (anti-b7 mAb)	inflammatory bowel disease Chron's disease ulcerative colitis	a4b7 integrin receptor on white blood cells (leukocytes)
SMART Anti-Gamma Interferon antibody	autoimmune disorders	Anti-Gamma Interferon
Verteportin	rheumatoid arthritis	
MDX-33	blood disorders caused by autoimmune reactions Idiopathic Thrombocytopenia Purpura (ITP) autoimmune hemolytic anemia	monoclonal antibody against FcRI receptors
MDX-CD4	treat rheumatoid arthritis and other autoimmunity	monoclonal antibody against CD4 receptor molecule
VX-497	autoimmune disorders multiple sclerosis rheumatoid arthritis inflammatory bowel disease lupus psoriasis	inhibitor of inosine monophosphate dehydrogenase (enzyme needed to make new RNA and DNA used in production of nucleotides needed for lymphocyte proliferation)
VX-740	rheumatoid arthritis	inhibitor of ICE

Antibody	Indication	Target Antigen
		interleukin-1 beta (converting enzyme controls pathways leading to aggressive immune response)
VX-745	specific to inflammation involved in chemical signalling of immune response onset and progression of inflammation	inhibitor of P38MAP kinase mitogen activated protein kinase
Enbrel (etanercept)		targets TNF (tumor necrosis factor)
IL-8		fully human monoclonal antibody against IL-8 (interleukin 8)
Apogen MP4		recombinant antigen selectively destroys disease associated T-cells induces apoptosis T-cells eliminated by programmed cell death no longer attack body's own cells specific apogens target specific T-cells

5.4.2.1 IMMUNOMODULATORY AGENTS AND ANTI-INFLAMMATORY AGENTS

[00279] The present invention provides methods of treatment for autoimmune diseases and inflammatory diseases comprising administration of the molecules with variant Fc regions having an enhanced affinity for Fc γ RIIB and a decreased affinity for Fc γ RIIA and/or Fc γ RIIA in conjunction with other treatment agents. Examples of immunomodulatory agents include, but are not limited to, methothrexate, ENBREL, REMICADETM, leflunomide, cyclophosphamide, cyclosporine A, and macrolide antibiotics (e.g., FK506 (tacrolimus)), methylprednisolone (MP), corticosteroids, steroids, mycophenolate mofetil, rapamycin (sirolimus), mizoribine, deoxyspergualin, brequinar, malononitriloamindes (e.g., leflunamide), T cell receptor modulators, and cytokine receptor modulators.

[00280] Anti-inflammatory agents have exhibited success in treatment of inflammatory and autoimmune disorders and are now a common and a standard treatment for such disorders. Any anti-inflammatory agent well-known to one of skill in the art can be used in the methods of the invention. Non-limiting examples of anti-inflammatory agents

include non-steroidal anti-inflammatory drugs (NSAIDs), steroidal anti-inflammatory drugs, beta-agonists, anticholinergic agents, and methyl xanthines. Examples of NSAIDs include, but are not limited to, aspirin, ibuprofen, celecoxib (CELEBREX™), diclofenac (VOLTAREN™), etodolac (LODINE™), fenoprofen (NALFON™), indomethacin (INDOCIN™), ketoralac (TORADOL™), oxaprozin (DAYPRO™), nabumentone (RELAFEN™), sulindac (CLINORIL™), tolmentin (TOLECTIN™), rofecoxib (VIOXX™), naproxen (ALEVE™, NAPROSYN™), ketoprofen (ACTRON™) and nabumetone (RELAFEN™). Such NSAIDs function by inhibiting a cyclooxygenase enzyme (e.g., COX-1 and/or COX-2). Examples of steroidal anti-inflammatory drugs include, but are not limited to, glucocorticoids, dexamethasone (DECADRON™), cortisone, hydrocortisone, prednisone (DELTASONE™), prednisolone, triamcinolone, azulfidine, and eicosanoids such as prostaglandins, thromboxanes, and leukotrienes.

5.4.3 INFECTIOUS DISEASE

[00281] The invention also encompasses methods for treating or preventing an infectious disease in a subject comprising administering a therapeutically or prophylactically effective amount of one or more molecules of the invention. Infectious diseases that can be treated or prevented by the molecules of the invention are caused by infectious agents including but not limited to viruses, bacteria, fungi, protozoa, and viruses.

[00282] Viral diseases that can be treated or prevented using the molecules of the invention in conjunction with the methods of the present invention include, but are not limited to, those caused by hepatitis type A, hepatitis type B, hepatitis type C, influenza, varicella, adenovirus, herpes simplex type I (HSV-I), herpes simplex type II (HSV-II), rinderpest, rhinovirus, echovirus, rotavirus, respiratory syncytial virus, papilloma virus, papova virus, cytomegalovirus, echinovirus, arbovirus, huntavirus, coxsackie virus, mumps virus, measles virus, rubella virus, polio virus, small pox, Epstein Barr virus, human immunodeficiency virus type I (HIV-I), human immunodeficiency virus type II (HIV-II), and agents of viral diseases such as viral meningitis, encephalitis, dengue or small pox.

[00283] Bacterial diseases that can be treated or prevented using the molecules of the invention in conjunction with the methods of the present invention, that are caused by bacteria include, but are not limited to, mycobacteria rickettsia, mycoplasma, neisseria, S. pneumonia, Borrelia burgdorferi (Lyme disease), Bacillus anthracis (anthrax), tetanus, streptococcus, staphylococcus, mycobacterium, tetanus, pertissus, cholera, plague, diptheria, chlamydia, S. aureus and legionella.

[00284] Protozoal diseases that can be treated or prevented using the molecules of the invention in conjunction with the methods of the present invention, that are caused by protozoa include, but are not limited to, leishmania, kokzidioa, trypanosoma or malaria.

[00285] Parasitic diseases that can be treated or prevented using the molecules of the invention in conjunction with the methods of the present invention, that are caused by parasites include, but are not limited to, chlamydia and rickettsia.

[00286] According to one aspect of the invention, molecules of the invention comprising variant Fc regions have an enhanced antibody effector function towards an infectious agent, *e.g.*, a pathogenic protein, relative to a comparable molecule comprising a wild-type Fc region. In a specific embodiment, molecules of the invention enhance the efficacy of treatment of an infectious disease by enhancing phagocytosis and/or opsonization of the infectious agent causing the infectious disease. In another specific embodiment, molecules of the invention enhance the efficacy of treatment of an infectious disease by enhancing ADCC of infected cells causing the infectious disease.

[00287] In some embodiments, the molecules of the invention may be administered in combination with a therapeutically or prophylactically effective amount of one or additional therapeutic agents known to those skilled in the art for the treatment and/or prevention of an infectious disease. The invention contemplates the use of the molecules of the invention in combination with antibiotics known to those skilled in the art for the treatment and or prevention of an infectious disease. Antibiotics that can be used in combination with the molecules of the invention include, but are not limited to, macrolide (*e.g.*, tobramycin (Tobi®)), a cephalosporin (*e.g.*, cephalexin (Keflex®), cephadrine (Velosef®), cefuroxime (Ceftin®), cefprozil (Cefzil®), cefaclor (Ceclor®), cefixime (Suprax®) or cefadroxil (Duricef®)), a clarithromycin (*e.g.*, clarithromycin (Biaxin®)), an erythromycin (*e.g.*, erythromycin (EMycin®)), a penicillin (*e.g.*, penicillin V (V-Cillin K® or Pen Vee K®)) or a quinolone (*e.g.*, ofloxacin (Floxin®), ciprofloxacin (Cipro®) or norfloxacin (Noroxin®)), aminoglycoside antibiotics (*e.g.*, apramycin, arbekacin, bambermycins, butirosin, dibekacin, neomycin, neomycin, undecylenate, netilmicin, paromomycin, ribostamycin, sisomicin, and spectinomycin), amphenicol antibiotics (*e.g.*, azidamfenicol, chloramphenicol, florfenicol, and thiamphenicol), ansamycin antibiotics (*e.g.*, rifamide and rifampin), carbacephem (*e.g.*, loracarbef), carbapenems (*e.g.*, biapenem and imipenem), cephalosporins (*e.g.*, cefaclor, cefadroxil, cefamandole, cefatrizine, cefazedone, cefozopran, cefpimizole, cefpiramide, and cefpirome), cephamycins (*e.g.*, cefbuperazone, cefmetazole, and cefminox), monobactams (*e.g.*, aztreonam, carumonam, and tigemonam), oxacephem (*e.g.*, flomoxef, and moxalactam), penicillins (*e.g.*,

amdinocillin, amdinocillin pivoxil, amoxicillin, bacampicillin, benzylpenicillinic acid, benzylpenicillin sodium, epicillin, fenbenicillin, floxacillin, penamccillin, penethamate hydriodide, penicillin o-benethamine, penicillin O, penicillin V, penicillin V benzathine, penicillin V hydrabamine, penimepicycline, and phencihicillin potassium), lincosamides (e.g., clindamycin, and lincomycin), amphomycin, bacitracin, capreomycin, colistin, enduracidin, enviomycin, tetracyclines (e.g., apicycline, chlortetracycline, clomocycline, and demeclocycline), 2,4-diaminopyrimidines (e.g., brodimoprim), nitrofurans (e.g., furaltadone, and furazolium chloride), quinolones and analogs thereof (e.g., cinoxacin,, clinafloxacin, flumequine, and grepagloxacin), sulfonamides (e.g., acetyl

5 (e.g., clindamycin, and lincomycin), amphomycin, bacitracin, capreomycin, colistin, enduracidin, enviomycin, tetracyclines (e.g., apicycline, chlortetracycline, clomocycline, and demeclocycline), 2,4-diaminopyrimidines (e.g., brodimoprim), nitrofurans (e.g., furaltadone, and furazolium chloride), quinolones and analogs thereof (e.g., cinoxacin,, clinafloxacin, flumequine, and grepagloxacin), sulfonamides (e.g., acetyl

10 sulfamethoxypyrazine, benzylsulfamide, noprilsulfamide, phthalylsulfacetamide, sulfachrysoidine, and sulfacytine), sulfones (e.g., diathymosulfone, glucosulfone sodium, and solasulfone), cycloserine, mupirocin and tuberin.

[00288] In certain embodiments, the molecules of the invention can be administered in combination with a therapeutically or prophylactically effective amount of one or more

15 antifungal agents. Antifungal agents that can be used in combination with the molecules of the invention include but are not limited to amphotericin B, itraconazole, ketoconazole, fluconazole, intrathecal, flucytosine, miconazole, butoconazole, clotrimazole, nystatin, terconazole, tioconazole, ciclopirox, econazole, haloprogrin, naftifine, terbinafine, undecylenate, and griseofulvin.

20 [00289] In some embodiments, the molecules of the invention can be administered in combination with a therapeutically or prophylactically effective amount of one or more anti-viral agent. Useful anti-viral agents that can be used in combination with the molecules of the invention include, but are not limited to, protease inhibitors, nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors and nucleoside

25 analogs. Examples of antiviral agents include but are not limited to zidovudine, acyclovir, gangcyclovir, vidarabine, idoxuridine, trifluridine, and ribavirin, as well as foscarnet, amantadine, rimantadine, saquinavir, indinavir, amprenavir, lopinavir, ritonavir, the alpha-interferons; adefovir, clevidine, entecavir, pleconaril.

5.5 VACCINE THERAPY

30 [00290] The invention further encompasses using a composition of the invention to induce an immune response against an antigenic or immunogenic agent, including but not limited to cancer antigens and infectious disease antigens (examples of which are disclosed *infra*). The vaccine compositions of the invention comprise one or more antigenic or immunogenic agents to which an immune response is desired, wherein the one or more

antigenic or immunogenic agents is coated with a variant antibody of the invention that has an enhanced affinity to FcγRIIIA. Although not intending to be bound by a particular mechanism of action, coating an antigenic or immunogenic agent with a variant antibody of the invention that has an enhanced affinity to FcγRIIIA, enhances the immune response to the desired antigenic or immunogenic agent by inducing humoral and cell-mediated responses. The vaccine compositions of the invention are particularly effective in eliciting an immune response, preferably a protective immune response against the antigenic or immunogenic agent.

[00291] In some embodiments, the antigenic or immunogenic agent in the vaccine compositions of the invention comprise a virus against which an immune response is desired. The viruses may be recombinant or chimeric, and are preferably attenuated. Production of recombinant, chimeric, and attenuated viruses may be performed using standard methods known to one skilled in the art. The invention encompasses a live recombinant viral vaccine or an inactivated recombinant viral vaccine to be formulated in accordance with the invention. A live vaccine may be preferred because multiplication in the host leads to a prolonged stimulus of similar kind and magnitude to that occurring in natural infections, and therefore, confers substantial, long-lasting immunity. Production of such live recombinant virus vaccine formulations may be accomplished using conventional methods involving propagation of the virus in cell culture or in the allantois of the chick embryo followed by purification.

[00292] In a specific embodiment, the recombinant virus is non-pathogenic to the subject to which it is administered. In this regard, the use of genetically engineered viruses for vaccine purposes may require the presence of attenuation characteristics in these strains. The introduction of appropriate mutations (*e.g.*, deletions) into the templates used for transfection may provide the novel viruses with attenuation characteristics. For example, specific missense mutations which are associated with temperature sensitivity or cold adaption can be made into deletion mutations. These mutations should be more stable than the point mutations associated with cold or temperature sensitive mutants and reversion frequencies should be extremely low. Recombinant DNA technologies for engineering recombinant viruses are known in the art and encompassed in the invention. For example, techniques for modifying negative strand RNA viruses are known in the art, see, *e.g.*, U.S. Patent No. 5,166,057, which is incorporated herein by reference in its entirety.

[00293] Alternatively, chimeric viruses with "suicide" characteristics may be constructed for use in the intradermal vaccine formulations of the invention. Such viruses would go through only one or a few rounds of replication within the host. When used as a

- vaccine, the recombinant virus would go through limited replication cycle(s) and induce a sufficient level of immune response but it would not go further in the human host and cause disease. Alternatively, inactivated (killed) virus may be formulated in accordance with the invention. Inactivated vaccine formulations may be prepared using conventional techniques
- 5 to "kill" the chimeric viruses. Inactivated vaccines are "dead" in the sense that their infectivity has been destroyed. Ideally, the infectivity of the virus is destroyed without affecting its immunogenicity. In order to prepare inactivated vaccines, the chimeric virus may be grown in cell culture or in the allantois of the chick embryo, purified by zonal ultracentrifugation, inactivated by formaldehyde or β -propiolactone, and pooled.
- 10 [00294] In certain embodiments, completely foreign epitopes, including antigens derived from other viral or non-viral pathogens can be engineered into the virus for use in the intradermal vaccine formulations of the invention. For example, antigens of non-related viruses such as HIV (gp160, gp120, gp41) parasite antigens (e.g., malaria), bacterial or fungal antigens or tumor antigens can be engineered into the attenuated strain.
- 15 [00295] Virtually any heterologous gene sequence may be constructed into the chimeric viruses of the invention for use in the intradermal vaccine formulations. Preferably, heterologous gene sequences are moieties and peptides that act as biological response modifiers. Preferably, epitopes that induce a protective immune response to any of a variety of pathogens, or antigens that bind neutralizing antibodies may be expressed by or
- 20 as part of the chimeric viruses. For example, heterologous gene sequences that can be constructed into the chimeric viruses of the invention include, but are not limited to, influenza and parainfluenza hemagglutinin neuraminidase and fusion glycoproteins such as the HN and F genes of human PIV3. In yet another embodiment, heterologous gene sequences that can be engineered into the chimeric viruses include those that encode
- 25 proteins with immuno-modulating activities. Examples of immuno-modulating proteins include, but are not limited to, cytokines, interferon type 1, gamma interferon, colony stimulating factors, interleukin -1, -2, -4, -5, -6, -12, and antagonists of these agents.
- [00296] In yet other embodiments, the invention encompasses pathogenic cells or viruses, preferably attenuated viruses, which express the variant antibody on their surface.
- 30 [00297] In alternative embodiments, the vaccine compositions of the invention comprise a fusion polypeptide wherein an antigenic or immunogenic agent is operatively linked to a variant antibody of the invention that has an enhanced affinity for Fc γ RIIIA. Engineering fusion polypeptides for use in the vaccine compositions of the invention is performed using routine recombinant DNA technology methods and is within the level of
- 35 ordinary skill.

[00298] The invention further encompasses methods to induce tolerance in a subject by administering a composition of the invention. Preferably a composition suitable for inducing tolerance in a subject, comprises an antigenic or immunogenic agent coated with a variant antibody of the invention, wherein the variant antibody has a higher affinity to Fc γ RIIB. Although not intending to be bound by a particular mechanism of action, such compositions are effective in inducing tolerance by activating the Fc γ RIIB mediated inhibitory pathway.

5.6 COMPOSITIONS AND METHODS OF ADMINISTERING

[00299] The invention provides methods and pharmaceutical compositions comprising molecules of the invention (*i.e.*, antibodies, polypeptides) comprising variant Fc regions. The invention also provides methods of treatment, prophylaxis, and amelioration of one or more symptoms associated with a disease, disorder or infection by administering to a subject an effective amount of a fusion protein or a conjugated molecule of the invention, or a pharmaceutical composition comprising a fusion protein or a conjugated molecule of the invention. In a preferred aspect, an antibody, a fusion protein, or a conjugated molecule, is substantially purified (*i.e.*, substantially free from substances that limit its effect or produce undesired side-effects). In a specific embodiment, the subject is an animal, preferably a mammal such as non-primate (*e.g.*, cows, pigs, horses, cats, dogs, rats *etc.*) and a primate (*e.g.*, monkey such as, a cynomolgous monkey and a human). In a preferred embodiment, the subject is a human. In yet another preferred embodiment, the antibody of the invention is from the same species as the subject.

[00300] Various delivery systems are known and can be used to administer a composition comprising molecules of the invention (*i.e.*, antibodies, polypeptides), comprising variant Fc regions, *e.g.*, encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the antibody or fusion protein, receptor-mediated endocytosis (*See, e.g.*, Wu and Wu, 1987, *J. Biol. Chem.* 262:4429-4432), construction of a nucleic acid as part of a retroviral or other vector, *etc.* Methods of administering a molecule of the invention include, but are not limited to, parenteral administration (*e.g.*, intradermal, intramuscular, intraperitoneal, intravenous and subcutaneous), epidural, and mucosal (*e.g.*, intranasal and oral routes). In a specific embodiment, the molecules of the invention are administered intramuscularly, intravenously, or subcutaneously. The compositions may be administered by any convenient route, for example, by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (*e.g.*, oral mucosa, rectal and intestinal mucosa, *etc.*)

and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, pulmonary administration can also be employed, *e.g.*, by use of an inhaler or nebulizer, and formulation with an aerosolizing agent. *See, e.g.*, U.S. Patent Nos. 6,019,968; 5,985,320; 5,985,309; 5,934,272; 5,874,064; 5,855,913; 5,290,540; 5 and 4,880,078; and PCT Publication Nos. WO 92/19244; WO 97/32572; WO 97/44013; WO 98/31346; and WO 99/66903, each of which is incorporated herein by reference in its entirety.

[00301] The invention also provides that the molecules of the invention (*i.e.*, antibodies, polypeptides) comprising variant Fc regions, are packaged in a hermetically sealed container such as an ampoule or sachette indicating the quantity of antibody. In one embodiment, the molecules of the invention are supplied as a dry sterilized lyophilized powder or water free concentrate in a hermetically sealed container and can be reconstituted, *e.g.*, with water or saline to the appropriate concentration for administration to a subject. Preferably, the molecules of the invention are supplied as a dry sterile lyophilized powder in a hermetically sealed container at a unit dosage of at least 5 mg, more preferably at least 10 mg, at least 15 mg, at least 25 mg, at least 35 mg, at least 45 mg, at least 50 mg, or at least 75 mg. The lyophilized molecules of the invention should be stored at between 2 and 8°C in their original container and the molecules should be administered within 12 hours, preferably within 6 hours, within 5 hours, within 3 hours, or within 1 hour after being reconstituted. In an alternative embodiment, molecules of the invention are supplied in liquid form in a hermetically sealed container indicating the quantity and concentration of the molecule, fusion protein, or conjugated molecule. Preferably, the liquid form of the molecules of the invention are supplied in a hermetically sealed container at least 1 mg/ml, more preferably at least 2.5 mg/ml; at least 5 mg/ml, at least 8 mg/ml, at least 10 mg/ml, at least 15 mg/kg, at least 25 mg/ml, at least 50 mg/ml, at least 100 mg/ml, at least 150 mg/ml, 25 at least 200 mg/ml of the molecules.

[00302] The amount of the composition of the invention which will be effective in the treatment, prevention or amelioration of one or more symptoms associated with a disorder can be determined by standard clinical techniques. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the condition, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

[00303] For antibodies encompassed by the invention, the dosage administered to a patient is typically 0.0001 mg/kg to 100 mg/kg of the patient's body weight. Preferably, the 35

dosage administered to a patient is between 0.0001 mg/kg and 20 mg/kg, 0.0001 mg/kg and 10 mg/kg, 0.0001 mg/kg and 5 mg/kg, 0.0001 and 2 mg/kg, 0.0001 and 1 mg/kg, 0.0001 mg/kg and 0.75 mg/kg, 0.0001 mg/kg and 0.5 mg/kg, 0.0001 mg/kg to 0.25 mg/kg, 0.0001 to 0.15 mg/kg, 0.0001 to 0.10 mg/kg, 0.001 to 0.5 mg/kg, 0.01 to 0.25 mg/kg or 0.01 to 0.10 mg/kg of the patient's body weight. Generally, human antibodies have a longer half-life within the human body than antibodies from other species due to the immune response to the foreign polypeptides. Thus, lower dosages of human antibodies and less frequent administration is often possible. Further, the dosage and frequency of administration of antibodies of the invention or fragments thereof may be reduced by enhancing uptake and tissue penetration of the antibodies by modifications such as, for example, lipidation.

[00304] In one embodiment, the dosage of the molecules of the invention administered to a patient are 0.01mg to 1000mg/day, when used as single agent therapy. In another embodiment the molecules of the invention are used in combination with other therapeutic compositions and the dosage administered to a patient are lower than when said molecules are used as a single agent therapy.

[00305] In a specific embodiment, it may be desirable to administer the pharmaceutical compositions of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion, by injection, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. Preferably, when administering a molecule of the invention, care must be taken to use materials to which the molecule does not absorb.

[00306] In another embodiment, the compositions can be delivered in a vesicle, in particular a liposome (*See Langer, Science 249:1527-1533 (1990); Treat et al., in Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353- 365 (1989); Lopez-Berestein, *ibid.*, pp. 3 17-327; *see generally ibid.*).

[00307] In yet another embodiment, the compositions can be delivered in a controlled release or sustained release system. Any technique known to one of skill in the art can be used to produce sustained release formulations comprising one or more molecules of the invention. *See, e.g.*, U.S. Patent No. 4,526,938; PCT publication WO 91/05548; PCT publication WO 96/20698; Ning *et al.*, 1996, "Intratumoral Radioimmunotherapy of a Human Colon Cancer Xenograft Using a Sustained-Release Gel," *Radiotherapy & Oncology* 39:179-189, Song *et al.*, 1995, "Antibody Mediated Lung Targeting of Long-Circulating Emulsions," *PDA Journal of Pharmaceutical Science & Technology*

50:372-397; Cleek *et al.*, 1997, "Biodegradable Polymeric Carriers for a bFGF Antibody for Cardiovascular Application," *Pro. Int'l. Symp. Control. Rel. Bioact. Mater.* 24:853-854; and Lam *et al.*, 1997, "Microencapsulation of Recombinant Humanized Monoclonal Antibody for Local Delivery," *Proc. Int'l. Symp. Control Rel. Bioact. Mater.* 24:759-760, each of which is incorporated herein by reference in its entirety. In one embodiment, a pump may be used in a controlled release system (*See* Langer, *supra*; Sefton, 1987, *CRC Crit. Ref. Biomed. Eng.* 14:20; Buchwald *et al.*, 1980, *Surgery* 88:507; and Saudek *et al.*, 1989, *N. Engl. J. Med.* 321:574). In another embodiment, polymeric materials can be used to achieve controlled release of antibodies (*see e.g.*, Medical Applications of Controlled Release, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); Controlled Drug Bioavailability, Drug Product Design and Performance, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, 1983, *J. Macromol. Sci. Rev. Macromol. Chem.* 23:61; *See also* Levy *et al.*, 1985, *Science* 228:190; During *et al.*, 1989, *Ann. Neurol.* 25:351; Howard *et al.*, 1989, *J. Neurosurg.* 71:105); U.S. Patent No. 5,679,377; U.S. Patent No. 5,916,597; U.S. Patent No. 5,912,015; U.S. Patent No. 5,989,463; U.S. Patent No. 5,128,326; PCT Publication No. WO 99/15154; and PCT Publication No. WO 99/20253). Examples of polymers used in sustained release formulations include, but are not limited to, poly(2-hydroxy ethyl methacrylate), poly(methyl methacrylate), poly(acrylic acid), poly(ethylene-co-vinyl acetate), poly(methacrylic acid), polyglycolides (PLG), polyanhydrides, poly(N-vinyl pyrrolidone), poly(vinyl-alcohol), polyacrylamide, poly(ethylene glycol), polylactides (PLA), poly(lactide-co-glycolides) (PLGA), and polyorthoesters. In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target (*e.g.*, the lungs), thus requiring only a fraction of the systemic dose (*see, e.g.*, Goodson, in *Medical Applications of Controlled Release, supra*, vol. 2, pp. 115-138 (1984)). In another embodiment, polymeric compositions useful as controlled release implants are used according to Dunn *et al.* (*See* U.S. 5,945,155). This particular method is based upon the therapeutic effect of the in situ controlled release of the bioactive material from the polymer system. The implantation can generally occur anywhere within the body of the patient in need of therapeutic treatment. In another embodiment, a non-polymeric sustained delivery system is used, whereby a non-polymeric implant in the body of the subject is used as a drug delivery system. Upon implantation in the body, the organic solvent of the implant will dissipate, disperse, or leach from the composition into surrounding tissue fluid, and the non-polymeric material will gradually coagulate or precipitate to form a solid, microporous matrix (*See* U.S. 5,888,533).

- [00308] Controlled release systems are discussed in the review by Langer (1990, *Science* 249:1527-1533). Any technique known to one of skill in the art can be used to produce sustained release formulations comprising one or more therapeutic agents of the invention. See, e.g., U.S. Patent No. 4,526,938; International Publication Nos. WO 91/05548 and WO 96/20698; Ning *et al.*, 1996, *Radiotherapy & Oncology* 39:179-189; Song *et al.*, 1995, *PDA Journal of Pharmaceutical Science & Technology* 50:372-397; Cleek *et al.*, 1997, *Pro. Int'l. Symp. Control. Rel. Bioact. Mater.* 24:853-854; and Lam *et al.*, 1997, *Proc. Int'l. Symp. Control Rel. Bioact. Mater.* 24:759-760, each of which is incorporated herein by reference in its entirety.
- 10 [00309] In a specific embodiment where the composition of the invention is a nucleic acid encoding an antibody, the nucleic acid can be administered *in vivo* to promote expression of its encoded antibody, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, e.g., by use of a retroviral vector (See U.S. Patent No. 4,980,286), or by direct injection, or by use of
- 15 microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in linkage to a homeobox-like peptide which is known to enter the nucleus (See e.g., Joliot *et al.*, 1991, *Proc. Natl. Acad. Sci. USA* 88:1864-1868), etc. Alternatively, a nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression by
- 20 homologous recombination.
- [00310] For antibodies, the therapeutically or prophylactically effective dosage administered to a subject is typically 0.1 mg/kg to 200 mg/kg of the subject's body weight. Preferably, the dosage administered to a subject is between 0.1 mg/kg and 20 mg/kg of the subject's body weight and more preferably the dosage administered to a subject is between
- 25 1 mg/kg to 10 mg/kg of the subject's body weight. The dosage and frequency of administration of antibodies of the invention may be reduced also by enhancing uptake and tissue penetration (e.g., into the lung) of the antibodies or fusion proteins by modifications such as, for example, lipidation.
- [00311] Treatment of a subject with a therapeutically or prophylactically effective
- 30 amount of molecules of the invention can include a single treatment or, preferably, can include a series of treatments. In a preferred example, a subject is treated with molecules of the invention in the range of between about 0.1 to 30 mg/kg body weight, one time per week for between about 1 to 10 weeks, preferably between 2 to 8 weeks, more preferably between about 3 to 7 weeks, and even more preferably for about 4, 5, or 6 weeks. In other
- 35 embodiments, the pharmaceutical compositions of the invention are administered once a

day, twice a day, or three times a day. In other embodiments, the pharmaceutical compositions are administered once a week, twice a week, once every two weeks, once a month, once every six weeks, once every two months, twice a year or once per year. It will also be appreciated that the effective dosage of the molecules used for treatment may increase or decrease over the course of a particular treatment.

5.6.1 PHARMACEUTICAL COMPOSITIONS

[00312] The compositions of the invention include bulk drug compositions useful in the manufacture of pharmaceutical compositions (*e.g.*, impure or non-sterile compositions) and pharmaceutical compositions (*i.e.*, compositions that are suitable for administration to a subject or patient) which can be used in the preparation of unit dosage forms. Such compositions comprise a prophylactically or therapeutically effective amount of a prophylactic and/or therapeutic agent disclosed herein or a combination of those agents and a pharmaceutically acceptable carrier. Preferably, compositions of the invention comprise a prophylactically or therapeutically effective amount of one or more molecules of the invention and a pharmaceutically acceptable carrier.

[00313] In one particular embodiment, the pharmaceutical composition comprises a therapeutically effective amount of one or more molecules of the invention comprising a variant Fc region, wherein said variant Fc region binds Fc γ RIIA and/or Fc γ RIIA with a greater affinity than a comparable molecule comprising a wild-type Fc region binds Fc γ RIIA and/or Fc γ RIIA and/or said variant Fc region mediates an effector function at least 2-fold more effectively than a comparable molecule comprising a wild-type Fc region, and a pharmaceutically acceptable carrier. In another embodiment, the pharmaceutical composition comprises a therapeutically effective amount of one or more molecules of the invention comprising a variant Fc region, wherein said variant Fc region binds Fc γ RIIA with a greater affinity than a comparable molecule comprising a wild-type Fc region binds Fc γ RIIA, and said variant Fc region binds Fc γ RIIB with a lower affinity than a comparable molecule comprising a wild-type Fc region binds Fc γ RIIB, and/or said variant Fc region mediates an effector function at least 2-fold more effectively than a comparable molecule comprising a wild-type Fc region, and a pharmaceutically acceptable carrier. In another embodiment, said pharmaceutical compositions further comprise one or more anti-cancer agents.

[00314] The invention also encompasses pharmaceutical compositions comprising a therapeutic antibody (*e.g.*, tumor specific monoclonal antibody) that is specific for a particular cancer antigen, comprising one or more amino acid modifications in the Fc region

as determined in accordance with the instant invention, and a pharmaceutically acceptable carrier.

[00315] In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant (e.g., Freund's adjuvant (complete and incomplete), excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like.

[00316] Generally, the ingredients of compositions of the invention are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

[00317] The compositions of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include, but are not limited to those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, *etc.*, and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, *etc.*

5.6.2 GENE THERAPY

[00318] In a specific embodiment, nucleic acids comprising sequences encoding molecules of the invention, are administered to treat, prevent or ameliorate one or more symptoms associated with a disease, disorder, or infection, by way of gene therapy. Gene therapy refers to therapy performed by the administration to a subject of an expressed or expressible nucleic acid. In this embodiment of the invention, the nucleic acids produce their encoded antibody or fusion protein that mediates a therapeutic or prophylactic effect.

[00319] Any of the methods for gene therapy available in the art can be used according to the present invention. Exemplary methods are described below.

10 [00320] For general reviews of the methods of gene therapy, see Goldspiel *et al.*, 1993, *Clinical Pharmacy* 12:488-505; Wu and Wu, 1991, *Biotherapy* 3:87-95; Tolstoshev, 1993, *Ann. Rev. Pharmacol. Toxicol.* 32:573-596; Mulligan, *Science* 260:926-932 (1993); and Morgan and Anderson, 1993, *Ann. Rev. Biochem.* 62:191-217; May, 1993, *TIBTECH* 11(5):155-215. Methods commonly known in the art of recombinant DNA technology
15 which can be used are described in Ausubel *et al.* (eds.), Current Protocols in Molecular Biology, John Wiley & Sons, NY (1993); and Kriegler, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY (1990).

[00321] In a preferred aspect, a composition of the invention comprises nucleic acids encoding an antibody, said nucleic acids being part of an expression vector that expresses
20 the antibody in a suitable host. In particular, such nucleic acids have promoters, preferably heterologous promoters, operably linked to the antibody coding region, said promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, nucleic acid molecules are used in which the antibody coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a
25 desired site in the genome, thus providing for intrachromosomal expression of the antibody encoding nucleic acids (Koller and Smithies, 1989, *Proc. Natl. Acad. Sci. USA* 86:8932-8935; and Zijlstra *et al.*, 1989, *Nature* 342:435-438).

[00322] In another preferred aspect, a composition of the invention comprises nucleic acids encoding a fusion protein, said nucleic acids being a part of an expression vector that
30 expresses the fusion protein in a suitable host. In particular, such nucleic acids have promoters, preferably heterologous promoters, operably linked to the coding region of a fusion protein, said promoter being inducible or constitutive, and optionally, tissue-specific. In another particular embodiment, nucleic acid molecules are used in which the coding sequence of the fusion protein and any other desired sequences are flanked by regions that

promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the fusion protein.

[00323] Delivery of the nucleic acids into a subject may be either direct, in which case the subject is directly exposed to the nucleic acid or nucleic acid-carrying vectors, or
5 indirect, in which case, cells are first transformed with the nucleic acids *in vitro*, then transplanted into the subject. These two approaches are known, respectively, as *in vivo* or *ex vivo* gene therapy.

[00324] In a specific embodiment, the nucleic acid sequences are directly administered *in vivo*, where it is expressed to produce the encoded product. This can be
10 accomplished by any of numerous methods known in the art, *e.g.*, by constructing them as part of an appropriate nucleic acid expression vector and administering it so that they become intracellular, *e.g.*, by infection using defective or attenuated retroviral or other viral vectors (*see* U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (*e.g.*, a gene gun; Biolistic, Dupont), or coating with lipids or
15 cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering them in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (*See, e.g.*, Wu and Wu, 1987, *J. Biol. Chem.* 262:4429-4432) (which can be used to target cell types specifically expressing the receptors), *etc.* In another embodiment, nucleic acid-
20 ligand complexes can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted *in vivo* for cell specific uptake and expression, by targeting a specific receptor (*See, e.g.*, PCT Publications WO 92/06180; WO 92/22635; W092/20316; W093/14188; WO 93/20221). Alternatively, the nucleic acid can
25 be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (Koller and Smithies, 1989, *Proc. Natl. Acad. Sci. USA* 86:8932-8935; and Zijlstra *et al.*, 1989, *Nature* 342:435-438).

[00325] In a specific embodiment, viral vectors that contain nucleic acid sequences encoding a molecule of the invention (*e.g.*, an antibody or a fusion protein) are used. For
30 example, a retroviral vector can be used (*See* Miller *et al.*, 1993, *Meth. Enzymol.* 217:581-599). These retroviral vectors contain the components necessary for the correct packaging of the viral genome and integration into the host cell DNA. The nucleic acid sequences encoding the antibody or a fusion protein to be used in gene therapy are cloned into one or more vectors, which facilitates delivery of the nucleotide sequence into a subject. More
35 detail about retroviral vectors can be found in Boesen *et al.*, (1994, *Biotherapy* 6:291-302),

which describes the use of a retroviral vector to deliver the *mdr 1* gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes *et al.*, 1994, *J. Clin. Invest.* 93:644-651; Klein *et al.*, 1994, *Blood* 83:1467-1473; Salmons and Gunzberg, 1993, *Human Gene Therapy* 4:129-141; and Grossman and Wilson, 1993, *Curr. Opin. in Genetics and Devel.* 3:110-114.

[00326] Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson (*Current Opinion in Genetics and Development* 3:499-503, 1993, present a review of adenovirus-based gene therapy. Bout *et al.*, (*Human Gene Therapy*, 5:3-10, 1994) demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld *et al.*, 1991, *Science* 252:431-434; Rosenfeld *et al.*, 1992, *Cell* 68:143-155; Mastrangeli *et al.*, 1993, *J. Clin. Invest.* 91:225-234; PCT Publication W094/12649; and Wang *et al.*, 1995, *Gene Therapy* 2:775-783. In a preferred embodiment, adenovirus vectors are used.

[00327] Adeno-associated virus (AAV) has also been proposed for use in gene therapy (*see, e.g.*, Walsh *et al.*, 1993, *Proc. Soc. Exp. Biol. Med.* 204:289-300 and U.S. Patent No. 5,436,146).

[00328] Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a subject.

[00329] In this embodiment, the nucleic acid is introduced into a cell prior to administration *in vivo* of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to, transfection, electroporation, microinjection, infection with a viral or bacteriophage vector, containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcellmediated gene transfer, spheroplast fusion, *etc.* Numerous techniques are known in the art for the introduction of foreign genes into cells (*See, e.g.*, Loeffler and Behr, 1993,

Meth. Enzymol. 217:599-618, Cohen *et al.*, 1993, *Meth. Enzymol.* 217:618-644; and *Clin. Pharma. Ther.* 29:69-92, 1985) and may be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to
5 the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

[00330] The resulting recombinant cells can be delivered to a subject by various methods known in the art. Recombinant blood cells (*e.g.*, hematopoietic stem or progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use
10 depends on the desired effect, patient state, *etc.*, and can be determined by one skilled in the art.

[00331] Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood
15 cells such as T lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, *e.g.*, as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, *etc.*

[00332] In a preferred embodiment, the cell used for gene therapy is autologous to
20 the subject.

[00333] In an embodiment in which recombinant cells are used in gene therapy, nucleic acid sequences encoding an antibody or a fusion protein are introduced into the cells such that they are expressible by the cells or their progeny, and the recombinant cells are then administered *in vivo* for therapeutic effect. In a specific embodiment, stem or
25 progenitor cells are used. Any stem and/or progenitor cells which can be isolated and maintained *in vitro* can potentially be used in accordance with this embodiment of the present invention (*See e.g.*, PCT Publication WO 94/08598; Stemple and Anderson, 1992, *Cell* 71:973-985; Rheinwald, 1980, *Meth. Cell Bio.* 21A:229; and Pittelkow and Scott, 1986, *Mayo Clinic Proc.* 61:771).

30 [00334] In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy comprises an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription.

5.6.3 KITS

[00335] The invention provides a pharmaceutical pack or kit comprising one or more containers filled with the molecules of the invention (*i.e.*, antibodies, polypeptides comprising variant Fc regions). Additionally, one or more other prophylactic or therapeutic agents useful for the treatment of a disease can also be included in the pharmaceutical pack or kit. The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

[00336] The present invention provides kits that can be used in the above methods. In one embodiment, a kit comprises one or more molecules of the invention. In another embodiment, a kit further comprises one or more other prophylactic or therapeutic agents useful for the treatment of cancer, in one or more containers. In another embodiment, a kit further comprises one or more cytotoxic antibodies that bind one or more cancer antigens associated with cancer. In certain embodiments, the other prophylactic or therapeutic agent is a chemotherapeutic. In other embodiments, the prophylactic or therapeutic agent is a biological or hormonal therapeutic.

20 5.7 CHARACTERIZATION AND DEMONSTRATION OF THERAPEUTIC UTILITY

[00337] Several aspects of the pharmaceutical compositions, prophylactic, or therapeutic agents of the invention are preferably tested *in vitro*, in a cell culture system, and in an animal model organism, such as a rodent animal model system, for the desired therapeutic activity prior to use in humans. For example, assays which can be used to determine whether administration of a specific pharmaceutical composition is desired, include cell culture assays in which a patient tissue sample is grown in culture, and exposed to or otherwise contacted with a pharmaceutical composition of the invention, and the effect of such composition upon the tissue sample is observed. The tissue sample can be obtained by biopsy from the patient. This test allows the identification of the therapeutically most effective prophylactic or therapeutic molecule(s) for each individual patient. In various specific embodiments, *in vitro* assays can be carried out with representative cells of cell types involved in an autoimmune or inflammatory disorder (*e.g.*, T cells), to determine if a pharmaceutical composition of the invention has a desired effect upon such cell types.

[00338] Combinations of prophylactic and/or therapeutic agents can be tested in suitable animal model systems prior to use in humans. Such animal model systems include, but are not limited to, rats, mice, chicken, cows, monkeys, pigs, dogs, rabbits, *etc.* Any animal system well-known in the art may be used. In a specific embodiment of the invention, combinations of prophylactic and/or therapeutic agents are tested in a mouse model system. Such model systems are widely used and well-known to the skilled artisan. Prophylactic and/or therapeutic agents can be administered repeatedly. Several aspects of the procedure may vary. Said aspects include the temporal regime of administering the prophylactic and/or therapeutic agents, and whether such agents are administered separately or as an admixture.

[00339] Preferred animal models for use in the methods of the invention are, for example, transgenic mice expressing human Fc γ R on mouse effector cells, *e.g.*, any mouse model described in U.S. 5,877,396 (which is incorporated herein by reference in its entirety) can be used in the present invention. Transgenic mice for use in the methods of the invention include, but are not limited to, mice carrying human Fc γ RIIIA; mice carrying human Fc γ RIIA; mice carrying human Fc γ RIIB and human Fc γ RIIIA; mice carrying human Fc γ RIIB and human Fc γ RIIA.

[00340] Preferably, mutations showing the highest levels of activity in the functional assays described above will be tested for use in animal model studies prior to use in humans. Antibodies harboring the Fc mutants identified using the methods of the invention and tested in ADCC assays, including ch4D5 and ch520C9, two anti-Erb-B2 antibodies, and chCC49, an anti-TAG72 antibody, are preferred for use in animal models since they have been used previously in xenograft mouse model (Hudsiak *et al.*, 1989, *Mol. Cell Biol.* 9: 1165-72; Lewis *et al.*, 1993, *Cancer Immunol. Immunother.* 37: 255-63; Bergman *et al.*, 2001 *Clin. Cancer Res.* 7: 2050-6; Johnson *et al.*, 1995, *Anticancer Res.* 1387-93). Sufficient quantities of antibodies may be prepared for use in animal models using methods described supra, for example using mammalian expression systems and IgG purification methods disclosed and exemplified herein. A typical experiment requires at least about 5.4 mg of mutant antibody. This calculation is based on average quantities of wild type antibody required to protect 8-10 30 g mice following a loading dose of 4 μ g/g and a weekly maintenance dose, 2 μ g/g, for ten weeks. invention encompasses tumor cell lines as a source for xenograft tumors, such as SK-BR-3, BT474 and HT29 cells which are derived from patients with breast adenocarcinoma. These cells have both Erb-B2 and the prolactin receptors on their surface. The SK-BR-3 cells have been used successfully in both ADCC and xenograft tumor models. In other assays OVCAR3 cells derived from a human ovarian

adenocarcinoma may be used. These cells express the antigen TAG72 on the cell surface and can be used in conjunction with the chCC49 antibody. The use of different antibodies and multiple tumor models will circumvent loss of any specific mutations due to an antibody specific Fc mutant incompatibility.

- 5 [00341] Mouse xenograft models may be used for examining efficacy of mouse antibodies generated against a tumor specific target based on the affinity and specificity of the CDR regions of the antibody molecule and the ability of the Fc region of the antibody to elicit an immune response (Wu et al., 2001, Trends Cell Biol. 11: S2-9). Transgenic mice expressing human FcγRs on mouse effector cells are unique and are tailor-made animal
- 10 models to test the efficacy of human Fc-FcγR interactions. Pairs of FcγRIIIA, FcγRIIIB and FcγRIIA transgenic mouse lines generated in the lab of Dr. Jeffrey Ravetch (Through a licensing agreement with Rockefeller U. and Sloan Kettering Cancer center) can be used such as those listed in the Table below.

Table 5: Mice Strains

Strain Background	Human FcR
Nude / CD16A KO	none
Nude / CD16A KO	FcγRIIIA
Nude / CD16A KO	FcγR IIA
Nude / CD16A KO	FcγR IIA and IIIA
Nude / CD32B KO	none
Nude / CD32B KO	FcγR IIB

- 15 [00342] Preferably Fc mutants showing both enhanced binding to FcγRIIIA and reduced binding to FcγRIIB, increased activity in ADCC and phagocytosis assays are tested in animal model experiments. The animal model experiments examine the increase in efficacy of Fc mutant bearing antibodies in FcγRIIIA transgenic, nude mCD16A knockout mice compared to a control which has been administered native antibody. Preferably,
- 20 groups of 8-10 mice are examined using a standard protocol. An exemplary animal model experiment may comprise the following steps: in a breast cancer model, $\sim 2 \times 10^6$ SK-BR-3 cells are injected subcutaneously on day 1 with 0.1 mL PBS mixed with Matrigel (Becton Dickinson). Initially a wild type chimeric antibody and isotype control are administered to establish a curve for the predetermined therapeutic dose, intravenous injection of 4D5 on
- 25 day 1 with an initial dose of 4 μg/g followed by weekly injections of 2 μg/g. Tumor volume is monitored for 6-8 weeks to measure progress of the disease. Tumor volume should increase linearly with time in animals injected with the isotype control. In contrast very little tumor growth should occur in the group injected with 4D5. Results from the standard

dose study are used to set an upper limit for experiments testing the Fc mutants. These studies are done using subtherapeutic doses of the Fc mutant containing antibodies. A one tenth dose was used on xenograft models in experiments done in FcγRIIB knockout mice, see, Clynes *et al.*, 2000, *Nat. Med.* 6: 443-6, with a resultant block in tumor cell growth.

- 5 Since the mutants of the invention preferably show an increase in FcγRIIA activation and reduction in FcγRIIB binding the mutants are examined at one tenth therapeutic dose. Examination of tumor size at different intervals indicates the efficacy of the antibodies at the lower dose. Statistical analysis of the data using *t* test provides a way of determining if the data is significant. Fc mutants that show increased efficacy are tested at incrementally lower doses to determine the smallest possible dose as a measure of their efficacy.

- 10 [00343] The anti-inflammatory activity of the combination therapies of invention can be determined by using various experimental animal models of inflammatory arthritis known in the art and described in Crofford L.J. and Wilder R.L., "Arthritis and Autoimmunity in Animals", in *Arthritis and Allied Conditions: A Textbook of Rheumatology*, McCarty *et al.* (eds.), Chapter 30 (Lee and Febiger, 1993). Experimental and spontaneous animal models of inflammatory arthritis and autoimmune rheumatic diseases can also be used to assess the anti-inflammatory activity of the combination therapies of invention. The following are some assays provided as examples, and not by limitation.

- 20 [00344] The principle animal models for arthritis or inflammatory disease known in the art and widely used include: adjuvant-induced arthritis rat models, collagen-induced arthritis rat and mouse models and antigen-induced arthritis rat, rabbit and hamster models, all described in Crofford L.J. and Wilder R.L., "Arthritis and Autoimmunity in Animals", in *Arthritis and Allied Conditions: A Textbook of Rheumatology*, McCarty *et al.* (eds.), Chapter 30 (Lee and Febiger, 1993), incorporated herein by reference in its entirety.

- 25 [00345] The anti-inflammatory activity of the combination therapies of invention can be assessed using a carrageenan-induced arthritis rat model. Carrageenan-induced arthritis has also been used in rabbit, dog and pig in studies of chronic arthritis or inflammation. Quantitative histomorphometric assessment is used to determine therapeutic efficacy. The methods for using such a carrageenan-induced arthritis model is described in Hansra P. *et al.*, "Carrageenan-Induced Arthritis in the Rat," *Inflammation*, 24(2): 141-155, (2000). Also commonly used are zymosan-induced inflammation animal models as known and described in the art.

- 30 [00346] The anti-inflammatory activity of the combination therapies of invention can also be assessed by measuring the inhibition of carrageenan-induced paw edema in the rat,

using a modification of the method described in Winter C. A. *et al.*, "Carrageenan-Induced Edema in Hind Paw of the Rat as an Assay for Anti-inflammatory Drugs" *Proc. Soc. Exp. Biol Med.* 111, 544-547, (1962). This assay has been used as a primary *in vivo* screen for the anti-inflammatory activity of most NSAIDs, and is considered predictive of human efficacy.

- 5 The anti-inflammatory activity of the test prophylactic or therapeutic agents is expressed as the percent inhibition of the increase in hind paw weight of the test group relative to the vehicle dosed control group.

[00347] Additionally, animal models for inflammatory bowel disease can also be used to assess the efficacy of the combination therapies of invention (Kim *et al.*, 1992, *Scand. J. Gastroenterol.* 27:529-537; Strober, 1985, *Dig. Dis. Sci.* 30(12 Suppl):3S-10S).
10 Ulcerative colitis and Crohn's disease are human inflammatory bowel diseases that can be induced in animals. Sulfated polysaccharides including, but not limited to amylopectin, carrageen, amylopectin sulfate, and dextran sulfate or chemical irritants including but not limited to trinitrobenzenesulphonic acid (TNBS) and acetic acid can be administered to
15 animals orally to induce inflammatory bowel diseases.

[00348] Animal models for autoimmune disorders can also be used to assess the efficacy of the combination therapies of invention. Animal models for autoimmune disorders such as type 1 diabetes, thyroid autoimmunity, sytemic lupus erthematosus, and glomerulonephritis have been developed (Flanders *et al.*, 1999, *Autoimmunity* 29:235-246;
20 Krogh *et al.*, 1999, *Biochimie* 81:511-515; Foster, 1999, *Semin. Nephrol.* 19:12-24).

[00349] Further, any assays known to those skilled in the art can be used to evaluate the prophylactic and/or therapeutic utility of the combinatorial therapies disclosed herein for autoimmune and/or inflammatory diseases.

[00350] Toxicity and efficacy of the prophylactic and/or therapeutic protocols of the
25 instant invention can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD₅₀/ED₅₀. Prophylactic and/or therapeutic agents that exhibit large
30 therapeutic indices are preferred. While prophylactic and/or therapeutic agents that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such agents to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

[00351] The data obtained from the cell culture assays and animal studies can be used
35 in formulating a range of dosage of the prophylactic and/or therapeutic agents for use in

humans. The dosage of such agents lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any agent used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC₅₀ (i.e., the concentration of the test compound that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[00352] The anti-cancer activity of the therapies used in accordance with the present invention also can be determined by using various experimental animal models for the study of cancer such as the SCID mouse model or transgenic mice or nude mice with human xenografts, animal models, such as hamsters, rabbits, etc. known in the art and described in *Relevance of Tumor Models for Anticancer Drug Development* (1999, eds. Fiebig and Burger); *Contributions to Oncology* (1999, Karger); *The Nude Mouse in Oncology Research* (1991, eds. Boven and Winograd); and *Anticancer Drug Development Guide* (1997 ed. Teicher), herein incorporated by reference in their entireties.

[00353] Preferred animal models for determining the therapeutic efficacy of the molecules of the invention are mouse xenograft models. Tumor cell lines that can be used as a source for xenograft tumors include but are not limited to, SKBR3 and MCF7 cells, which can be derived from patients with breast adenocarcinoma. These cells have both erbB2 and prolactin receptors. SKBR3 cells have been used routinely in the art as ADCC and xenograft tumor models. Alternatively, OVCAR3 cells derived from a human ovarian adenocarcinoma can be used as a source for xenograft tumors.

[00354] The protocols and compositions of the invention are preferably tested *in vitro*, and then *in vivo*, for the desired therapeutic or prophylactic activity, prior to use in humans. Therapeutic agents and methods may be screened using cells of a tumor or malignant cell line. Many assays standard in the art can be used to assess such survival and/or growth; for example, cell proliferation can be assayed by measuring ³H-thymidine incorporation, by direct cell count, by detecting changes in transcriptional activity of known genes such as proto-oncogenes (e.g., fos, myc) or cell cycle markers; cell viability can be assessed by trypan blue staining, differentiation can be assessed visually based on changes in morphology, decreased growth and/or colony formation in soft agar or tubular network formation in three-dimensional basement membrane or extracellular matrix preparation, etc.

[00355] Compounds for use in therapy can be tested in suitable animal model systems prior to testing in humans, including but not limited to in rats, mice, chicken, cows, monkeys, rabbits, hamsters, *etc.*, for example, the animal models described above. The compounds can then be used in the appropriate clinical trials.

- 5 [00356] Further, any assays known to those skilled in the art can be used to evaluate the prophylactic and/or therapeutic utility of the combinatorial therapies disclosed herein for treatment or prevention of cancer, inflammatory disorder, or autoimmune disease.

6. EXAMPLES

[00357] Using a yeast display system, mutant human IgG1 heavy chain Fc regions were screened for modified affinity to different Fc receptors. In particular, a mutant Fc library was generated by error prone PCR (Genemorph, Stratagene), and then the mutant Fc proteins were fused to the Aga2p cell wall protein, which allowed the fusion protein to be secreted extracellularly and displayed on the yeast cell wall.

[00358] Soluble forms of the human receptors (Fc γ RIIA and Fc γ RIIB) were cloned. Detection of the IgG1 Fc domains on the yeast cell surface, however, is hindered due to the low affinity of Fc γ R for its ligand. In order to circumvent this limitation, soluble Fc γ R tetrameric complexes were formed using an AVITAG sequence which could be enzymatically biotinylated and subsequently reacted with streptavidin conjugated to phycoerythrin (SA-PE; Molecular Probes) to form soluble tetrameric Fc γ R complexes. ELISA assays confirmed that the soluble Fc γ R tetrameric complexes had a higher avidity for human IgG1 relative to the monomeric Fc γ R. Fc fusion proteins on the yeast cell surface also bound the soluble Fc γ R tetrameric complexes as assessed by FACS analysis.

[00359] The differential binding of the Fc fusion proteins expressed on the yeast cell surface to soluble tetrameric Fc γ R complexes was monitored by a FACS analysis. Fc fusion proteins with altered affinities for one or more soluble tetrameric Fc γ R complexes were thus identified and were then incorporated into a complete immunoglobulin and expressed in mammalian cells. The mammalian expressed product was used in ELISA assays to confirm the results obtained in the yeast surface display system. Finally, the mutant Fc regions were sequenced to confirm the altered residue(s).

6.1 CLONING, EXPRESSION AND PURIFICATION OF Fc γ RIIA

25 MATERIALS AND METHODS

[00360] Soluble Fc γ RIIB and Fc γ RIIA were cloned as follows. The cDNA clones for the human Fc γ R genes (Fc γ RIIB and Fc γ RIIA) were obtained (gift from Ravetch lab). Soluble region of the Fc γ RIIA gene (amino acids 7- 203) was amplified by PCR (Table 5), digested with BamHI/HindIII and ligated into the pET25vector (Novagen). This vector was digested with Sall/NotI and a 370 by fragment was gel isolated. The vector hu3A, (gift from J. Ravetch) was digested with BamHI/Sall and a 270 by fragment containing the N-terminus of Fc γ RIIA was isolated. Both fragments were coligated into pcDNA3.1 cut with BamH/NotI to create pcDNA3-Fc γ RIIA (amino acids 1- 203). The soluble region of

Fc γ RIIB (amino acids 33 - 180) was amplified by PCR (Table 5), digested with BglII/HindIII and ligated into pET25b(+) (Novagen). This vector was digested with BamHI/NotI and a 140 bp fragment was gel isolated. The vector huRIIb1 (gift from J. Ravetch) was digested with BamHI/EcoRI and a 440 bp N-terminal Fc γ RIIB fragment was isolated. Both of these fragments were coligated into pcDNA3.1 cut with BamHI/NotI to create pcDNA3-Fc γ RIIB (amino acids 1- 180). Recombinant clones were transfected into 293H cells, supernatants were collected from cell cultures, and soluble recombinant Fc γ R (rFc γ R) proteins were purified on an IgG sepharose column.

RESULTS

Recombinant soluble Fc γ RIIA (rFc γ RIIA) and recombinant soluble Fc γ RIIB (rFc γ RIIB) were purified to homogeneity

[00361] Subsequent to expression and purification of the recombinant soluble Fc γ R proteins on an IgG sepharose column, the purity and apparent molecular weight of the recombinant purified soluble receptor proteins were determined by SDS-PAGE. As shown in FIG. 1, soluble rFc γ RIIA (FIG. 1, lane 1) had the expected apparent molecular weight of ~35KDa and soluble rFc γ RIIB (FIG. 1, lane 4) had the expected apparent molecular weight of ~20KDa. As shown in FIG. 1, soluble rFc γ RIIA migrates as a diffuse "fuzzy" band which has been attributed to the high degree of glycosylation normally found on Fc γ RIIA (Jefferis, *et al.*, 1995 *Immunol Lett.* 44, 111-117).

6.1.1 CHARACTERIZATION OF PURIFIED RECOMBINANT SOLUBLE Fc γ RIIA

MATERIALS AND METHODS

[00362] Purified soluble rFc γ RIIA, which was obtained as described above, was analyzed for direct binding against human monomeric or aggregated IgG using an ELISA assay. The plate is coated with 10ng of soluble rFc γ RIIA overnight in 1X PBS. Subsequent to coating, the plate is washed three times in 1X PBS/0.1% Tween 20. Human IgG, either biotinylated monomeric IgG or biotinylated aggregated IgG, is added to the wells at a concentration ranging from 0.03 mg/mL to 2 mg/mL, and allowed to bind to the soluble rFc γ RIIA. The reaction is carried out for one hour at 37°C. The plate is washed again three times with 1X PBS/0.1% Tween 20. The binding of human IgG to soluble rFc γ RIIA is detected with streptavidin horseradish peroxidase conjugate by monitoring the absorbance at 650nm. The absorbance at 650nm is proportional to the bound aggregated IgG.

[00363] In a blocking ELISA experiment, the ability of an Fc γ RIIIA monoclonal antibody, 3G8, a mouse anti-Fc γ RIIIA antibody (Pharmingen), to block the binding of the receptor to aggregated IgG is monitored. The washing and incubation conditions were the same as described above, except that prior to IgG addition, a 5-fold molar excess of 3G8 was added and allowed to incubate for 30 minutes at 37°C.

RESULTS

[00364] Purified, recombinant soluble Fc γ RIIIA binds aggregated IgG specifically

[00365] The direct binding of purified recombinant soluble Fc γ RIIIA to aggregated and monomeric IgG was tested using an ELISA assay (FIG. 2). At an IgG concentration of 2 μ g/ml, strong binding to the aggregated IgG was observed. However, at a similar concentration, no binding was detected to the monomeric IgG. The binding to aggregated IgG was blocked by 3G8, a mouse anti-Fc γ RIIIA monoclonal antibody that blocks the ligand binding site, indicating that the aggregated IgG binding is via that of the normal Fc γ RIIIA ligand binding site (FIG. 2). Soluble rFc γ RIIB was also characterized and shown to bind to IgG with similar characteristics as the soluble rFc γ RIIIA (data not shown).

6.2 FORMATION OF SOLUBLE Fc γ R TETRAMERIC COMPLEXES

MATERIALS AND METHODS

Construction of plasmids for expression of soluble FcR γ IIIA and FcR γ IIB fused to the AVITAG peptide.

[00366] To generate soluble Fc γ R tetrameric complexes, the soluble region of the human FcR γ IIIA gene (amino acids 7-203) was amplified by PCR (Table 1), digested with BamHI/HindIII and ligated into the pET25b(+) (Novagen). This vector was digested with SalI/NotI, and a 370 bp fragment was isolated by agarose gel electrophoresis. The vector hu3A, (gift from J. Ravetch) was digested with BamHI/SalI, and a 270 bp fragment containing the N-terminus of FcR γ IIIA was isolated. Both fragments were coligated into pcDNA3.1 (Invitrogen), which had been digested with BamH/NotI to create pcDNA3-FcR γ IIIA (amino acids 1-203).

[00367] The soluble region of FcR γ IIB (amino acids 33 - 180) was amplified by PCR (Table I), digested with BglII/HindIII and ligated into pET25b(+) (Novagen). This vector was digested with BamHI/NotI, and a 140 bp fragment was isolated by agarose gel electrophoresis. The vector huRIIB₁ (gift from J. Ravetch) was digested with BamHI/EcoRI, and a 440 bp FcR γ IIB N-terminal fragment was isolated. Both of these fragments were co-ligated into pcDNA3.1, which had been digested with BamHI/NotI to

create pcDNA3-FcR γ IIB (amino acids 1-180). Subsequently, the linker-AVITAG sequence was fused to the C-terminus of both FcR γ IIIA and FcR γ IIB. To generate the FcR γ IIIA-linker-avitag and FcR γ IIB-linker-avitag constructs, the pcDNA3.1 FcR γ IIIA and FcR γ IIB constructs were digested with Not I and XbaI (both cut in the vector sequence) and a 86 base pair double stranded oligonucleotide consisting of NotI site at the 5' end and XbaI at the 3' end was ligated into the vector. This 86 bp fragment was generated by annealing two 5' phosphorylated reverse complement oligonucleotides (shown in Table 8 as 5' and 3' linker.avitag primers) with the restrictions sites for NotI and XbaI already pre-designed. Equal volumes of each primer at 100 ng per ul were mixed and the DNA heated to 90 °C for 15 minutes and cooled at room temperature for an hour to anneal. This created a double-stranded DNA fragment ready to be ligated to the pcDNA3.1-FcR γ IIIA and FcR γ IIB constructs digested with the respective enzymes. Therefore, the *pcDNA3.1-FcR γ IIIA-linker-AVITAG* and *pcDNA3.1-FcR γ IIB-linker-AVITAG*, were constructed.

TABLE 8: PRIMERS USED FOR CONSTRUCTION OF Fc γ R AND IgG VECTORS

Oligomer	Sequence
5' linker.avitag (SEQ. ID NO. 1)	GGCCGCAGGTGGTGGTGGTTCTGGTGGTGGTGGTTCTGGTCTGAACGACATCTT CGAGGCTCAGAAAA TCGAATGGCACGAATGAT
3' linker.avitag (SEQ. ID NO. 2)	CTAGATCATTCGTGCCATTTCGATTTTCTGAGCCTCGAAGATGTCGTTTCAGACCAG AACCACCAACCACCAAGAACCACCAACCACCTGC
FcR γ IIIA left (SEQ. ID NO. 3)	G TTG GAT CCT CCA ACT GCT CTG CTA CTT CTA GTT T
FcR γ IIIA right (SEQ. ID NO. 4)	GAA AAG CTT AAA GAA TGA TGA GAT GGT TGA CAC T
FcR γ IIBright (SEQ. ID NO. 5)	GAA GTC GAC AAT GAT CCC CAT TGG TGA AGA G
FcR γ IIBleft (SEQ. ID NO. 6)	G TTA GAT CTT GCT GTG CTA TTC CTG GCT CC
IgG1 right (SEQ. ID NO. 7)	ATA GTC GAC CAC TGA TTT ACC CGG AGA
IgG1left (SEQ. ID NO. 8)	GGAA TTC AAC ACC AAG GTG GAC AAG AAA GTT
mcr025;chl (f') (SEQ. ID NO. 9)	AAA GGATCC GCG AGC TCA GCC TCC ACC AAG G

H021 (SEQ. ID NO. 10)	GTCTGCTCGAAGCATTAACC
-----------------------------	----------------------

Biotinylation by BirA

- [00368] Soluble Fc receptors (Fc γ R) fused to the 15 amino acid AVITAG sequence (Avidity, CO) (Schatz P.J., 1993, *Biotechnology*, 11:1138-1143) at the C-terminus of the protein cloned into pcDNA3.1 were generated by transiently transfecting 293H cells using Lipofectamine 2000 reagent (Invitrogen, CA). Supernatants were collected from the cultures and soluble FcR proteins were purified by passing the supernatants over an IgG sepharose column. Concentration of the soluble FcR-AVITAG fusion protein was quantitated by absorbance at 280 nm. The AVITAG present on the soluble FcR proteins was biotinylated according to the manufacturer's protocol (Avidity, CO) with the *E. coli* BirA enzyme, a biotin ligase. A 1:100 final dilution of a cocktail of protease inhibitors (Sigma catalog #P8849) and 1 mg/ml final concentration of Leupeptin (Sigma L-8511) were added to the mixture to prevent degradation of the proteins. The BirA reaction was incubated at room temperature overnight, following which the solution was concentrated using a Biomax 10K-ultrafiltration device (Millipore) by centrifugation at 3500 rpm at 4°C. The protein was loaded onto an FPLC Superdex 200 HR 10/30 column (Pharmacia Biotech) in Tris-HCl (20 mM, pH 8.0), 50 mM NaCl to separate the labeled soluble Fc γ R from free biotin.

Determination of the extent of biotinylation by streptavidin shift assay

- [00369] Approximately 80-85% of the protein was biotinylated by the BirA enzyme (Avidity, CO). The streptavidin-shift assay was used to determine the extent of biotinylation of the protein. Biotinylated protein was incubated with streptavidin (MW 60,000 Daltons) in different ratios. Unbiotinylated protein alone and streptavidin alone are included as controls to determine the extent of biotinylation. The incubation is carried out either on ice for 2 hours or overnight at 4°C. Samples are analyzed on a 4-12% SDS-PAGE Bis-Tris (Invitrogen, CA) with reducing agent and without boiling of the samples. Streptavidin bound biotinylated protein migrates as a high molecular weight band. The extent of biotinylation is estimated by the amount of monomeric protein left in the sample. Absence of monomeric low molecular weight species and presence of a complex with molecular weight greater than streptavidin alone indicates a high degree of biotinylation.

Formation of FcγR tetrameric complexes

[00370] Formation of FcγR tetrameric complexes was performed according to previously established methodologies for MHC class I tetramers (*See* Busch, D. H. et al., 1998 *Immunity* 8:353-362; Altman, J. D. et al., 1996, *Science* 274: 94-96). The

5 concentration of the biotinylated monomeric FcγR was calculated based on absorbance at 280 nm. One molecule of streptavidin-phycoerythrin (SA-PE) (Molecular Probes, OR) has the capacity to bind 4 molecules of biotin. A 5:1 molar ratio of monomeric biotinylated FcγR to SA-PE (5X monomeric biotinylated FcγR: 1X SA-PE) was used to ensure an excess of biotinylated protein. The calculated molecular weight of SA-PE is 300,000

10 Daltons, therefore 303 mL of a 1 mg/mL solution of streptavidin-PE has 1 nmole of SA-PE, which was added to 5 nmole of protein. Efficient formation of tetrameric protein requires SA-PE to be added in step-wise increments. Half the amount of SA-PE was added upfront, and the remaining SA-PE was added in small aliquots every 20-30 minutes at 4°C in the dark. The intervals for the addition of the remaining SA-PE is flexible. After the addition

15 of SA-PE was complete, the solution was concentrated and loaded over an FPLC size exclusion column as above in phosphate buffered saline, at pH 7.4. The fraction that eluted in the void volume with a molecular weight greater than SA-PE alone was collected. Protease inhibitors were replenished to prevent protein degradation. The solution was concentrated and additional protease inhibitors were added to the final complex for storage.

20 The final concentration of the soluble FcγR tetrameric complex was calculated based on the starting concentration of the biotinylated monomeric protein. For example, if 500 μg of biotinylated protein was used to make the tetrameric complex and the final concentrated tetramers were in a volume of 500 μL, the concentration is estimated to be approximately 1 mg/mL (The losses incurred during concentration are not taken into account as it is difficult

25 to accurately determine how much is lost during each step of the formation of the tetramers. It is also not possible to take an absorbance at 280 nm to measure the concentration due to interference from the PE). Soluble FcγR tetrameric complexes were dispensed in small aliquots at -80°C for long term storage with protease inhibitors. Sodium azide was not added to these preparations as the tetramers were used for screening a yeast display library.

30 On thawing an aliquot, the tetramers were stored at 4°C for up to 1 week.

ELISA assay for Characterizing the Tetrameric FcγR Complexes

[00371] An ELISA was used to characterize the tetrameric FcγR complexes. Maxisorb F96 well plate (Nunc) was coated with 25 ng of human IgG in PBS buffer, and incubated overnight at 4 °C. The plates were washed with PBS/0.5% BSA/0.1% Tween 20

- (wash and diluent buffer) before adding the combination of Fc γ RIIIA tetramers and test antibodies to determine blocking with 3G8, a mouse anti-human Fc γ RIIIA antibody as described below: The blocking step was performed as follows: soluble Fc γ RIIIA tetramers at a fixed 0.5 mg/ml final concentration were pre-incubated with antibodies for 1 h at room temperature in buffer, PBS/0.5% BSA/0.1% Tween 20. The final concentrations of the antibodies ranged from 60 mg/mL to 0.25 mg/mL. 3G8 is a mouse anti-human Fc γ RIIIA antibody, and for the purpose of this experiment, a chimeric version was used, *i.e.*, the variable region of the antibody is a mouse anti-human Fc γ RIIIA and the constant region of the heavy and light chains is from the IgG1 human region. A chimeric 4.4.20. D265A was also used in this experiment, which is an anti-fluorescein antibody, such that the Fc region contains a mutation at position 265, where an aspartic acid is substituted with alanine in the human IgG1, which results in a reduced binding to Fc γ R. This antibody has been characterized previously (*See Clynes et al.*, 2000, *Nat. Med.* 6: 443-446; *Shields et al.*, 2001, *J. Biol. Chem.*, 276: 6591-6604). This antibody was used as negative isotype control.
- 15 [00372] The antibodies were allowed to bind to Fc γ RIIIA tetramers, by pre-incubation for 1 hour at room temperature. The mixture was then added to the IgG on the washed plate and incubated for an additional hour at room temperature. The plate was washed with buffer and DJ130c (a mouse anti-human Fc γ RIIIA antibody available from DAKO, Denmark; its epitope is distinct from that of the 3G8 antibody) at 1:5000 dilution was added and allowed to incubate for 1 hr. at room temperature in order to detect the bound Fc γ RIIIA tetramers. Unbound antibodies were washed out with buffer and the bound DJ130c was detected with goat anti-mouse peroxidase (Jackson laboratories). This reagent will not detect the human Fc. After washing out the unbound peroxidase-conjugated antibody, the substrate, TMB reagent (BioF_x), was added to detect the extent of blocking with 3G8 versus the isotype control and the developed color was read at 650 nm.
- 25 [00373] For direct binding of soluble tetrameric Fc γ RIIIA to IgG by ELISA, maxisorb plates were coated with 25 ng IgG as described above. The soluble tetrameric Fc γ RIIIA were added from 20 mg/mL to 0.1 mg/mL and the biotinylated monomeric soluble tetrameric Fc γ RIIIA were added at concentrations ranging from 20 mg/mL to 0.16 mg/mL. Detection was the same as above with DJ130c, followed by goat anti-mouse-peroxidase antibody. Color developed with the TMB reagent and the plate was read at 650 nm.
- 30

RESULTS

Soluble Fc γ RIIIA tetrameric complex binds monomeric human IgG via its normal ligand binding site

[00374] Soluble Fc γ RIIIA-AVITAG fusion proteins were generated, isolated, and
5 analyzed as described in the Material and Methods section using an ELISA assay and were shown to have similar properties as the non-AVITAG soluble Fc γ RIIIA protein (data not shown). The fusion proteins were biotinylated, and the tetrameric complexes were generated as described above.

[00375] The soluble Fc γ R tetrameric complex was then assessed for binding its
10 ligand, monomeric human IgG, using an ELISA assay. Analysis by ELISA showed the soluble tetrameric Fc γ R complexes bind monomeric human IgG specifically. As shown in FIG. 3A, binding of soluble tetrameric Fc γ RIIIA to monomeric human IgG is blocked by 3G8, a mouse anti-human Fc γ RIIA monoclonal antibody, as monitored by the absorbance at 650nm. On the other hand, the 4-4-20 monoclonal antibody harboring the D265A mutation
15 was not able to block the binding of soluble tetrameric Fc γ RIIIA to monomeric human IgG (FIG. 3A). This experiment thus confirms that binding of the soluble tetrameric Fc γ RIIIA complex occurs through the native ligand binding site.

Soluble Fc γ RIIIA tetrameric complex binds monomeric human IgG with a greater avidity than monomeric soluble Fc γ RIIA

20 [00376] The direct binding of soluble tetrameric Fc γ RIIIA to aggregated human IgG was assessed using an ELISA assay and compared to the direct binding of soluble monomeric Fc γ RIIIA to monomeric human IgG. As shown in FIG. 3B, soluble tetrameric Fc γ RIIIA binds human IgG with a higher avidity (8-10 fold) than the soluble monomeric receptor, as monitored by the absorbance at 450 nm.

25 [00377] The binding of soluble Fc γ RIIIA tetrameric complex was also assayed using magnetic beads coated with Fc Fragment purified from IgG1 (FIG. 4). Soluble Fc γ RIIIA tetrameric complex binds to the IgG1 Fc-coated beads, under conditions in which monomer binding is not detected. Specificity of binding was shown by pre-incubating the receptor complex, with an anti-Fc γ RIIIA monoclonal antibody, LNK16, which blocks Fc binding.
30 This assay further confirms that soluble Fc γ RIIIA tetrameric complex binds monomeric IgG through its normal ligand binding site, and the avidity of the receptor is increased due to multiple binding sites within the complex.

6.3 CONSTRUCTION OF YEAST STRAIN FOR DISPLAY OF MUTANT IgG1 Fc DOMAINS

MATERIALS AND METHODS

[00378] The pYD1 vector (Invitrogen) is derived directly from a yeast replicating
5 vector, pCT302 (Shusta, *et al.*, 2000 *Nat. Biotechnol.* 18: 754-759, that has been
successfully used to display T-cell receptors and a number of scFVs. This plasmid is
centromeric and harbors the TRP1 gene enabling a relatively constant copy number of 1-2
plasmids per cell in a *trp1* yeast strain. Directional cloning into the polylinker places the
gene of interest under the control of the *GAL1* promoter and in-frame with *AGA2*. Fusion of
10 the IgG Fc domain to the yeast Aga2p results in the extracellular secretion of the Aga2-Fc
fusion protein and subsequent display of the Fc protein on the cell wall via disulfide
bonding to the yeast Aga1p protein, which is an integral cell wall protein.

[00379] In order to optimize the display levels, different fragments from the IgG1
heavy chain were amplified by PCR and cloned into pYD1. Specifically, the Fc region of
15 the IgG1 heavy chain (allotype IG1m(a); amino acids 206- 447) was amplified by PCR
(Table 1) from the IMAGE clone 182740, digested with EcoRI/SaII and ligated into the
pYD1 vector (Invitrogen). The initial clone from IMAGE contained a deletion of a single
nucleotide at position 319 which was corrected by *in vitro* site directed mutagenesis to
construct pYD-GIF206 (Quickchange, Stratagene).

20 [00380] The CH1-CH3 fragment (amino acids 118-447) was amplified from the
heavy chain clone of the MAb B6.2 in the pCINEO vector using a 5' oligo (mcr025;chl(f))
and a 3' oligo (H021) (See Table 8). The fragment was digested with BamHI/NotI and
ligated into the pYD1 vector to construct pYD-CH1.

[00381] FIG. 5, shows a schematic presentation of the constructs. The CH1-CH3
25 construct contains the CH1 domain in addition to the hinge-CH2-CH3 domains of the heavy
chain, GIF206 contains 6 amino acid residues upstream of the hinge and GIF227 starts
within the hinge region at an endogenous proteolytic cleavage site (Jendeborg *et al.*, 1997 *J.*
Immunol. Meth. 201: 25-34).

6.4 IMMUNOLOCALIZATION AND CHARACTERIZATION OF Fc DOMAINS ON THE YEAST CELL WALL

MATERIALS AND METHODS

[00382] Constructs containing the Aga2p-Fc fusion proteins and a control vector,
pYDI, lacking any insert, were transformed into the yeast strain EBY100 (Invitrogen),

MATa ura3-52 trp1 leu2Δ1 his3Δ200 pep4::HIS3 prb1Δ1.6R can1 GAL::GAL-AGA1, using a standard lithium acetate yeast transformation protocol (Gietz *et al.*, 1992 *Nucleic Acids Res.* 20: 1425) Subsequently, tryptophan prototrophs were selected on defined media. Amplification of independent cell populations and induction of Agalp and the Aga2p-Fc fusion proteins were accomplished by growth in glucose, followed by growth in media containing galactose as the primary carbon source for 24-48 hrs at 20°C. Growth in galactose induces expression of the Aga2-Fc fusion proteins via the GAL1 promoter, which subsequently leads to the display of the Fc fusion proteins on the yeast cell surface.

RESULTS

10 FACS Analysis of Fc Fusion Proteins

[00383] Expression of Fc fusion proteins on the yeast cell surface was analyzed by immunostaining using a PE-conjugated polyclonal F(ab)₂ goat anti-human FcγR and HP6017 (Sigma) antibody (Jackson ImmunoResearch Laboratories, Inc.). Fluorescence microscopy shows peripheral staining for the three Fc fusion proteins. The control strain, harboring vector alone, shows little or no staining (data not shown). FACS analysis was used to quantitate the staining (FIG. 6). The yeast strain containing the CH1-CH3 fusion demonstrated the highest percentage of cells stained with both antibodies (FIG. 6B and F). The GIF227 construct showed the greatest mean fluorescence intensity (FIG. 6, panels C and G).

20 Characterization of the Binding of Fc Fusion Proteins Expressed on the Yeast cell Surface

[00384] The natural context of the Fc and FcγR proteins places the receptor on the cell surface and the Fc as the soluble ligand; however, the yeast Fc surface display reverses the geometry of the natural interaction. Detection of the IgG1 Fc proteins on the surface of the yeast cell wall is complicated by both the low affinity of the FcγR for its ligand and the reverse geometry inherent in the display system. Although the latter point cannot be altered, the avidity of the ligand was improved as explained above by forming soluble FcγR tetrameric complexes, which allows detection of FcγR binding to the Fc fusion proteins expressed on the surface yeast cell wall.

30 [00385] To characterize the binding of soluble tetrameric FcγR complexes to the surface displayed Fc fusion proteins, yeast cells expressing different Fc constructs were incubated with the soluble rFcγRIIIA tetrameric complex and analyzed by FACS. Yeast cells harboring pYD-CH1, displaying the wild type CH1-CH3 construct were bound by the soluble rFcγRIIIA tetrameric complex as shown by FACS analysis. The GIF206 and

GIF227 strains, however, showed little or no binding to the soluble rFc γ RIIIA tetrameric complex as shown by FACS analysis (data not shown).

[00386] Mutations in the Fc region that block binding to the Fc γ Rs have been identified (Shields *et al.*, 2001; *J Biol.Chem.* 276: 6591-6604). One of these mutations, D265A, was incorporated into pYD-CH1 and this mutant was expressed on the yeast cell surface. These cells were incubated with the soluble Fc γ RIIIA tetrameric complex using a high concentration of ligand (0.15 mM of Fc; 7.5 mM of D265A) FACS analysis indicated that soluble Fc γ RIIIA tetrameric complex bound to wild type Fc (FIG. 7A) but soluble Fc γ RIIIA tetrameric complex did not bind to the D265A-Fc mutant indicating that Fc γ R is interacting with the normal FcR binding site in the lower hinge-CH2 region (FIG. 7B).

[00387] Antibodies against the Fc γ RIIIA ligand binding site blocked binding of the soluble Fc γ RIIIA tetrameric complex to the wild type Fc protein displayed on the yeast cell surface wall, as analyzed by FACS (FIG. 8). The binding of soluble Fc γ RIIIA tetrameric complex was blocked by the 3G8 antibody, as well as the LNK16 antibody, another anti-Fc γ RIIIA monoclonal antibody (Advanced Immunological) (Tam *et al.*, 1996 *J. Immunol.* 157:, 1576-1581) and was not blocked by an irrelevant isotype control. Therefore, binding of soluble Fc γ RIIIA tetrameric complex to the Fc proteins displayed on the yeast cell surface occurs through the normal ligand binding site. The limited binding of the Fc γ RIIIA tetrameric complex indicates that a subpopulation of cells have a correctly folded Fc that is accessible to Fc γ R. There are numerous reasons why only a subpopulation of cells may be able to bind the ligand, for example, they may be at different stages of cell cycle or the fusion proteins may not have been exported.

[00388] In order to determine the dissociation constant of the Fc γ RIIIA-tetramer binding to the Fc fusion proteins on the yeast cell surface, the binding of a range of Fc γ RIIIA tetrameric complex was analyzed using FACS. Fc γ RIIIA tetrameric complex was titrated at concentrations of 1.4 μ M to 0.0006 μ M. Using the mean fluorescence intensity as a measure of binding affinity and nonlinear regression analysis, the K_D was determined to be 0.006 μ M (+/- 0.001) (data not shown).

6.5 CONSTRUCTION OF Fc MUTANT LIBRARY

[00389] A mutant Fc library was constructed using primers flanking the Fc fragment in the Fc-CH1 construct and error-prone PCR (Genemorph, Stratagene). The CH1-CH3 insert in vector pYD-CHI was amplified using a mutagenic PCR (Genemorph, Stratagene). Five reactions were carried out using the pYD-upstream and pYD-downstream primers (Invitrogen). The resultant amplified fragment was digested with XHOI/BamHI and ligated

into pYD1. The ligation reaction was then transformed into XL10 ultracompetent cells (Stratagene), which resulted in $\sim 1 \times 10^6$ transformants, with 80% of the transformants containing inserts.

[00390] Sequence analysis of 28 random plasmids from the library indicated a mutation frequency ~ 2 -3 mutations/kb with a breakdown of 40% conserved nucleotide changes and 60% of the mutations resulting in amino acid changes.

[00391] The library was transformed into the yeast strain EBY100, MAT α *ura3-52 trp1 leu2 Δ 1 his3 Δ 200 pep4::HIS3 prb1 Δ 1.6R can1 GAL GAL-AGA 1::URA3* to a high efficiency, $\sim 3.3 \times 10^5$ transformants/ μ g, in 30 independent transformation reactions to create a total of $\sim 10^7$ yeast transformants (Gietz, *et al.*, 1992, *Nucleic Acids Res.* 20: 1425). The library was pooled and amplified by growth in glucose.

6.6 SELECTION AND ANALYSIS OF Fc MUTANTS

MATERIALS AND METHODS

ELISA assay for screening Fc mutants

[00392] ELISA plates (Nunc F96 MaxiSorp Immunoplate) were coated with 50 ml/well of 0.5 mg/ml BSA-FITC in carbonate buffer at 4°C, and allowed to incubate overnight. Plates were washed with 1X PBS/0.1% Tween 20 (PBST) 3 times. 200 ml/well of PBST/0.5% BSA was added and the plates were incubated for 30 mins at room temperature. Plates were washed three additional times with PBST. 50 ml/well of 1:4 diluted 4-4-20 antibody (approximately 3 mg/mL which would lead to a final concentration of 0.7-0.8 mg/well) either wild type or containing an Fc mutant, was added from conditional medium in PBST/0.5% BSA and allowed to incubate for 2 hrs at room temperature. Plates were washed with PBST three times. Purified, biotinylated monomeric Fc γ RIIIA at 3 mg/ml (in PBST/0.5% BSA) was added (50 μ l/well) to the plates and allowed to incubate for 1.5 hours at room temperature. Plates were washed with PBST three times. 50 ml/well of a 1:5000 dilution of Streptavidin-HRP (Pharmacia, RPN 123v) in PBST/0.5% BSA was added and the plates were incubated for 30 minutes at room temperature. Plates were washed with PBST three times. 80 ml/well of TMB reagent (BioFX) was then added to the plates, and allowed to incubate for 10-15 minutes at room temperature in a dark place. The reactions were finally stopped by adding 40 ml/well of stop solution (0.18 M sulfuric acid). Plates were then monitored for absorbance at 450 nm. After the first screen, the interesting candidates were further confirmed by serial titration of 4-4-20-Fc mutants in the immuno-complex based binding ELISA. A few modifications were made in this ELISA. For coating the plates, 2 mg/ml BSA-FITC was used. Based on IgG quantitation results,

diluted 4-4-20Fc (wild type or mutants) from conditional medium was added to a final concentration of 1, 0.5, 0.25, 0.125, 0.063, and 0 mg/ml in PBST-/0.5% BSA.

FACS Screen for the Cell Surface displayed Fc proteins

[00393] Cells were grown in at least 10 mls of HSM-Trp-Ura pH 5.5 with glucose for 5
16-24 hrs or until OD₆₀₀ was greater than 2.0. Cells were spun down at ~2000 rpm for 5
minutes. Cells were resuspended in an equal volume of HSM-Trp-Ura, pH 7.0 with
galactose. In a 125 ml flask, 36 mls of galactose media was added, and inoculated with 9
mls of culture, which was incubated at 20°C with shaking for 24-48 hrs. Growth was
monitored by measuring OD₆₀₀ at 8-16 hr intervals. Cells were harvested at 2K rpm for 5
10 minutes, and resuspended in an equal volume of 1XPBS, pH 7.4.

[00394] *Equilibrium screen:* An appropriate amount of cells was incubated while
maintaining an excess of ligand. For example, it is preferred to start with a number of cells
needed to ensure 10-fold coverage of the library. For the first sort with a library containing
10⁷ transformants, 10⁸ cells should be used. In fact it is best to start with 10⁹ cells to
15 compensate for loss during the staining protocol.

[00395] Incubation was typically done in a 1.5 mL tube in volumes of 20-100 mls for
1 hour at 4°C in the dark on a rotator (incubation buffer: 1XPBS pH7.4; 1 mg/ml BSA).
Cells were washed once in 500 ml of incubation buffer and spun down at 4K rpm for 2.5
minutes. Cells were resuspended in 100 ml incubation buffer and incubated with the second
20 staining reagent. For Fc-CH1, a F(ab)₂ goat anti-hFc F(ab)₂-FITC antibody (Jackson
Immunoresearch Laboratories, Inc.) can be used to stain for CH1 expression. Staining was
done with 1 mL for 30 minutes. Cells were washed additionally in 500 mL of incubation
buffer and spun down at 4K rpm for 2.5 minutes, resuspended in 1 mL IX PBS 1 mg/mL
BSA and analyzed by FACS.

25 [00396] Typical equilibrium screen sort gates and number of cells collected are
shown in Table 9.

TABLE 9. SORT GATES AND NUMBER OF CELLS SORTED

Sort	Gate	total cells screened	cells collected
1 st	5%	10 ⁸	5 x 10 ⁶
2 nd	1%	10 ⁷	1 x 10 ⁵
3 rd	0.2%	10 ⁷	2 x 10 ⁴
4 th	0.2%	10 ⁷	2 x 10 ⁴

[00397] After the 3rd and 4th sorts, cells were plated directly on -trp-ura plates to identify individual mutants. This typically recovered ~200-400 colonies per plate. After collection the cells were placed in 10 mLs of glucose media in a 50 mL conical tube and grown at 30°C. The whole procedure was repeated iteratively.

5 RESULTS

FACS analysis of Fc Mutants

[00398] After induction in galactose media, cells were harvested and co-stained with soluble Fc γ R11A tetrameric complex-PE labeled and F(ab₂) of mouse anti-human Fc-FITC labeled (Jackson Immunoresearch Laboratories, Inc.). Cells were analyzed by FACS and
10 sort gates were used to select the cells that showed the highest affinity for the soluble Fc γ R11A tetrameric complex relative to the amount of Fc expression on the cell surface (FIG. 9). For example, a cell containing a mutant Fc that binds better to the soluble Fc γ R11A tetrameric complex may express fewer Fc fusion proteins on the yeast cell surface, and this cell will be in the lower left hand corner of the sort gate.

15 [00399] Four consecutive sorts were done to enrich for those mutants that showed the highest affinity for the soluble Fc γ R11A tetrameric complex. The gates for each successive sort were 5.5%, 1%, 0.2% and 0.1%. After the last sort, cells were plated onto selective media and individual colonies were isolated. Each individual colony represented a clonal population of cells harboring a single Fc mutant within the Aga2-Fc fusion protein. Initially
20 32 independent colonies were picked and tested by FACS for binding to soluble Fc γ R11A tetrameric complex (FIG. 10). Eighteen mutants showed an increase in binding intensity as measured by the percentage of cells bound by soluble Fc γ R11A tetrameric complex and the mean fluorescence intensity of the bound cells.

[00400] Mutations showing an increase in binding to Fc γ R11A were also tested for
25 binding to soluble Fc γ R11B tetrameric complex (FIG. 10). Most mutations that lead to an increase in binding to the soluble Fc γ R11A tetrameric complex also resulted in detection of Fc γ R11B tetrameric complex staining (FIG. 10). Based on both previous physical and genetic data, some mutations that increase binding to Fc γ R11A, are expected to also increase binding to Fc γ R11B (Shields *et al.*, 2001, *J Biol.Chem.* 276: 6591-6604;
30 Sondermann *et al.*, 2000, *Nature* 406: 267-273).

Analysis of mutants in a 4-4-20 MAb produced in a human cell line.

[00401] Isolation and analysis of mutations in the yeast system allows for fast identification of novel mutant alleles. The use of a heterologous system to isolate mutations could result in the identification of mutations that enhance binding through an alteration that

results in misfolding or alteration in glycosylation that is specific to yeast. To analyze the Fc mutations in an immunoglobulin molecule that is produced in human cells, the mutants were subcloned into a mammalian expression vector, containing the heavy chain of the anti-fluorescein monoclonal antibody, 4-4-20 (Kranz *et al.*, 1982 *J.Biol.Chem*, 257(12): 6987-6995). The mutant 4-4-20 heavy chains were transiently coexpressed with the light chain clones in the human kidney cell line (293H). Supernatants were collected and analyzed by ELISA (FIG. 11).

[00402] According to the ELISA assay, the majority of the mutants that were identified as having an enhanced affinity for the soluble monomeric Fc γ RIIA complex, in the secondary FACS analysis, also showed an increase in binding to the soluble Fc γ RIIA tetrameric complex when present in the Fc region of the 4-4-20 monoclonal antibody produced in the human cell line (FIG. 11A). Two mutants, number 16 and number 19, however, showed a decrease in binding to the soluble Fc γ RIIA monomeric complex.

[00403] Table 10, summarizes the mutations that have been identified and their corresponding binding characteristics to Fc γ RIIA and Fc γ RIIB, as determined by both yeast display based assays and ELISA. In Table 6, the symbols represent the following: • corresponds to a 1-fold increase in affinity; + corresponds to a 50% increase in affinity; - corresponds to a 1-fold decrease in affinity; → corresponds to no change in affinity compared to a comparable molecule comprising a wild-type Fc region.

20 **TABLE 10: MUTATIONS IDENTIFIED AND BINDING CHARACTERISTICS**

Clone #	Mutation sites	Domain	IIIA binding	IIIB binding
4	A339V, Q347H	CH2, CH3	+	+
5	L251P, S415I	CH2, CH3	+	+
7	Aga2p-T43I	Note: This is a mutation in Aga2P that enhances display.		Aga2p-T43I
8	V185M, K218N, R292L, D399E	CH1,hinge,CH2, CH3	no change	-
12	K290E, L142P	CH1,CH2	+	not tested
16	A141V, H268L, K288E, P291S	CH1,CH2	-	not tested
19	L133M, P150Y, K205E, S383N, N384K	CH1,CH2,CH3	-	not tested

Clone #	Mutation sites	Domain	IIIA binding	IIB binding
21	P396L	CH3	•	•+
25	P396H	CH3	•••	••
6	K392R	CH3	no change	no change
15	R301C, M252L, S192T	CH1,CH2	-	not tested
17	N315I	CH2	no change	not tested
18	S132I	CH1	no change	not tested
26	A162V	CH1	no change	not tested
27	V348M, K334N, F275I, Y202M, K147T	CH1,Ch2	+	+
29	H310Y, T289A, G337E	CH2	-	not tested
30	S119F, G371S, Y407N, E258D	CH1,CH2,CH3	+	no change
31	K409R, S166N	CH1,CH3	no change	not tested
20	S408I, V215I, V125I	CH1,hinge,CH3	+	no change
24	G385E, P247H	CH2, CH3	•••	+
16	V379M	CH3	••	no change
17	S219Y	Hinge	•	-
18	V282M	CH2	•	-
31	F275I, K334N, V348M	CH2	+	no change
35	D401V	CH3	+	no change
37	V280L, P395S	CH2	+	-
40	K222N	Hinge	•	no change
41	K246T, Y319F	CH2	•	no change
42	F243I, V379L	CH2,CH3	•+	-
43	K334E	CH2	•+	-
44	K246T, P396H	CH2,CH3	•	••+
45	H268D, E318D	CH2	•+	•••••

Clone #	Mutation sites	Domain	IIIA binding	IIB binding
49	K288N, A330S, P396L	CH2,CH3
50	F243L, R255L, E318K	CH2	.	-
53	K334E, T359N, T366S	CH2,CH3	.	no change
54	I377F	CH3	•+	+
57	K334I	CH2	.	no change
58	P244H, L358M, V379M, N384K, V397M	CH2,CH3	•+	•+
59	K334E, T359N, T366S (independent isolate)	CH2,CH3	•+	no change
61	I377F (independent isolate)	CH3	...	••+
62	P247L	CH2	••	••+
64	P217S, A378V, S408R	Hinge, CH3	••	••••+
65	P247L, I253N, K334N	CH2	...	••+
66	K288M, K334E	CH2	...	-
67	K334E, E380D	CH2,CH3	•+	-
68	P247L (independent isolate)	CH2	+	••••
69	T256S, V305I, K334E, N390S	CH2,CH3	•+	no change
70	K326E	CH2	•+	••+
71	F372Y	CH3	+	••••+
72	K326E (independent isolate)	CH2	+	••
74	K334E, T359N, T366S (independent isolate)	CH2,CH3	••	no change
75	K334E (independent isolate)	CH2	••+	no change
76	P396L (independent isolate)	CH3	•+	no change
78	K326E (independent isolate)	CH2	••	•••+
79	K246L, K334N	CH2	.	••••
80	K334E (independent isolate)	CH2	.	no change
81	T335N, K370E, A378, T394M, S424L	CH2,CH3	.	no change
82	K320E, K326E	CH2	.	.

Clone #	Mutation sites	Domain	IIIA binding	IIB binding
84	H224L	Hinge	•	•••••
87	S375C, P396L	CH3	•+	••••+
89	E233D, K334E	CH2	•+	no change
91	K334E (independent isolate)	CH2	•	no change
92	K334E (independent isolate)	CH2	•	no change
94	K334E, T359N, T366S, Q386R	CH2	•	no change

[00404] Analysis of soluble Fc γ R IIB tetrameric complex binding shows that 7 out of the 8 mutants that showed an increase in binding to the soluble Fc γ R IIIA tetrameric complex also had an increased binding to the soluble Fc γ R IIB tetrameric complex (FIG. 11B). One mutant, number 8, showed a decrease in binding to the soluble Fc γ R IIB tetrameric complex. Three of the mutants show no difference in binding to either the soluble Fc γ R IIIA tetrameric complex or the soluble Fc γ R IIB tetrameric complex, possibly due to mutations that result in yeast specific alterations.

6.7 ADCC ASSAY OF Fc MUTANTS

[00405] *Effector cell preparation:* Peripheral blood mononuclear cells (PBMC) were purified by Ficoll-Paque (Pharmacia, 17-1440-02) Ficoll-Paque density gradient centrifugation from normal peripheral human blood (Biowhittaker/Poietics, 1W-406). Blood was shipped the same day at ambient temperature, and diluted 1:1 in PBS and glucose (1g/1L) and layered onto Ficoll in 15 mL conical tubes (3 mL Ficoll; 4 mL PBS/blood) or 50mL conical tubes (15mL: Ficoll; 20mL PBS/blood). Centrifugation was done at 1500 rpm (400 rcf) for 40 minutes at room temperature. The PBMC layer was removed (approximately 4-6 mL from 50 mL conical tube) and diluted 1:10 in PBS (which contains no Ca²⁺ or Mg²⁺) in a 50 mL conical tube, and spun for an additional ten minutes at 1200 rpm (250 rcf) at room temperature. The supernatant was removed and the pellets were resuspended in 10-12 mL PBS (which contains no Ca²⁺ or Mg²⁺), transferred to 15 mL conical tubes, and spun for another 10 minutes at 1200 rpm at room temperature. The supernatant was removed and the pellets were resuspended in a minimum volume (1-2 mL) of media (Isocove's media (IMDM) + 10% fetal bovine serum (FBS), 4 mM Gln, Penicillin/Streptomycin (P/S)). The resuspended PBMC were diluted to the appropriate volume for the ADCC assay; two fold dilutions were done in an ELISA 96 well plate (Nunc

F96 MaxiSorp Immunoplate). The yield of PBMC was approximately 3×10^7 cells per 40-50 mL of whole blood.

[00406] *Target cell preparation:* Target cells used in the assay were SK-BR-3 (ATCC Accession number HTB-30; Trempe *et al.*, 1976, *Cancer Res.* 33-41), Raji (ATCC Accession number CCL-86; Epstein *et al.*, 1965, *J. Natl. Cancer Inst.* 34: 231-40), or Daudi cells (ATCC Accession number CCL-213; Klein *et al.*, 1968, *Cancer Res.* 28: 1300-10) (resuspended in 0.5 mL IMDM media) and they were labeled with europium chelate bis(acetoxymethyl) 2,2':6',2" terpyridine 6,6' dicarboxylate (BATDA reagent; Perkin Elmer DELFIA reagent; C136-100). K562 cells (ATCC Accession number CCL-243) were used as control cells for NK activity. The Daudi and Raji cells were spun down; the SK-BR-3 cells were trypsinized for 2-5 minutes at 37°C, 5% CO₂ and the media was neutralized prior to being spun down at 200-350 G. The number of target cells used in the assays was about 4×10^6 cells and it did not exceed 5×10^6 since labeling efficiency was best with as few as 2×10^6 cells. Once the cells were spun down, the media was aspirated to 0.5 mL in 15 mL Falcon tubes. 2.5 µl of BATDA reagent was added and the mixture was incubated at 37°C, 5% CO₂ for 30 minutes. Cells were washed twice in 10 mL PBS and 0.125 mM sulfinpyrazole ("SP"; SIGMA S-9509); and twice in 10 mL assay media (cell media + 0.125 mM sulfinpyrazole). Cells were resuspended in 1 mL assay media, counted and diluted.

[00407] When SK-BR-3 cells were used as target cells after the first PBS/SP wash, the PBS/SP was aspirated and 500 µg/mL of FITC was added (PIERCE 461110) in IMDM media containing SP, Gln, and P/S and incubated for 30 minutes at 37°C, 5% CO₂. Cells were washed twice with assay media; resuspended in 1 mL assay media, counted and diluted.

[00408] *Antibody Opsonization:* Once target cells were prepared as described *supra*, they were opsonized with the appropriate antibodies. In the case of Fc variants, 50 µL of 1×10^5 cells/mL were added to 2x concentration of the antibody harboring the Fc variant. Final concentrations were as follows: Ch-4-4-20 final concentration was 0.5-1 µg/mL; and Ch4D5 final concentration was 30 ng/mL-1 ng/mL.

[00409] Opsonized target cells were added to effector cells to produce an effector:target ratio of 75:1 in the case of the 4-4-20 antibodies with Fc variants. In the case of the Ch4D5 antibodies with Fc variants, effector: target ratio of 50:1 or 75:1 were achieved. Effective PBMC gradient for the assay ranges from 100:1 to 1:1. Spontaneous release (SR) was measured by adding 100 µL of assay media to the cells; maximal release (MR) was measured by adding 4% TX-100. Cells were spun down at 200 rpm in a

Beckman centrifuge for 1 minute at room temperature at 57 G. Cells were incubated for 3-3.5 hours at 37°C, 5%CO₂. After incubation, the cells were spun at 1000 rpm in a Beckman centrifuge (about 220xg) for five minutes at 10°C. 20 µl of supernatant was collected; 200µL of Eu solution was added and the mixture was shaken for 15 minutes at room
5 temperature at 120 rpm on a rotary shaker. The fluorescence was quantitated in a time resolved fluormeter (Victor 1420, Perkin Elmer)

RESULTS

[00410] As described above, the variant Fc regions were subcloned into a mammalian
10 expression vector, containing the heavy chain of the anti-fluoresceine monoclonal antibody, 4-4-20 (Kranz *et al.*, 1982 *J.Biol.Chem.*, 257(12): 6987-6995). The variant 4-4-20 heavy chains were transiently coexpressed with the light chain clones in the human kidney cell line (293H). Supernatants were collected and analyzed using the ADCC assay. FIG. 12 shows that ADCC activity of the mutants is concentration-dependent. As summarized in
15 Table 8, five immunoglobulins with variant Fc regions had an enhanced ADCC activity relative to wild type ch 4-4-20. The five mutants were as follows: MGFc-27 (G316D, A378V, D399E); MGFc-31 (P247L, N421K); MGFc-10 (K288N, A330S, P396L); MGFc-28 (N315I, V379M, T394M); MGFc-29 (F243I, V379L, G420V).

[00411] Additonal 4-4-20 immunoglobulins with variant Fc regions were assayed for
20 their ADCC activity relative to a 4-4-20 immunoglobulin with a wild-type Fc region. These results are summarized in Table 11.

[00412] ADCC assays were also carried out using the same protocol as previously described for the 4-4-20 antibody, however, the variant Fc regions were cloned into a humanized antibody (Ab4D5) which is specific for the human epidermal growth factor
25 receptor 2 (HER2/neu). In this case, SK-BR-3 cells were used as the target cells that were opsonized with a HER2/neu antibody carrying a variant Fc region. HER2/neu is endogenously epxressed by the SK-BR-3 cells and therefore present on the surface these cells. FIG. 13 shows the ADCC activity of HER2/neu antibodies carrying variant Fc regions. Table 12 summarizes the results of ADCC activity of the mutants in the context of
30 the HER2/neu antibody. Normalization was carried out by comparing the concentration of the mutant to the wildtype antibody required for a specific value of percent cell lysis.

[00413] As shown in FIG. 13A, MGFc-5 (V379M), MGFc-9 (P243I, V379L), MGFc-10 (K288N, A330S, P396L), MGFc-13 (K334E, T359N, T366S), and MGFc-27

(G316D, A378V, D399E) mutants that were cloned in to the humanized anti-HER2/neu antibody exhibited a higher % specific lysis of SK-BR-3 cells relative to the wild antibody.

Table 11. SUMMARY OF ADCC ACTIVITY OF MUTANTS

Label	Fc Variant		1ug/ml		ADCC		0.5ug/ml	
	Ref	Amino Acid Variation	% specific lysis	Normalized	% specific lysis	Normalized	% specific lysis	Normalized
MGFc-27	2C4	G316D, A378V, D399E	33%	2.24	22%	3.60		
MGFc-31	3B9	P247L, N421K	30%	2.05	17%	2.90		
MGFc-10	1E1	K288N, A330S, P396L	24%	1.66	10%	1.67		
MGFc-28	2C5	N315I, V379M, T394M	20%	1.37	10%	1.69		
MGFc-29	3D11	F243I, V379L, G420V	20%	1.35	7%	1.17		
ch4-4-20 (P54008)			15%	1.00	6%	1.00		
MGFc-35	3D2	R255Q, K326E	11%	0.79	3%	0.53		
MGFc-36	3D3	K218R, G281D, G385R	10%	0.67	5%	0.78		
MGFc-30	3A8	F275Y	9%	0.64	2%	0.37		
MGFc-32	3C8	D280E, S354F, A431D, L441I	9%	0.62	4%	0.75		
MGFc-33	3C9	K317N, F423deleted	3%	0.18	-1%	-0.22		
MGFc-34	3B10	F241L, E258G	-1%	-0.08	-4%	-0.71		
MGFc-26		D265A	1%	0.08	-3%	-0.45		

Table 12: SUMMARY OF MUTANTS

Fc Variant	Amino Acid changes	FCR3A, K_D/K_{off}	FCR2B, K_D/K_{off}	ELISA IIIA binding	ELISA IIB binding	Phagocytosis (mutant/WT)	4-4-20 ADCC (mutant/wt)	Anti-HER2 ADCC (mutant/wt)
Wt	none	198/0.170	94/0.094	1	1	1	1	1
MGFc 5	V379M	160/0.167	70/0.10	2X	N/C	0.86	2.09	1.77
MGFc 9	P243I, V379L	99.7/0.105	120/0.113	1.5X	reduced	?	2.25	2.04
MGFc 10	K288N, A330S, P396L	128/0.115	33.4/0.050	5X	3X	1.2	2.96	2.50
MGFc 11	F243L, R255L	90/0.075	74.7/0.09	1x	reduced	0.8	2.38	1.00
MGFc 13	K334E, T359N, T366S	55.20.128	72/0.11	1.5X	N/C	1	1.57	3.67
MGFc 14	K288M, K334E	75.4/0.1	95.6/0.089	3X	reduced	1	1.74	1.6
MGFc 23	K334E, R292L	70.2/0.105	108/0.107			1	2.09	6.88
MGFc 27	G316D, A378V, D399E	72/0.117	46/0.06	1.5X	14X	1.4	3.60	
MGFc 28	N315I, A379M, D399E			1X	9X	1.37	1.69	1.00
MGFc 29	P243I, V379L, G420V	108/0.082	93.4/0.101	2.5X	7X	0.93	1.17	1.00
MGFc 31	P247L, N421K	62/0.108	66/0.065	3X	N/C	1.35	2.90	1.00
MGFc 37	K248M	154/0.175	100/0.091	1.4X	reduced	0.98	3.83	0.67
MGFc 38	K392T, P396L	84/0.104	50/0.041	4.5X	2.5X	1.4	3.07	2.50
MGFc 39	E293V, Q295E, A327T	195/0.198	86/0.074	1.4X	reduced	1.5	4.29	0.50
MGFc 40	K248M	180/0.186	110/0.09	1.4X	reduced	1.14	4.03	
MGFc 41	H268N, P396L	178/0.159	46.6/0.036	2.2X	4.5X	1.96	2.24	0.67
MGFc 43	Y319F, P352L, P396L	125/0.139	55.7/0.041	3.5X	2X	1.58	1.09	

6.8 ANALYSIS OF KINETIC PARAMETERS OF Fc MUTANTS

[00414] Kinetic parameters of the binding of ch4-4-20 antibodies harboring Fc mutants to FcγRIIIA and FcγRIIB were analyzed using a BIAcore assay (BIAcore instrument 1000, BIAcore Inc., Piscataway, N.J.). The FcγRIIIA used in this assay was a
5 soluble monomeric protein, the extracellular region of FcγRIIIA joined to the linker-AVITAG sequence as described in Section 6.2 *supra*. The FcγRIIB used in this assay was a soluble dimeric protein prepared in accordance with the methodology described in U.S. Provisional Application No. 60/439,709 filed on January 13, 2003, which is incorporated herein by reference. Briefly, the FcγRIIB used was the extracellular domain of FcγRIIB
10 fused to the hinge-CH2-CH3 domain of human IgG2.

[00415] BSA-FITC (36 μg/mL in 10mM Acetate Buffer at pH 5.0) was immobilized on one of the four flow cells (flow cell 2) of a sensor chip surface through amine coupling chemistry (by modification of carboxymethyl groups with mixture of NHS/EDC) such that
about 5000 response units (RU) of BSA-FITC was immobilized on the surface. Following
15 this, the unreacted active esters were "capped off" with an injection of 1M Et-NH₂. Once a suitable surface was prepared, ch 4-4-20 antibodies carrying the Fc mutations were passed over the surface by one minute injections of a 20 μg/mL solution at a 5 μL/mL flow rate. The level of ch-4-4-20 antibodies bound to the surface ranged between 400 and 700 RU. Next, dilution series of the receptor (FcγRIIIA and FcγRIIB-Fc fusion protein) in HBS-P
20 buffer (10mM HEPES, 150 mM NaCl, .005% Surfactant P20, 3mM EDTA, pH 7.4) were injected onto the surface at 100 μL/min. Antibody regeneration between different receptor dilutions was carried out by single 5 second injections of 100mM NaHCO₃ pH 9.4; 3M NaCl.

[00416] The same dilutions of the receptor were also injected over a BSA-FITC
25 surface without any ch-4-4-20 antibody at the beginning and at the end of the assay as reference injections.

[00417] Once an entire data set was collected, the resulting binding curves were globally fitted using computer algorithms supplied by the manufacturer, BIAcore, Inc. (Piscataway, NJ). These algorithms calculate both the K_{on} and K_{off} , from which the
30 apparent equilibrium binding constant, K_D is deduced as the ratio of the two rate constants (*i.e.*, K_{off}/K_{on}). More detailed treatments of how the individual rate constants are derived can be found in the BIAevaluation Software Handbook (BIAcore, Inc., Piscataway, NJ).

- [00418] Binding curves for two different concentrations (200 nM and 800 nM for FcγRIIIA and 200 nM and 400nM for FcγRIIB fusion protein) were aligned and responses adjusted to the same level of captured antibodies, and the reference curves were subtracted from the experimental curves. Association and dissociation phases were fitted separately.
- 5 Dissociation rate constant was obtained for interval 32-34 sec of the dissociation phase; association phase fit was obtained by a 1:1 Langmuir model and base fit was selected on the basis R_{\max} and χ^2 criteria.

RESULTS

- 10 [00419] FIG. 14 shows the capture of ch 4-4-20 antibodies with mutant Fc regions on the BSA-FTIC-immobilized sensor chip. 6 μL of antibodies at a concentration of about 20 μg/mL were injected at 5 μL/min over the BSA-FITC surface. FIG. 15 is a sensogram of real time binding of FcγRIIIA to ch-4-4-20 antibodies carrying variant Fc regions. Binding of FcγRIIIA was analyzed at 200 nM concentration and resonance signal responses were
- 15 normalized at the level of the response obtained for the wild type ch-4-4-20 antibody. Kinetic parameters for the binding of FcγRIIIA to ch-4-4-20 antibodies were obtained by fitting the data obtained at two different FcγRIIIA concentrations, 200 and 800 nM (FIG. 16). The solid line represents the association fit which was obtained based on the K_{off} values calculated for the dissociation curves in interval 32-34 seconds. K_D and K_{off}
- 20 represent the average calculated from the two different FcγRIIIA concentrations used. FIG. 17 is a sensogram of real time binding of FcγRIIB-Fc fusion protein to ch-4-4-20 antibodies carrying variant Fc regions. Binding of FcγRIIB-Fc fusion protein was analyzed at 200 nM concentration and resonance signal responses were normalized at the level of the response obtained for the wild type ch-4-4-20 antibody. Kinetic parameters for the binding
- 25 of FcγRIIB-Fc fusion protein to ch-4-4-20 antibodies were obtained by fitting the data obtained at two different FcγRIIB-Fc fusion protein concentrations, 200 and 400 nM (FIG. 18). The solid line represents the association fit which was obtained based on the K_{off} calculated for the dissociation curves in interval 32-34 seconds. K_D and K_{off} represent the average from the two different FcγRIIB-Fc fusion protein concentrations used.
- 30 [00420] The kinetic parameters (K_{on} and K_{off}) that were determined from the BIAcore analysis correlated with the binding characteristic of the mutants as determined by an ELISA assay and the functional activity of the mutants as determined in an ADCC assay. Specifically, as seen in Table 13, mutants that had an enhanced ADCC activity relative to

the wild-type protein, and had an enhanced binding to FcγRIIIA as determined by an ELISA assay had an improved K_{off} for FcγRIIIA (*i.e.*, a lower K_{off}). Therefore, a lower K_{off} value for FcγRIIIA for a mutant Fc protein relative to a wild type protein may be likely to have an enhanced ADCC function. On the other hand, as seen in Table 14, mutants that had an enhanced ADCC activity relative to the wild-type protein, and had a reduced binding for FcγRIIB-Fc fusion protein as determined by an ELISA assay had a higher K_{off} for FcγRIIB - Fc fusion protein.

[00421] Thus, the K_{off} values for FcγRIIIA and FcγRIIB can be used as predictive measures of how a mutant will behave in a functional assay such as an ADCC assay. In fact, ratios of K_{off} values for FcγRIIIA and FcγRIIB-Fc fusion protein of the mutants to the wild type protein were plotted against ADCC data (FIG 19). Specifically, in the case of K_{off} values for FcγRIIIA, the ratio of K_{off} (wt)/ K_{off} (mutant) was plotted against the ADCC data; and in the case of K_{off} values for FcγRIIB, the ratio of K_{off} (mut)/ K_{off} (wt) was plotted against the ADCC data. Numbers higher than one (1) show a decreased dissociation rate for FcγRIIIA and an increased dissociation rate for FcγRIIB -Fc relative to wild type. Mutants that fall within the indicated box have a lower off rate for FcγRIIIA binding and a higher off-rate for FcγRIIB -Fc binding, and possess an enhanced ADCC function.

Table 13. Kinetic parameters of FcRIIIa binding to ch4-4-20Ab obtained by "separate fit" of 200nM and 800nM binding curves

Ch4-4-20Ab	BIAcore Kd,nM	K_{on} 1/Ms	K_{off} , 1/s	ELISA ₅₀ OD	ADCC, %
Wt(0225)	319	6.0×10^5	0.170	0.5	17.5
Mut1(0225)	90	8.22×10^5	0.075	0.37	32
Mut5(0225)	214	8.2×10^5	0.172	0.75	26
Mut6(0225)	264	6.67×10^5	0.175	0.6	23
Mut8(0225)	234	8.3×10^5	0.196	0.5	22
Mut10(0225)	128	9.04×10^5	0.115	1.0	41
Mut12(0225)	111	1.04×10^6	0.115	1.0	37
Mut15(0225)	67.9	1.97×10^6	0.133	1.0	15
Mut16(0225)	84.8	1.60×10^6	0.133	1.0	15
Mut18(0225)	92	1.23×10^6	0.112	1.0	28
Mut25(0225)	48.6	2.05×10^6	0.1	1.0	41
Mut14(0225)	75.4	1.37×10^6	0.1	1.1	28
Mut17(0225)	70.5	1.42×10^6	0.1	1.25	30
Mut19(0225)	100	1.20×10^6	0.120	0.75	11
Mut20(0225)	71.5	1.75×10^6	0.126	0.5	10

Mut23(0225)	70.2	1.43×10^6	0.105	1.25	25

Highlighted mutants do not fit to the group by ELISA or ADCC data.

Table 14. Kinetic parameters of FcRIIB-Fc binding to wild type and mutant ch4-4-20Ab obtained by "separate fit" of 200 nM and 800 nM binding curves.

Ch4-4-20Ab	BIAcore K _d , nM	K _{on} 1/Ms	K _{off} , 1/s	ELISA, OD	ADCC, %
Wt(0225)	61.4		0.085	0.4	17.5
Mut11(0225)	82.3		0.1	0.08	32
Mut5(0225)	50		0.057	0.6	26
Mut6(0225)	66.5		0.060	0.35	23
Mut8(0225)	44.2		0.068	0.25	22
Mut10(0225)	41.3		0.05	1.2	41
Mut12(0225)	40.1		0.051	0.4	37
Mut15(0225)	37.8		0.040	1.55	15
Mut16(0225)	40		0.043	1.55	15
Mut18(0225)	51.7		0.043	1.25	28
Mut25(0225)			0.112	0.08	41
Mut14(0225)	95.6		0.089	0.13	28
Mut17(0225)	55.3		0.056	0.38	30
Mut19(0225)	45.3		0.046	1.0	11
Mut20(0225)	24.1		0.028	0.8	10
Mut23(0225)	108		0.107	0.1	25

5 6.9 SCREENING FOR Fc MUTANTS USING MULTIPLE ROUNDS OF ENRICHMENT USING A SOLID PHASE ASSAY

[00422] The following mutant screens were aimed at identifying additional sets of mutants that show improved binding to FcγRIIIA and reduced binding to FcγRIIB.

Secondary screening of selected Fc variants was performed by ELISA followed by testing
 10 for ADCC in the 4-4-20 system. Mutants were then selected primarily based on their ability to mediate ADCC via 4-4-20 using Fluorescein coated SK-BR3 cells as targets and isolated PBMC from human donors as the effector cell population. Fc mutants that showed a relative increase in ADCC, e.g., an enhancement by a factor of 2 were then cloned into
 15 anti-HER2/neu or anti-CD20 chAbs and tested in an ADCC assay using the appropriate tumor cells as targets. The mutants were also analyzed by BIAcore and their relative K_{off} were determined.

[00423] *Screen 1: Sequential solid phase depletion and selection using Magnetic beads coated with FcγRIIB followed by selection with magnetic beads coated with FcγRIIA.* The aim of this screen was identification of Fc mutants that either no longer bind FcγRIIB or show reduced binding to FcγRIIB. A 10-fold excess of the naïve library (~10⁷ cells) was incubated with magnetic beads ("My One", Dynal) coated with FcγRIIB. Yeast bound to beads were separated from the non-bound fraction by placing the tube containing the mixture in a magnetic field. Those yeast cells that were not bound to the beads were removed and placed in fresh media. They were next bound to beads that were coated with FcγRIIA. Yeast bound to beads were separated from the nonbound fraction by placing the tube containing the mixture in a magnetic field. Nonbound yeast were removed and the bound cells were removed by vigorous vortexing. The recovered cells were regrown in glucose containing media and reinduced in selective media containing galactose. The selection process was repeated. The final culture was then used to harvest DNA. Inserts containing the Fc domain were amplified by PCR and cloned into 4-4-20. Approximately 90 Fc mutants were screened by 4-4-20 ELISA and ADCC assays and the resultant positive mutants are shown in Table 15.

Table 15: Mutants selected by sequential solid phase depletion and selection using Magnetic beads coated with FcγRIIB followed by selection with magnetic beads coated with FcγRIIA.

Mutant	Amino Acid changes
MgFc37	K248M
MgFc38	K392T, P396L
MgFc39	E293V, Q295E, A327T
MgFc41	H268N, P396LN
MgFc43	Y319F, P352L, P396L
	D221E, D270E, V308A, Q311H, P396L,
MgFc42	G402D

20

[00424] *Screens 2&3: Mutants Selected by FACS, Equilibrium and Kinetic Screening:* The first library screen identified a mutation at position 396, changing the amino acid from Proline to Leucine (P396L). This Fc variant showed increased binding to both FcγRIIA and FcγRIIB. A second library was constructed using P396L as a base line. PCR mutagenesis was used to generate ~10⁷ mutants each of which contained the P396L mutation and contained additional nucleotide changes. The P396L library was screened using two sets of conditions.

[00425] An equilibrium screen was performed using biotinylated FcγRIIA -linker-avitag as a monomer, using methods already described. Approximately 10-fold excess of

library (10^8 cells) was incubated in a 0.5 mL of approximately 7 nM Fc γ RIIIA for 1 hr. The mixture was sorted by FACS, selecting top 1.2% of binders. Selected yeast cells were grown in selective media containing glucose and reinduced in selective media containing galactose. The equilibrium screen was repeated a second time and the sort gate was set to collect the top 0.2% of binders. The selected yeast cells were then grown under selective conditions in glucose. This culture was then used to harvest DNA. Inserts containing the Fc domain were amplified by PCR and cloned into the nucleotide sequence encoding 4-4-20 variable domain using methods already described. Approximately 90 Fc mutants were screened by 4-4-20 ELISA and ADCC and the resultant positive mutants are shown in Table 16.

Table 16: Mutants selected by FACS using an Equilibrium screen with concentrations of Fc γ RIIIA of approximately 7 nM.

Mutant	Amino Acid changes
MgFc43b	K288R, T307A, K344E, P396L
MgFc44	K334N, P396L
MgFc46	P217S, P396L
MgFc47	K210M, P396L
MgFc48	V379M, P396L
MgFc49	K261N, K210M, P396L
MgFc60	P217S, P396L

[00426] A kinetic screen was also implemented to identify mutants with improved K_{off} in binding Fc γ RIIIA. Conditions were established for screening the P396L library using a strain with the P396L Fc variant displayed on the yeast surface. Briefly cells grown under inducing conditions were incubated with 0.1 μ M biotinylated Fc γ RIIIA-linker-avitag monomer for 1 hr. The cells were washed to remove the labeled ligand. Labeled cells were then incubated for different times with 0.1 μ M unlabeled Fc γ RIIIA-linker-avitag monomer, washed and then stained with SA:PE for FACS analysis (FIG. 20). Cells were also stained with goat anti-human Fc to show that the Fc display was maintained during the experiment.

[00427] Based on the competition study it was determined that a 1 minute incubation resulted in approximately 50% loss of cell staining. This time point was chosen for the kinetic screen using the P396L library. Approximately 10-fold excess of library (10^8 cells) was incubated with 0.1 μ M biotinylated Fc γ RIIIA-linker-avitag monomer in a 0.5 mL volume. Cells were washed and then incubated for 1 minute with unlabeled ligand. Subsequently the cells were washed and labeled with SA:PE. The mixture was sorted by FACS, selecting the top 0.3% of binders. Selected yeast cells were grown in selective media containing glucose and reinduced in selective media containing galactose. The kinetic screen was repeated a second time and the sort gate was set to collect the top 0.2%

of binders. The nonselected P396L library was compared to the yeast cells selected for improved binding by FACS (FIG. 21). The histograms show the percentage of cells that are costained with both Fc γ RIIIA /PE and goat anti-human Fc/FITC (upper right).

- [00428] The selected yeast cells from the second sort were then grown under selective conditions in glucose. This culture was then used to harvest DNA. Inserts containing the Fc domain were amplified by PCR and cloned into the nucleotide sequence encoding 4-4-20 variable domain using methods described above. Approximately 90 Fc mutants were screened by 4-4-20 ELISA and ADCC and the resultant positive mutants are shown in Table 17.

Table 17: Mutants selected by FACS using a Kinetic screen using equimolar amounts of unlabeled CD16A for 1 minute.

Mutants	Amino Acid changes
MgFc50	P247S, P396L
MgFc51	Q419H, P396L
MgFc52	V240A, P396L
MgFc53	L410H, P396L
MgFc54	F243L, V305I, A378D, F404S, P396L
MgFc55	R255I, P396L
MgFc57	L242F, P396L
MgFc59	K370E, P396L

Screens 4 and 5: Combining the Solid Phase Fc γ RIIB Depletion Step with Fc γ RIIIA

Selection by FACS Sort, using the Fc γ RIIIA 158V allele

- [00429] Analysis of Fc variants from Screen 1 showed that the mutations that were selected from the secondary screen had improved binding to both Fc γ RIIIA and Fc γ RIIB. Therefore, the data suggested that sequential depletion and selection using magnetic beads (solid phase) under the established conditions did not efficiently select for differential binding of Fc γ RIIIA and Fc γ RIIB. Therefore, in order to screen more effectively for mutants that bind Fc γ RIIIA, while having reduced or no binding to Fc γ RIIB, the solid phase Fc γ RIIB depletion step was combined with Fc γ RIIIA selection by FACS sort. This combination identified Fc variants that bind Fc γ RIIIA with greater or equal affinity than wild-type Fc.
- [00430] A 10-fold excess of the naïve library ($\sim 10^7$) was incubated with magnetic beads coated with Fc γ RIIB. Yeast bound to beads were separated from the non-bound fraction by placing the tube containing the mixture in a magnetic field. Those yeast cells that were not bound to the beads were removed and placed in fresh media and subsequently reinduced in media containing galactose. The Fc γ RIIB depletion by magnetic beads was

repeated 5 times. The resulting yeast population was analyzed and found to show greater than 50% cell staining with goat anti-human Fc and a very small percentage of cells were stained with Fc γ RIIIA. These cells were then selected twice by a FACS sort using 0.1 μ M biotinylated Fc γ RIIIA linker-avitag (data not shown). The Fc γ RIIIA was the 158V allotype.

- 5 Yeast cells were analyzed for both Fc γ RIIIA and Fc γ RIIB binding after each sort and compared to binding by wild-type Fc domain (FIG. 22).

[00431] The selected yeast cells from the second sort were then grown under selective conditions in glucose. This culture was then used to harvest DNA. Inserts containing the Fc domain were amplified by PCR and cloned into the nucleotide sequence encoding 4-4-20 variable domain. Approximately 90 Fc mutants were screened by 4-4-20
10 ELISA and ADCC and the resultant positive mutants are shown in Table 18 (mutants 61-66).

15

Table 18: Mutants selected by magnetic bead depletion using beads coated with CD32B and final selection by FACS using Fc γ RIIIA 158Valine or 158Phenylalanine

Mutants	Amino Acid Changes
MgFc61	A330V
MgFc62	R292G
MgFc63	S298N, K360R, N361D
MgFc64	E233G
MgFc65	N276Y
MgFc66	A330V, V427M
MgFc67	V284M, S298N, K334E, R355W, R416T

- [00432] *Screening of Fc mutants using the 158F allele of Fc γ RIIIA:* Two different
20 alleles of Fc γ RIIIA receptor exist that have different binding affinities for the IgG1 Fc domain (Koene *et al.*, 1997, *Blood* 90: 1109-1114; Wu *et al.*, 1997, *J. Clin. Invest.* 100: 1059-70). The 158F allele binds to the Fc domain with a binding constant 5-10 fold lower than the 158V allele. Previously all of the Fc screens using yeast display were done using the high binding 158V allele as a ligand. In this experiment, Fc mutants were selected from
25 the Fc γ RIIB depleted yeast population using biotinylated Fc γ RIIIA158F-linker-avitag monomer as a ligand. The sort gate was set to select the top 0.25 percent Fc γ RIIIA 158F binders. The resulting enriched population was analyzed by FACS (FIG. 23A). Individual clones were then isolated and their binding to different Fc γ Rs were analyzed by FACS (FIG. 23B). Analysis of individual clones from the population resulted in the identification

of a single mutant harboring 5 mutations MgFc67 (V284M, S298N, K334E, R355W, R416S), which had an enhanced binding to FcγRIIIA and a reduced binding to FcγRIIB.

Secondary Screen of Mutants by an ADCC assay For Screens 1, 2, and 3:

- [00433] Mutants that were selected in the above screens were then analyzed using a standard ADCC assay to determine the relative rates of lysis mediated by ch4-4-20 harboring the Fc mutants. ch4-4-20 antibodies carrying the Fc variants were constructed using methods already described above. SK-BR3 cells were used as targets and effector cells were PBMC that were isolated from donors using a Ficoll gradient, as described supra (Section 6.7). The ADCC activity results for the mutants are summarized in Table 19.
- 10 [00434] As seen in Table 19, mutants isolated using the above primary and secondary screens based on FcγRIIB depletion and FcγRIIIA selection showed enhanced ADCC activity relative to wild-type.

15 **Table 19: Analysis of ADCC mediated by 4-4-20 anti-Fluorescein antibody on SKBR3 cells coated with fluorescein.**

Mutant	Amino Acid Change	Relative rate of lysis
MgFc37	K248M	3.83
MgFc38	K392T, P396L	3.07
MgFc39	E293V, Q295E, A327T	4.29
MgFc41	H268N, P396LN	2.24
MgFc43	Y319F, P352L, P396L	1.09
MgFc42	D221E, D270E, V308A, Q311H, P396L, G402D	3.17
MgFc43b	K288R, T307A, K344E, P396L	3.3
MgFc44	K334N, P396L	2.43
MgFc46	P217S, P396L	2.04
MgFc47	K210M, P396L	2.02
MgFc48	V379M, P396L	2.01
MgFc49	K261N, K210M, P396L	2.06
MgFc50	P247S, P396L	2.1
MgFc51	Q419H, P396L	2.24
MgFc52	V240A, P396L	2.35
MgFc53	L410H, P396L	2
MgFc54	F243L, V305I, A378D, F404S, P396L	3.59
MgFc55	R255I, P396L	2.79
MgFc57	L242F, P396L	2.4
MgFc59	K370E, P396L	2.47
MgFc60	P217S, P396L	1.44

[00435] Mutants 37, 38, 39, 41, 43 were analyzed using 0.5 µg/mL ch4-4-20. All other antibodies were tested at 1 µg/mL. All rates were normalized to wild type ch4-4-20 (IgG1).

[00436] Mutants were additionally cloned into the heavy chain of antitumor
5 monoclonal antibody 4D5 (anti-HER2/neu) and anti-CD20 monoclonal antibody 2H7 by replacing the Fc domain of these monoclonal antibodies. These chimeric monoclonal antibodies were expressed and purified and tested in an ADCC assay using standard methods by transient transfection into 293H cells and purification over protein G column. The chimeric 4D5 antibodies were tested in an ADCC assay using SK-BR3 cells as targets
10 (FIG. 24), whereas the chimeric 2H7 antibodies were tested in an ADCC assay using Daudi cells as targets (FIG. 25).

[00437] *Secondary Screen of Mutants via BIAcore:* Mutants that were selected in the above screens were then analyzed by BIAcore to determine the kinetic parameters for binding FcγRIIIA(158V) and FcγRIIB. The method used was similar to that disclosed in
15 Section 6.8, supra.

[00438] The data displayed are K_{off} values relative to wild type off rates as determined from experiments using the Fc mutants in the ch4-4-20 monoclonal antibody. Relative numbers greater than one indicate a decrease in K_{off} rate. Numbers less than one indicate an increase in off rate.

20 [00439] Mutants that showed a decrease in off rates for FcγRIIIA were MgFc38 (K392, P396L), MgFc43(Y319F, P352L, P396L), MgFc42(D221E, D270E, V308A, Q311H, P396L, G402D), MgFc43b (K288R, T307A, K344E, P396L), MgFc44 (K334N, P396L), MgFc46 (P217S, P396L), MgFc49 (K261N, K210M, P396L). Mutants that showed a decrease in off rate for FcγRIIB were, MgFc38(K392, P396L), MgFc39 (E293V, Q295E,
25 A327T), MgFc43 (K288R, T307A, K344E, P396L), MgFc44 (K334N, P396L). The BIAcore data is summarized in Table 20.

Table 20: BIAcore data.

[00440] The invention described and claimed herein is not to be limited in scope by

Fc mutant	AA residues	FcγRIIIA158V (Koff WT/ Mut)	FcγRIIB (Koff WT/ Mut)
MgFc37	K248M	0.977	1.03
MgFc38	K392T, P396L	1.64	2.3
MgFc39	E293V, Q295E, A327T	0.86	1.3
MgFc41	H268N, P396LN	0.92	1.04
MgFc43	Y319F, P352L, P396L	1.23	2.29
	D221E, D270E, V308A, Q311H, P396L,		
MgFc42	G402D	1.38	
MgFc43b	K288R, T307A, K344E, P396L	1.27	0.89
MgFc44	K334N, P396L	1.27	1.33
MgFc46	P217S, P396L	1.17	0.95
MgFc47	K210M, P396L		
MgFc48	V379M, P396L		
MgFc49	K261N, K210M, P396L	1.29	0.85
MgFc50	P247S, P396L		
MgFc51	Q419H, P396L		
MgFc52	V240A, P396L		
MgFc53	L410H, P396L		
MgFc54	F243L, V305I, A378D, F404S, P396L		
MgFc55	R255I, P396L		
MgFc57	L242F, P396L		
MgFc59	K370E, P396L		
MgFc60	P217S, P396L		
MgFc61	A330V	1	0.61
MgFc62	R292G	1	0.67
MgFc63	S298N, K360R, N361D	1	0.67
MgFc64	E233G	1	0.54
MgFc65	N276Y	1	0.64
MgFc66	A330V, G427M,	1	0.62
MgFc67	V284M, S298N, K334E, R355W, R416S		

the specific embodiments herein disclosed since these embodiments are intended as

illustration of several aspects of the invention. Any equivalent embodiments are intended to
 5 be within the scope of this invention. Indeed, various modifications of the invention in
 addition to those shown and described herein will become apparent to those skilled in the art
 from the foregoing description. Such modifications are also intended to fall within the
 scope of the appended claims.

[00441] Throughout this application various publications are cited. Their contents
 10 are hereby incorporated by reference into the present application in their entireties for all
 purposes

WHAT IS CLAIMED IS:

1. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable
5 polypeptide comprising the wild-type Fc region binds Fc γ R11A, provided that said at least one amino acid modification is not solely a substitution at positions 329, 331, or 332, or does not have an alanine at any of positions 256, 290, 298, 312, 326, 333, 334, 359, 360, or 430; a lysine at position 330; a threonine at position 339; a methionine at position 320; a
10 serine at position 326; an asparagine at position 326; an aspartic acid at position 326; a glutamic acid at position 326; a glutamine at position 334; a glutamic acid at position 334; a methionine at position 334; a histidine at position 334; a valine at position 334; a lysine at position 330; a lysine at position 335; or a leucine at position 334.
2. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
15 said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises a substitution at position 339 with valine and at position 347 with histidine.
3. A polypeptide comprising a variant Fc region, wherein said variant Fc region
20 comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 251 with proline and at position 415 with isoleucine.
- 25 4. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid
30 modification comprises substitution at position 185 with methionine, and at position 218 with asparagine, and at position 292 with leucine, and at position 399 with glutamic acid.
5. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that

said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 290 with proline and at position 142 with proline.

- 5 6. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 141 with valine, at position 268 with
10 leucine, at position 288 with glutamic acid, and at position 291 with serine.
7. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid
15 modification comprises substitution at position 133 with methionine, at position 149 with tyrosine, at position 205 with glutamic acid, at position 334 with asparagine, and at position 384 with lysine.
8. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
20 said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 125 with leucine, at position 215 with isoleucine, and at position 408 with isoleucine.
9. A polypeptide comprising a variant Fc region, wherein said variant Fc region
25 comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 395 with isoleucine.
10. A polypeptide comprising a variant Fc region, wherein said variant Fc region
30 comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable

polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position of 247 with histidine.

11. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
5 said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 396 with histidine.
12. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
10 said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 392 with arginine.
13. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
15 said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 415 with isoleucine and at position 251 with phenylalanine.
14. A polypeptide comprising a variant Fc region, wherein said variant Fc region
20 comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 301 with cysteine, at position 252 with leucine, and at position 192 with threonine.
15. A polypeptide comprising a variant Fc region, wherein said variant Fc region
25 comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 315 with isoleucine.
16. A polypeptide comprising a variant Fc region, wherein said variant Fc region
30 comprises at least one amino acid modification relative to a wild-type Fc region, such that

said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 132 with isoleucine.

17. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 162 with valine.

18. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 348 with methionine, at position 334 with asparagine, at position 275 with isoleucine, at position 202 with methionine, and at position 147 with threonine.

19. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region; such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 310 with tyrosine, at position 289 with alanine, and at position 337 with glutamic acid.

20. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 119 with phenylalanine, at position 371 with serine, at position 407 with valine, and at position 258 with aspartic acid.

21. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid

modification comprises substitution at position 409 with arginine and at position 166 with asparagine.

22. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
5 said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 408 with isoleucine, at position 215 with isoleucine, and at position 125 with isoleucine.
23. A polypeptide comprising a variant Fc region, wherein said variant Fc region
10 comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 396 with leucine.
24. A polypeptide comprising a variant Fc region, wherein said variant Fc region
15 comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 385 with glutamic acid and at position 247 with histidine.
- 20 25. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 379 with methionine.
- 25 26. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 219 with tyrosine.
- 30 27. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that

said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 282 with methionine.

28. A polypeptide comprising a variant Fc region, wherein said variant Fc region
5 comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 276 with isoleucine, at position 334 with asparagine, and at position 348 with methionine.
- 10 29. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 401 with valine.
- 15 30. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid
20 modification comprises substitution at position 280 with leucine and at position 395 with serine.
31. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid
25 modification comprises substitution at position 222 with asparagine.
32. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid
30 modification comprises substitution at position 246 with threonine and at position 319 with phenylalanine.

33. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid
5 modification comprises substitution at position 243 with isoleucine and at position 379 with leucine.
34. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable
10 polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 246 with threonine and at position 396 with histidine.
35. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
15 said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 268 with aspartic acid and at position 318 with aspartic acid.
36. A polypeptide comprising a variant Fc region, wherein said variant Fc region
20 comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 288 with asparagine, at position 330 with serine, and at position 396 with leucine.
- 25 37. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid
30 modification comprises substitution at position 243 with leucine, at position 255 with leucine, and at position 318 with lysine.
38. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that

said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 334 with glutamic acid, at position 359 with asparagine, and at position 366 with serine.

- 5 39. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 377 with phenylalanine.
- 10 40. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 334 with isoleucine.
- 15 41. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 244 with histidine, at position 358 with
- 20 methionine, at position 379 with methionine, at position 384 with lysine, and at position 378 with methionine.
42. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable
- 25 polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 247 with leucine.
43. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable
- 30 polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 217 with serine, at position 378 with valine, and at position 408 with arginine.

44. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid
5 modification comprises substitution at position 247 with leucine, at position 253 with asparagine, and at position 334 with asparagine.

45. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable
10 polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 288 with methionine and at position 334 with glutamic acid.

46. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
15 said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 334 with glutamic acid and at position 380 with aspartic acid.

47. A polypeptide comprising a variant Fc region, wherein said variant Fc region
20 comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 256 with serine, at position 305 with isoleucine, at position 334 with glutamic acid, and at position 390 with serine.

25 48. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 372 with tyrosine.

30 49. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable

polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 246 with isoleucine and at position 334 with asparagine.

50. A polypeptide comprising a variant Fc region, wherein said variant Fc region
5 comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 335 with asparagine, at position 370 with glutamic acid, at position 378 with glutamic acid, at position 394 with methionine, and at
10 position 424 with leucine.

51. A polypeptide comprising a variant Fc region, wherein said variant Fc region
comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable
polypeptide comprising the wild-type Fc region, wherein said at least one amino acid
15 modification comprises substitution at position 320 with glutamic acid and at position 326 with glutamic acid.

52. A polypeptide comprising a variant Fc region, wherein said variant Fc region
comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable
20 polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 224 with leucine.

53. A polypeptide comprising a variant Fc region, wherein said variant Fc region
comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable
25 polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 375 with cysteine and at position 396 with leucine.

54. A polypeptide comprising a variant Fc region, wherein said variant Fc region
comprises at least one amino acid modification relative to a wild-type Fc region, such that
30 said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid

modification comprises substitution at position 233 with aspartic acid and at position 334 with glutamic acid.

55. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
- 5 said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 334 with glutamic acid, at position 359 with asparagine, at position 366 with serine, and at position 386 with arginine.
56. A polypeptide comprising a variant Fc region, wherein said variant Fc region
- 10 comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region binds Fc γ R11A, and said polypeptide further specifically binds Fc γ R11B with a lower affinity than a comparable polypeptide comprising the wild-type Fc region binds Fc γ R11B, provided that said variant Fc region
- 15 does not have an alanine at any of positions 256, 298, 333, or 334.
57. The polypeptide of claim 1, wherein said modification comprises at least one amino acid substitution in the Fc region.
58. The polypeptide of claim 1, wherein said amino acid modification comprises at least one amino acid modification in the CH2 domain of the Fc region.
- 20 59. The polypeptide of claim 58, wherein said amino acid modification comprises substitution of Pro247 with another amino acid at that position.
60. The polypeptide of claim 1, wherein said amino acid modification comprises at least one amino acid modification in the CH3 domain of the Fc region.
61. The polypeptide of claim 60, wherein said amino acid modification comprises
- 25 substitution at position 396 with another amino acid at that position.
62. The polypeptide of claim 1, wherein said amino acid modification comprises at least one amino acid modification in the CH2 domain and at least one amino acid modification in the CH3 domain of the Fc region.

63. The polypeptide of claim 62, wherein said amino acid modification in the CH2 domain comprises substitution at position 251, 292, 268, 288, 291, or 247 with another amino acid at that position.
64. The polypeptide of claim 62, wherein said amino acid modification in the CH3 domain comprises substitution at position 347, 415, 399, 383, 384, 407, 395, or 396 with another amino acid at that position.
65. The polypeptide of claim 64, further comprising at least one amino acid modification in the hinge region of the Fc region.
66. The polypeptide of claim 1, wherein said amino acid modification comprises at least one amino acid modification in the hinge region of the Fc region.
67. The polypeptide of any of claims 1 or 56, wherein the Fc region of the parent polypeptide is a human IgG Fc region.
68. The polypeptide of claim 67, wherein the human IgG Fc region is a human IgG1, IgG2, IgG3, or IgG4 Fc region.
69. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region binds Fc γ RIIIA, provided that said variant at least one amino acid modification is not solely a substitution at positions 329, 331, or 332, or does not have an alanine at any of positions 256, 290, 298, 312, 333, 334, 359, 360, or 430; a lysine at position 330; a threonine at position 339; a methionine at position 320; a serine at position 326; an asparagine at position 326; an aspartic acid at position 326; a glutamic acid at position 326; a glutamine at position 334; a glutamic acid at position 334; a methionine at position 334; a histidine at position 334; a valine at position 334; or a leucine at position 334.
70. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region binds Fc γ RIIIA, and said antibody further specifically binds Fc γ RIIB with a lower affinity than a comparable antibody comprising the wild-type

Fc region binds Fc γ RIIB, provided that said variant Fc region does not have an alanine at any of positions 256, 298, 333, or 334.

71. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
5 said antibody specifically binds Fc γ RIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises a substitution at position 339 with valine and at position 347 with histidine.

72. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
10 said antibody specifically binds Fc γ RIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 251 with proline and at position 415 with isoleucine.

73. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
15 said antibody specifically binds Fc γ RIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 185 with methionine, and at position 218 with asparagine, and at position 292 with leucine, and at position 399 with glutamic acid.

74. An antibody comprising a variant Fc region, wherein said variant Fc region
20 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 290 with proline and at position 142 with proline.

75. An antibody comprising a variant Fc region, wherein said variant Fc region
25 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 141 with valine, at position 268 with leucine, at position 288 with glutamic acid, and at position 291 with serine.

30 76. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that

said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 133 with methionine, at position 149 with tyrosine, at position 205 with glutamic acid, at position 334 with asparagine, and at position 384 with lysine.

77. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position of 247 with histidine.

78. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 396 with histidine.

79. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 392 with arginine.

80. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 301 with cysteine, at position 252 with leucine, and at position 192 with threonine.

81. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 315 with isoleucine.

82. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification
5 comprises substitution at position 132 with isoleucine.

83. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification
10 comprises substitution at position 162 with valine.

84. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification
15 comprises substitution at position 348 with methionine, at position 334 with asparagine, at position 275 with isoleucine, at position 202 with methionine, and at position 147 with threonine.

85. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
20 said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 310 with tyrosine, at position 289 with alanine, and at position 337 with glutamic acid.

86. An antibody comprising a variant Fc region, wherein said variant Fc region
25 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 119 with phenylalanine, at position 371 with serine, at position 407 with valine, and at position 258 with aspartic acid.

30 87. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody

comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 409 with arginine and at position 166 with asparagine.

88. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
5 said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 408 with isoleucine, at position 215 with isoleucine, and at position 125 with isoleucine.

89. An antibody comprising a variant Fc region, wherein said variant Fc region
10 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 396 with leucine.

90. An antibody comprising a variant Fc region, wherein said variant Fc region
15 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 385 with glutamic acid and at position 247 with histidine.

91. An antibody comprising a variant Fc region, wherein said variant Fc region
20 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 379 with methionine.

92. An antibody comprising a variant Fc region, wherein said variant Fc region
25 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 219 with tyrosine.

93. An antibody comprising a variant Fc region, wherein said variant Fc region
30 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody

comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 282 with methionine.

94. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
5 said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 276 with isoleucine, at position 334 with asparagine, and at position 348 with methionine.

95. An antibody comprising a variant Fc region, wherein said variant Fc region
10 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 401 with valine.

96. An antibody comprising a variant Fc region, wherein said variant Fc region
15 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 280 with leucine and at position 395 with serine.

97. An antibody comprising a variant Fc region, wherein said variant Fc region
20 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 222 with asparagine.

98. An antibody comprising a variant Fc region, wherein said variant Fc region
25 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 246 with threonine and at position 319 with phenylalanine.

30 99. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that

said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 243 with isoleucine and at position 379 with leucine.

100. An antibody comprising a variant Fc region, wherein said variant Fc region
5 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 334 with glutamic acid.

101. An antibody comprising a variant Fc region, wherein said variant Fc region
10 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 246 with threonine and at position 396 with histidine.

102. An antibody comprising a variant Fc region, wherein said variant Fc region
15 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 268 with aspartic acid and at position 318 with aspartic acid.

20 103. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 288 with asparagine, at position 330 with serine, and at
25 position 396 with leucine.

104. An antibody comprising a variant Fc region, wherein said variant Fc region
comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification
30 comprises substitution at position 243 with leucine, at position 255 with leucine, and at position 318 with lysine.

105. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification
5 comprises substitution at position 334 with glutamic acid, at position 359 with asparagine, and at position 366 with serine.

106. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody
10 comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 377 with phenylalanine.

107. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody
15 comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 334 with isoleucine.

108. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody
20 comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 244 with histidine, at position 358 with methionine, at position 379 with methionine, at position 384 with lysine, and at position 397 with methionine.

109. An antibody comprising a variant Fc region, wherein said variant Fc region
25 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 247 with leucine.

110. An antibody comprising a variant Fc region, wherein said variant Fc region
30 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification

comprises substitution at position 217 with serine, at position 378 with valine, and at position 408 with arginine.

111. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 247 with leucine, at position 253 with asparagine, and at position 334 with asparagine.

112. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 288 with methionine and at position 334 with glutamic acid.

113. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 334 with glutamic acid and at position 380 with aspartic acid.

114. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 256 with serine, at position 305 with isoleucine, at position 334 with glutamic acid, and at position 390 with serine.

115. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 326 with glutamic acid.

116. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification
5 comprises substitution at position 372 with tyrosine.

117. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification
10 comprises substitution at position 246 with isoleucine and at position 334 with asparagine.

118. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification
15 comprises substitution at position 335 with asparagine, at position 370 with glutamic acid, at position 378 with valine, at position 394 with methionine, and at position 424 with leucine.

119. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
20 said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 320 with glutamic acid and at position 326 with glutamic acid.

120. An antibody comprising a variant Fc region, wherein said variant Fc region
25 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 224 with leucine.

121. An antibody comprising a variant Fc region, wherein said variant Fc region
30 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody

comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 375 with cysteine and at position 396 with leucine.

122. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
5 said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 233 with aspartic acid and at position 334 with glutamic acid.

123. An antibody comprising a variant Fc region, wherein said variant Fc region
10 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 334 with glutamic acid, at position 359 with asparagine, at position 366 with serine, and at position 386 with arginine.

15 124. The antibody of claims 69 or 70, wherein said antibody is a monoclonal antibody.

125. The antibody of claim 69 or 70, wherein said antibody is a humanized antibody.

126. The antibody of claim 69 or 70, wherein said antibody is a human antibody.

127. A nucleic acid comprising a nucleotide sequence encoding the polypeptide of any of claims 1 or 56.

20 128. A vector comprising the nucleic acid of claim 127.

129. The vector of claim 128 which is an expression vector.

130. A host cell comprising the nucleic acid of claim 127.

131. A method for recombinantly producing the polypeptide of claims 1 or 56, said method comprising: (i) culturing in a medium a host cell comprising a nucleic acid
25 encoding said polypeptide, under conditions suitable for the expression of said polypeptide; and (ii) recovery of said polypeptide from said medium.

132. The antibody of claims 69 or 70, wherein said antibody specifically binds Fc γ RIIIA with at least two times greater affinity than a comparable antibody comprising the wild-type Fc region binds Fc γ RIIIA.

133. A therapeutic antibody that specifically binds a cancer antigen, said therapeutic antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said therapeutic antibody specifically binds Fc γ RIIIA with a greater affinity than the therapeutic antibody comprising the wild-type Fc region binds Fc γ RIIIA, provided that said variant Fc region does not have a substitution at positions 329, 331, or 332, and does not have an alanine at any of positions 256, 290, 298, 312, 333, 334, 359, 360, or 430; a lysine at position 330; a threonine at position 339; a methionine at position 320; a serine at position 326; an asparagine at position 326; an aspartic acid at position 326; a glutamic acid at position 326; a glutamine at position 334; a glutamic acid at position 334; a methionine at position 334; a histidine at position 334; a valine at position 334; or a leucine at position 334.

134. A therapeutic antibody that specifically binds a cancer antigen, said therapeutic antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said therapeutic antibody specifically binds Fc γ RIIIA with a greater affinity than a therapeutic antibody comprising the wild-type Fc region binds Fc γ RIIIA, and said therapeutic antibody further specifically binds Fc γ RIIB with a lower affinity than a therapeutic antibody comprising the wild-type Fc region binds Fc γ RIIB, provided that said variant Fc region does not have an alanine at any of positions 256, 298, 333, or 334.

135. The therapeutic antibody of claim 133 or 134, wherein said therapeutic antibody mediates antibody-dependent cell mediated cytotoxicity 2-fold more effectively, than the therapeutic antibody comprising a wild-type Fc region.

136. The therapeutic antibody of claim 133 or 134, wherein said therapeutic antibody is Herceptin®, Rituxan®, IC14, PANOREX™, IMC-225, VITAXIN™, Campath 1H/LDP-03, LYMPHOCIDE™, or ZEVLIN™.

137. The therapeutic antibody of claim 133 or 134, wherein said cancer antigen is MAGE-1, MAGE-3, BAGE, GAGE-1, GAGE-2, N-acetylglucosaminyltransferase, p15, beta-catenin, MUM-1, CDK4, HER-2/neu, human papillomavirus-E6, human papillomavirus-E7, or MUC-1.

138. A method of treating cancer in a patient having a cancer characterized by a cancer antigen, said method comprising administering to said patient a therapeutically effective amount of a therapeutic antibody of claim 133 or 134.
139. The method of claim 138, wherein said cancer antigen is MAGE-1, MAGE-3,
5 BAGE, GAGE-1, GAGE-2, N-acetylglucosaminyltransferase, p15, beta-catenin, MUM-1, CDK4, HER-2/neu, human papillomavirus-E6, human papillomavirus-E7, or MUC-1.
140. The method of claim 138, wherein said cancer antigen is a breast, ovarian, prostate, cervical, or pancreatic carcinoma antigen.
141. The method of claim 138 further comprising the administration of one or more
10 additional cancer therapies.
142. The method of claim 141, wherein said additional cancer therapy is selected from the group consisting of chemotherapy, immunotherapy, radiation therapy, hormonal therapy, or surgery.
143. The method of claim 138, wherein said patient is human.
- 15 144. A pharmaceutical composition comprising a therapeutically effective amount of one or more of the polypeptides of claims 1 or 56, and a pharmaceutically acceptable carrier.
145. A pharmaceutical composition comprising a therapeutically effective amount of one or more of the antibodies of claims 69 or 70, and a pharmaceutically acceptable carrier.
146. A pharmaceutical composition comprising a therapeutically effective amount of one
20 or more of the therapeutic antibodies of claims 133-137, and a pharmaceutically acceptable carrier.
147. The pharmaceutical composition of claims 133-137, further comprising one or more additional anti-cancer agents.
148. The pharmaceutical compositions of claim 147, wherein said anti-cancer agents is a
25 chemotherapeutic agent, a radiation therapeutic agent, a hormonal therapeutic agent, or an immunotherapeutic agent.
149. A nucleic acid comprising a nucleotide sequence encoding a heavy chain of an antibody of any of claims 69 or 70.

150. A vector comprising the nucleic acid of claim 149.
151. The vector of claim 150 which is an expression vector.
152. A host cell comprising the nucleic acid of claim 149.
153. A method for recombinantly producing the antibody of claim 69 or 70, said method
5 comprising: (i) culturing in a medium a host cell comprising a nucleic acid encoding said
antibody, under conditions suitable for the expression of said antibody; and (ii) recovery of
said antibody from said medium.
154. A method for producing a tetrameric Fc γ R complex, wherein said tetrameric
complex has an enhanced affinity for an Fc region, relative to the affinity of a monomeric
10 Fc γ R for the Fc region, comprising:
- (a) producing a fusion protein, such that a 15 amino acid AVITAG sequence is
operably linked to the soluble region of Fc γ R;
 - (b) biotinylating the protein produced in step (a);
 - (c) mixing the biotinylated protein produced in step (b) with streptaividn-
15 phycoerythin in an appropriate molar ratio,
- such that a tetrameric Fc γ R complex is formed.
155. The tetrameric Fc γ R complex produced by the method of claim 154.
156. The tetrameric Fc γ R complex of claim 155, wherein said complex binds an Fc
region with an 8-fold higher affinity than a monomeric Fc γ R complex binds the Fc region.
- 20 157. The tetrameric Fc γ R complex of claim 155, wherein said complex binds an Fc
region with a 10-fold higher affinity than a monomeric Fc γ R complex binds the Fc region.
158. The method of claim 154, wherein said tetrameric Fc γ R complex is a tetrameric
Fc γ RIIA complex, and said soluble region used in step (a) is the soluble region of
Fc γ RIIA.
- 25 159. The method of claim 154, wherein said tetrameric Fc γ R complex is a tetrameric
Fc γ RIIB complex, and said soluble region used in step (a) is the soluble region of Fc γ RIIB.

160. The method of claim 154, wherein the protein produced in step (a) is biotinylated enzymatically.
161. The method of claim 154, wherein the protein produced in step (a) is biotinylated with *E. coli* Bir A enzyme.
- 5 162. The method of claim 154, wherein the biotinylated protein produced in step (a) is mixed with streptavidin-phycoerythrin in a 1:5 molar ratio (1X streptavidin-phycoerythrin : 5X biotinylated protein).
163. A method of treating or managing cancer in a patient having a cancer characterized by a cancer antigen, said method comprising administering to said patient a therapeutically effective amount of one or more of the antibodies of claims 69, 70, 133, or 134.
- 10 164. A method of treating cancer in a patient having a cancer characterized by a cancer antigen, said method comprising administering to said patient a therapeutically effective amount of one or more of the antibodies of claim 69, 70, 133, or 134 that specifically bind said cancer antigen.
- 15 165. A method of treating an autoimmune disorder in a patient in need thereof, said method comprising administering to said patient a therapeutically effective amount of a molecule comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild type Fc region, such that said molecule specifically binds Fc γ RIIB with a greater affinity than a comparable molecule comprising the wild type Fc region, and said molecule further specifically binds Fc γ RIIA with a lower affinity than a comparable molecule comprising the wild type Fc region.
- 20 166. The method of claim 165, wherein said autoimmune disorder is rheumatoid arthritis, psoriatic arthritis, ankylosing spondylitis, Rieter's Syndrome, psoriasis, or lupus erythematosus.
- 25 167. The method of claim 165 further comprising administering to said patient a therapeutically effective amount one or more anti-inflammatory agents.
168. The method of claim 165 further comprising administering to said patient a therapeutically effective amount one or more immunomodulatory agents.

169. The method of claim 168, wherein at least one immunomodulatory agent is a small organic molecule.
170. The method of claim 169, wherein the small organic molecule is methotrexate, leflunomide, cyclophosphamide, cyclosporin A, FK506, mycophenolate mofetil, rapamycin,
5 mizoribine, deoxyspergualin, brequinar, malonitrolamide, steroid, or corticosteroid.
171. The method of claim 167, wherein at least one anti-inflammatory agents is a non-steroidal anti-inflammatory drug.
172. The method of claim 171, wherein the non-steroidal anti-inflammatory drug is aspirin, ibuprofen, diclofenac, nabumetone, naproxen, or ketoprofen.
- 10 173. A method of treating an infectious disease in a patient in need thereof, said method comprising administering to said patient a therapeutically effective amount of a molecule comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild type Fc region, such that said molecule specifically binds Fc γ R1IB with a greater affinity than a comparable molecule comprising the wild type
15 Fc region, and said molecule further specifically binds Fc γ R1IIIA with a lower affinity than a comparable molecule comprising the wild type Fc region.
174. The polypeptide of claims 1 or 56, wherein said polypeptide specifically binds Fc γ R1IIIA with a 4-fold greater affinity than a comparable polypeptide comprising the wild type Fc region binds Fc γ R1IIIA.
- 20 175. The polypeptide of claims 1 or 56, wherein said polypeptide specifically binds Fc γ R1IIIA with an 8-fold greater affinity than a comparable polypeptide comprising the wild type Fc region binds Fc γ R1IIIA
176. The polypeptide of claims 1 or 56, wherein said polypeptide specifically binds Fc γ R1IIIA with a 10-fold greater affinity than a comparable polypeptide comprising the wild
25 type Fc region binds Fc γ R1IIIA
177. The polypeptide of claims 1 or 56, wherein said polypeptide specifically binds Fc γ R1IIIA with a 100-fold greater affinity than a comparable polypeptide comprising the wild type Fc region binds Fc γ R1IIIA

178. The antibody of claim 69 or 70, wherein said antibody specifically binds Fc γ RIIIA with a 4-fold greater affinity than a comparable antibody comprising the wild type Fc region binds Fc γ RIIIA.
179. The antibody of claim 69 or 70, wherein said antibody specifically binds Fc γ RIIIA with an 8-fold greater affinity than a comparable antibody comprising the wild type Fc binds Fc γ RIIIA.
180. The antibody of claim 69 or 70, wherein said antibody specifically binds Fc γ RIIIA with a 10-fold greater affinity than a comparable antibody comprising the wild type Fc binds Fc γ RIIIA.
181. The antibody of claim 69 or 70, wherein said antibody specifically binds Fc γ RIIIA with a 100-fold greater affinity than a comparable antibody comprising the wild type Fc binds Fc γ RIIIA.
182. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIIA with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 247 with leucine and at position 420 with valine.
183. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIIA with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution 334 with glutamic acid and at position 292 with leucine.
184. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIIA with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, and said polypeptide further specifically binds FcRIIB with a lower affinity than a comparable polypeptide comprising the wild type Fc region binds FcRIIB, wherein said at least one amino acid modification comprises substitution at position 243 with isoleucine and at position 379 with leucine.

185. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, and said polypeptide further specifically
5 binds FcR11B with a lower affinity than a comparable polypeptide comprising the wild type Fc region binds FcR11B, wherein said at least one amino acid modification comprises substitution at position 288 with asparagine, at position 330 with serine and at position 396 with leucine.

186. A polypeptide comprising a variant Fc region, wherein said variant Fc region
10 comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, and said polypeptide further specifically binds FcR11B with a lower affinity than a comparable polypeptide comprising the wild type Fc region binds FcR11B, wherein said at least one amino acid modification comprises
15 substitution at position 243 with leucine and at position 255 with leucine.

187. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, and said polypeptide further specifically
20 binds FcR11B with a lower affinity than a comparable polypeptide comprising the wild type Fc region binds FcR11B, wherein said at least one amino acid modification comprises substitution at position 288 with methionine and at position 334 with glutamic acid.

188. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
25 said polypeptide specifically binds Fc γ R11A with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, and said polypeptide further specifically binds FcR11B with a lower affinity than a comparable polypeptide comprising the wild type Fc region binds FcR11B, wherein said at least one amino acid modification comprises substitution at position 316 with aspartic acid, at position 378 with valine, and at position
30 399 with glutamic acid.

189. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that

said polypeptide specifically binds Fc γ RIIIA with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, and said polypeptide further specifically binds FcRIIB with a lower affinity than a comparable polypeptide comprising the wild type Fc region binds FcRIIB, wherein said at least one amino acid modification comprises substitution at position 315 with isoleucine, at position 379 with methionine, and at position 399 with glutamic acid.

190. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIIA with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, and said polypeptide further specifically binds FcRIIB with a lower affinity than a comparable polypeptide comprising the wild type Fc region binds FcRIIB, wherein said at least one amino acid modification comprises substitution at position 243 with isoleucine, at position 379 with leucine, and at position 420 with valine.

191. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIIA with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, and said polypeptide further specifically binds FcRIIB with a lower affinity than a comparable polypeptide comprising the wild type Fc region binds FcRIIB, wherein said at least one amino acid modification comprises substitution at position 392 with threonine and at position 396 with leucine.

192. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIIA with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, and said polypeptide further specifically binds Fc γ RIIB with a lower affinity than a comparable polypeptide comprising the wild type Fc region binds Fc γ RIIB, wherein said at least one amino acid modification comprises substitution at position 293 with valine, at position 295 with glutamic acid, and at position 327 with threonine.

193. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds Fc γ RIIIA with a greater affinity than a comparable

polypeptide comprising the wild-type Fc region, and said polypeptide further specifically binds FcRIIB with a lower affinity than a comparable polypeptide comprising the wild type Fc region binds FcRIIB, wherein said at least one amino acid modification comprises substitution at position 268 with asparagine and at position 396 with leucine.

- 5 194. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said polypeptide specifically binds FcγRIIIA with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, and said polypeptide further specifically binds FcRIIB with a lower affinity than a comparable polypeptide comprising the wild type
10 Fc region binds FcRIIB, wherein said at least one amino acid modification comprises substitution at position 319 with phenylalanine, at position 352 with leucine, and at position 396 with leucine.

195. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
15 said polypeptide specifically binds FcγRIIIA with a greater affinity than a comparable polypeptide comprising the wild-type Fc region, and said polypeptide further specifically binds FcRIIB with a lower affinity than a comparable polypeptide comprising the wild type Fc region binds FcRIIB, wherein said at least one amino acid modification comprises substitution at position 248 with methionine.

- 20 196. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds FcγRIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution at position 247 with leucine and at position 420 with valine.

- 25 197. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds FcγRIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, wherein said at least one amino acid modification comprises substitution 334 with glutamic acid and at position 292 with leucine.

- 30 198. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds FcγRIIIA with a greater affinity than a comparable antibody

comprising the wild-type Fc region, and said antibody further specifically binds Fc γ RIIB with a lower affinity than a comparable antibody comprising the wild type Fc region binds Fc γ RIIB, wherein said at least one amino acid modification comprises substitution at position 243 with isoleucine and at position 379 with leucine.

- 5 199. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, and said antibody further specifically binds Fc γ RIIB with a lower affinity than a comparable antibody comprising the wild type Fc region binds
10 Fc γ RIIB, wherein said at least one amino acid modification comprises substitution at position 288 with asparagine, at position 330 with serine and at position 396 with leucine.

200. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIA with a greater affinity than a comparable antibody
15 comprising the wild-type Fc region, and said antibody further specifically binds Fc γ RIIB with a lower affinity than a comparable antibody comprising the wild type Fc region binds Fc γ RIIB, wherein said at least one amino acid modification comprises substitution at position 243 with leucine and at position 255 with leucine.

201. An antibody comprising a variant Fc region, wherein said variant Fc region
20 comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, and said antibody further specifically binds Fc γ RIIB with a lower affinity than a comparable antibody comprising the wild type Fc region binds Fc γ RIIB, wherein said at least one amino acid modification comprises substitution at
25 position 288 with methionine and at position 334 with glutamic acid.

202. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, and said antibody further specifically binds Fc γ RIIB
30 with a lower affinity than a comparable antibody comprising the wild type Fc region binds Fc γ RIIB, wherein said at least one amino acid modification comprises substitution at

position 316 with aspartic acid, at position 378 with valine, and at position 399 with glutamic acid.

203. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, and said antibody further specifically binds Fc γ R11B with a lower affinity than a comparable antibody comprising the wild type Fc region binds Fc γ R11B, wherein said at least one amino acid modification comprises substitution at position 315 with isoleucine, at position 379 with methionine, and at position 399 with glutamic acid.

204. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, and said antibody further specifically binds Fc γ R11B with a lower affinity than a comparable antibody comprising the wild type Fc region binds Fc γ R11B, wherein said at least one amino acid modification comprises substitution at position 243 with isoleucine, at position 379 with leucine, and at position 420 with valine.

205. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, and said antibody further specifically binds Fc γ R11B with a lower affinity than a comparable antibody comprising the wild type Fc region binds Fc γ R11B, wherein said at least one amino acid modification comprises substitution at position 392 with threonine and at position 396 with leucine.

206. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ R11A with a greater affinity than a comparable antibody comprising the wild-type Fc region, and said antibody further specifically binds Fc γ R11B with a lower affinity than a comparable antibody comprising the wild type Fc region binds Fc γ R11B, wherein said at least one amino acid modification comprises substitution at position 293 with valine, at position 295 with glutamic acid, and at position 327 with threonine.

207. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, and said antibody further specifically binds Fc γ RIIB with a lower affinity than a comparable antibody comprising the wild type Fc region binds Fc γ RIIB, wherein said at least one amino acid modification comprises substitution at position 268 with asparagine and at position 396 with leucine.

208. An antibody e comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, and said antibody further specifically binds Fc γ RIIB with a lower affinity than a comparable antibody comprising the wild type Fc region binds Fc γ RIIB, wherein said at least one amino acid modification comprises substitution at position 319 with phenylalanine, at position 352 with leucine, and at position 396 with leucine.

209. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that said antibody specifically binds Fc γ RIIIA with a greater affinity than a comparable antibody comprising the wild-type Fc region, and said antibody further specifically binds Fc γ RIIB with a lower affinity than a comparable antibody comprising the wild type Fc region binds Fc γ RIIB, wherein said at least one amino acid modification comprises substitution at position 248 with methionine.

210. The antibody of any of claims 196-209, wherein said antibody has an enhanced ADCC activity relative to a comparable antibody comprising the wild type fc region.

211. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification, wherein said at least one amino acid modification comprises substitution at position 255 with leucine and at position 396 with leucine.

212. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification, wherein said at least one amino acid

modification comprises substitution at position 370 with glutamic acid and at position 396 with leucine.

213. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification, wherein said at least one amino acid
5 modification comprises substitution at position 392 with threonine and at position 396 with leucine.

214. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification, wherein said at least one amino acid
10 modification comprises substitution at position 221 with glutamic acid, at position 270 with glutamic acid, at position 308 with alanine, at position 311 with histidine, at position 396 with leucine, and at position 402 with aspartic acid.

215. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification, wherein said at least one amino acid
15 modification comprises substitution at position 243 with leucine, at position 305 with isoleucine, at position 378 with aspartic acid, at position 404 with serine, and at position 396 with leucine.

216. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification, wherein said at least one amino acid
20 modification comprises substitution at position 284 with methionine, at position 298 with asparagine, at position 334 with glutamic acid, at position 355 with tryptophan, and at position 416 with threonine.

217. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification, wherein said at least one amino acid
25 modification comprises substitution at position 255 with leucine and at position 396 with leucine.

218. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification, wherein said at least one amino acid
modification comprises substitution at position 370 with glutamic acid and at position 396 with leucine.

30 219. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification, wherein said at least one amino acid

modification comprises substitution at position 392 with threonine and at position 396 with leucine.

220. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification, wherein said at least one amino acid
5 modification comprises substitution at position 221 with glutamic acid, at position 270 with glutamic acid, at position 308 with alanine, at position 311 with histidine, at position 396 with leucine, and at position 402 with aspartic acid.

221. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification, wherein said at least one amino acid
10 modification comprises substitution at position 243 with leucine, at position 305 with isoleucine, at position 378 with aspartic acid, at position 404 with serine, and at position 396 with leucine.

222. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification, wherein said at least one amino acid
15 modification comprises substitution at position 284 with methionine, at position 298 with asparagine, at position 334 with glutamic acid, at position 355 with tryptophan, and at position 416 with threonine.

223. A polypeptide comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
20 said polypeptide binds an FcγR with an altered affinity relative to a polypeptide comprising a wild-type Fc region, provided that said at least one amino acid modification is not solely a substitution at any of positions 255, 258, 267, 269, 270, 276, 278, 280, 283, 285, 289, 292, 293, 294, 295, 296, 300, 303, 305, 307, 309, 322, 329, 332, 331, 337, 338, 340, 373, 376, 416, 419, 434, 435, 437, 438, 439 or does not have an alanine at any of positions 256, 290,
25 298, 312, 333, 334, 359, 360, 326, or 430; a lysine at position 330; a threonine at position 339; a methionine at position 320; a serine at position 326; an asparagine at position 326; an aspartic acid at position 326; a glutamic acid at position 326; a glutamine at position 334; a glutamic acid at position 334; a methionine at position 334; a histidine at position 334; a valine at position 334; or a leucine at position 334; a lysine at position 335 an asparagine at
30 position 268; a glutamine at position 272; a glutamine, serine, or aspartic acid at position 286; a serine at position 290; a methionine, glutamine, glutamic acid, or arginine at position

320; a glutamic acid at position 322; a serine, glutamic acid, or aspartic acid at position 326; a lysine at position 330; a glutamine at position 335; or a methionine at position 301.

224. An antibody comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, such that
- 5 said antibody binds an FcγR with an altered affinity relative to an antibody comprising a wild-type Fc region, provided that said at least one amino acid modification is not solely a substitution at any of positions 255, 258, 267, 269, 270, 276, 278, 280, 283, 285, 289, 292, 293, 294, 295, 296, 300, 303, 305, 307, 309, 322, 329, 332, 331, 337, 338, 340, 373, 376, 416, 419, 434, 435, 437, 438, 439 or does not have an alanine at any of positions 256, 290,
- 10 298, 312, 333, 334, 359, 360, 326, or 430; a lysine at position 330; a threonine at position 339; a methionine at position 320; a serine at position 326; an asparagine at position 326; an aspartic acid at position 326; a glutamic acid at position 326; a glutamine at position 334; a glutamic acid at position 334; a methionine at position 334; a histidine at position 334; a valine at position 334; or a leucine at position 334; a lysine at position 335 an asparagine at
- 15 position 268; a glutamine at position 272; a glutamine, serine, or aspartic acid at position 286; a serine at position 290; a methionine, glutamine, glutamic acid, or arginine at position 320; a glutamic acid at position 322; a serine, glutamic acid, or aspartic acid at position 326; a lysine at position 330; a glutamine at position 335; or a methionine at position 301

1 2 3 4




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19

FACS analysis of FcR3A competition




FIG. 20




FIG. 21





FIG. 22

A.



B.




FIG. 23




FIG. 24




FIG. 25

SEQUENCE LISTING

<110> MacroGenics, Inc.

<120> IDENTIFICATION AND ENGINEERING OF
ANTIBODIES WITH VARIANT Fc REGIONS AND METHODS OF USING SAME

<130> 11183-004-228

<140> to be assigned

<141>

<150> 60/439,498

<151> 2003-01-09

<150> 60/456,041

<151> 2003-03-19

<150> 60/514,549

<151> 2003-10-23

<160> 10

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 86

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer: 5' linker.avitag

<400> 1

ggccgcaggt ggtggtggtt ctggtggtgg tggttctggt ctgaacgaca tcttcgaggc 60
tcagaaaatc gaatggcacg aatgat 86

<210> 2

<211> 86

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer: 3' linker.avitag

<400> 2

ctagatcatt cgtgccattc gattttctga gctcgaaga tgcgttcag accagaacca 60
ccaccaccag aaccaccacc acctgc 86

<210> 3

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer: FcR3A left

<400> 3

gttgatcct ccaactgctc tgctacttct agttt

35

<210> 4
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer: FcR3A Right

<400> 4
gaaaagctta aagaatgatg agatggttga cact

34

<210> 5
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer: FcR2B right

<400> 5
gaagtcgaca atgatcccca ttggtgaaga g

31

<210> 6
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer: FcR2B left

<400> 6
gttagatott gctgtgctat tcctggctcc

30

<210> 7
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer: IgG1 right

<400> 7
atagtcgacc actgatttac ccggaga

27

<210> 8
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer: IgG1 left

<400> 8
ggaattcaac accaaggtgg acaagaaagt t

31

<210> 9
<211> 31
<212> DNA
<213> Artificial Sequence

<220>

<223> Primer: mcr025;chl (f')

<400> 9

aaaggatccg cgagctcagc ctccaccaag g

31

<210> 10

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer: H021

<400> 10

gtctgctcga agcattaacc

20

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
29 July 2004 (29.07.2004)

PCT

(10) International Publication Number
WO 2004/063351 A3

(51) International Patent Classification⁷: **A61K 39/395**,
C07K 16/28

(21) International Application Number:
PCT/US2004/000643

(22) International Filing Date: 9 January 2004 (09.01.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/439,498 9 January 2003 (09.01.2003) US
60/456,041 19 March 2003 (19.03.2003) US
60/514,549 23 October 2003 (23.10.2003) US

(71) Applicant (for all designated States except US): **MACRO-
GENICS, INC.** [US/US]; 1500 East Gude Drive,
Rockville, MD 20850-5307 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **STAVENHAGEN,
Jeffrey** [US/US]; 19417 Treadway Road, Brookville, MD
20833 (US). **VIJH, Sujata** [US/US]; 912-100 Orchard
Ridge Drive, Gaithersburg, MD 20878 (US).

(74) Agents: **INSOGNA, Anthony, M. et al.**; Jones Day, 222
East 41st Street, New York, NY 10017-6702 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,
GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(88) Date of publication of the International search report:
31 March 2005

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: IDENTIFICATION AND ENGINEERING OF ANTIBODIES WITH VARIANT Fc REGIONS AND METHODS OF
USING SAME

(57) Abstract: The present invention relates to molecules, particularly polypeptides, more particularly immunoglobulins (e.g., antibodies), comprising a variant Fc region, wherein said variant Fc region comprises at least one amino acid modification relative to a wild-type Fc region, which variant Fc region binds FcγRIIIA and/or FcγRIIA with a greater affinity, relative to a comparable molecule comprising the wild-type Fc region. The molecules of the invention are particularly useful in preventing, treating, or ameliorating one or more symptoms associated with a disease, disorder, or infection. The molecules of the invention are particularly useful for the treatment or prevention of a disease or disorder where an enhanced efficacy of effector cell function (e.g., ADCC) mediated by FcγR is desired, e.g., cancer, infectious disease, and in enhancing the therapeutic efficacy of therapeutic antibodies the effect of which is mediated by ADCC.

WO 2004/063351 A3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US04/00643

A. CLASSIFICATION OF SUBJECT MATTER IPC(7) : A61K 39/395 C07K 16/28 US CL : 530/387.3, 388.22, 424/133.1, 143.1 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 530/387.3, 388.22, 424/133.1, 143.1 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Please See Continuation Sheet		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	US 2004/0002587 A1 (WATKINS et al) 01 January 2004 (01.01.2004), see pages 2, 21-22.	1, 15, 40, 57-58, 60
P,Y	US 2004/0185045 A1 (KOENIG et al) 23 September 2004 (23.09.2004), see pages 3-8.	165-173
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 05 January 2005 (05.01.2005)		Date of mailing of the international search report 02 FEB 2005
Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US Commissioner for Patents P.O. Box 1450 Alexandria, Virginia 22313-1450 Facsimile No. (703)305-3230		Authorized officer <i>David J Blanchard</i> David J Blanchard Telephone No. (571) 272-0827

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US04/00643

Continuation of B. FIELDS SEARCHED Item 3:

WEST, Medline, EMBASE, Cancerlit, CAPLUS, Biotechno, Biosis.

Search terms:Fc variant/mutant, Fc gammaRIIIA, Fc gammaRIIB, CH2, CH3, autoimmune, infectious disease, cancer or tumor, Fc residues.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US04/00643

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 146-148
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)